

IN FOREST PLANTINGS:

Sprayer Calibration for Herbicide Application

Extension Bulletin E-916

Reprinted June 1985

by Randall Heiligmann, Extension Specialist in Forestry

Control of competing vegetation at the time of planting and for several years following planting is often essential for good survival and growth of the trees. The most economical and effective method of accomplishing this is with herbicides. Many of the herbicides used in forestry are applied with a sprayer, and their accurate application at prescribed rates requires calibrating the sprayer. If the sprayer is not accurately calibrated, too little, or too much, herbicide may be applied resulting in unsatisfactory weed control or damage or death of the seedlings.

Herbicides are generally prescribed in pounds or quarts per acre. A recommendation might, for example, be 3 lbs./A (acre) or 1/2 qt./A. Sprayer calibration involves determining how much herbicide and water should be mixed in the sprayer to deliver the herbicide at the desired rate, whether the area sprayed is 1 acre or 1/1000 acre.

The first step in calibrating a sprayer is to determine the volume of spray delivered by your sprayer to a known area. This may be done in one of two ways:

1. Fill the sprayer with a known volume of water, spray a known area of ground as if you were applying herbicide, and measure the amount of water needed to refill the sprayer. For example, a small compression tank sprayer might use 1 gallon (128 fluid ounces) to spray an area 4 ft. wide and 85 ft. long (340 sq. ft. or .0078/A).*

2. Fill the sprayer with a known amount of water, spray as if you were applying herbicide until the tank is empty, and then measure the square foot of area sprayed.

Make this determination several times and use the average value. The amount of herbicide to add to a particular volume of water can now be calculated as follows:

$$\begin{array}{l} \text{Weight of} \\ \text{Herbicide} \\ \text{Needed} \end{array} = \begin{array}{l} \text{Recommended} \\ \text{Herbicide Rate} \\ \text{in Pounds/Acre} \end{array} \times \begin{array}{l} \text{Acreage Covered} \\ \text{with Known Volume} \\ \text{of Water} \end{array}$$

Using our example sprayer, which sprays .0078/A with 1 gallon and a recommended herbicide application rate of 2 lbs./A (32 ounces/A), the calculation would be as follows:

$$\begin{array}{l} \text{Weight of} \\ \text{Herbicide} \\ \text{Needed} \end{array} = (32 \text{ ounces/A}) \times (.0078 \text{ A}) = .25 \text{ ounces}$$

To apply herbicide with our sprayer at a rate of 2 pounds acre, 1/4 ounce of herbicide should be added to each gallon of water. The amount of herbicide to mix with a full sprayer of water is then calculated as follows:

$$\begin{array}{l} \text{Weight of} \\ \text{Herbicide to} \\ \text{Add to Full} \\ \text{Sprayer} \end{array} = \begin{array}{l} \text{Weight of Herbicide} \\ \text{to Add to Known} \\ \text{Volume of Water} \end{array} \times \frac{\text{Volume of Sprayer}}{\text{Known Volume of Water}}$$

In our example, if our sprayer's total capacity was 2 1/2 gallons:

$$\begin{array}{l} \text{Weight of} \\ \text{Herbicide to} \\ \text{Add to Full} \\ \text{Sprayer} \end{array} = (\frac{1}{4} \text{ ounce}) \times \frac{2\frac{1}{2} \text{ gallons}}{1 \text{ gallon}} = .625 \text{ ounces}$$

Herbicide recommendations are often made in pounds of active ingredients per acre (lbs. a.i./A) instead of the usual pounds of total material per acre (lbs./A). When this is the case, the amount of herbicide needed is determined by dividing the "Weight of Herbicide Needed," as calculated above, by the percent active ingredients in the herbicide.** Using our example above, if the recommendation had been for 2 lbs. a.i./A instead of 2 lbs./A, and the herbicide had been 80% active ingredient, we would have needed to add 0.31 ounces to each gallon of water or 0.78 ounces to our full sprayer to obtain the recommended rate of 2 lbs. a.i./A.

$$\frac{.25 \text{ ounces}}{.8} = .31 \text{ ounces} \quad \frac{.625 \text{ ounces}}{.8} = .78 \text{ ounces}$$

*There are 43,560 sq. ft./A. Therefore, 340 sq. ft. =

$$\frac{340 \text{ sq. ft.}}{43,560 \text{ sq. ft.}} = .0078/\text{A}$$

**Percent active ingredient is stated on the label.

For adding herbicide to a sprayer, it is convenient to either find a container which holds exactly the desired amount of herbicide, determine how many teaspoons or tablespoons of herbicide equals that amount, or mark on a larger container exactly how full it should be to contain the desired amount of herbicide.


Several factors significantly affect the spray rate of a sprayer:

Spraying Pressure. Spraying should be done at the same tank pressure as calibrating. If, for example, a small compression tank sprayer is calibrated at 25-30 p.s.i. pressure, spraying should be done at that pressure. Spraying at a higher or lower pressure will lead to more or less herbicide being applied as the sprayer passes over the ground.

Nozzle Size. Sprayers must be calibrated for each nozzle size. The larger the nozzle, the more spray put out as the sprayer passes over the ground. Thus, to maintain a particular rate of application, when larger nozzles are used, smaller amounts of herbicide will be added to the water.

Speed of Travel. Spraying should be done at a uniform speed and at the speed at which the sprayer was calibrated. Slower or faster speeds will result in higher or lower rates of herbicide application. For this reason, hand sprayers, such as the small compression tank sprayers, should be calibrated for use by each individual who uses them. Each person has a different speed and pattern of spraying. One person may for example, use 1 gallon to spray 340 sq. ft. while another uses $\frac{3}{4}$ of a gallon and another $1\frac{1}{2}$ gallons.

MICHIGAN STATE UNIVERSITY



COOPERATIVE
EXTENSION
SERVICE

MSU is an Affirmative Action/Equal Opportunity Institution. Cooperative Extension Service programs are open to all without regard to race, color, national origin, sex, or handicap.

Issued in furtherance of cooperative extension work in agriculture and home economics, acts of May 8, and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Gordon E. Guyer, Director, Cooperative Extension Service, Michigan State University, E. Lansing, MI 48824.

This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by the Cooperative Extension Service or bias against those not mentioned. This bulletin becomes public property upon publication and may be reprinted verbatim as a separate or within another publication with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company.

2P-6:85-7.5M-KMF-UP, Price 10 cents, Single Copy Free to Michigan Residents.