



*Plots with better turfgrass quality were treated preventatively with either mancozeb (Dithane T/O or Fore) or chlorothalnil (daconil 2787) fungicides at 14-day intervals.*

## Management of blue-green algae on putting greens

MONICA ELLIOTT AND MARCUS PREVATTE

UNIVERSITY OF FLORIDA

FORT LAUDERDALE RESEARCH AND EDUCATION CENTER

The two most common types of terrestrial algae are green algae, often called true algae, and blue-green algae. The latter are actually a type of bacteria and are also referred to as cyanobacteria. In general, the blue-green algae appear to be the predominant species associated with bermudagrass putting greens and tees in the southeastern U.S. A preliminary survey of putting greens in south Florida demonstrated that the dominant blue-green algae species were *Oscillatoria*, *Lyngbya*, and *Nostoc*. This was also found to be true on bermudagrass greens in Mississippi (Maddox and Krans, 1991). In general, the algal species associated with aquatic environments, such as lakes and waterways on the golf course, are not the same species associated with the greens and tees.

There are two types of algal problems on putting greens and tees, surface algae and the "black layer" phenomenon. "Black layer" is a more complex problem that includes not only blue-green algae but also sulfur-reducing bacteria and specific anaerobic soil conditions created in the soil. The controls for black layer are quite different from controls for the algal slime or crusts observed on bermudagrass putting greens surfaces. The research discussed below was conducted on surface algae as that appears to be the primary problem observed in Florida.

Algae are not plant pathogens. They do not infect (penetrate) the turfgrass plant and cause a disease. If the algae population

increases significantly, they will "slime" the turf and create thin to thick crusts on the soil surface, but they have not physically attacked the turfgrass. While it is possible that blue-green algae produce toxins which may inhibit turfgrass growth, this has not been proven to date. In most cases, it appears that the algal crusts are simply preventing the bermudagrass from physically growing into this area.

Why do blue-green algae "bloom" on putting greens? Because we (Mother Nature and humans) provide them with the perfect environment. By mowing the putting greens extremely short, especially in the summer months, the soil where the algae live all year is exposed to sunlight. Add in excessive rainfall or irrigation, frequent nitrogen applications and a high soil surface pH and an ideal breeding spot for blue-green algae is created.

You must determine the cause of the algal problem before you attempt to fix it. The cultural controls are fairly obvious - dry out the soil surface, break up the algal crusts and alter management or environmental factors which are conducive for algae development. A shady area that never dries out is a perfect place for algae. Document the problem with pictures and facts to convince the management or membership that the landscape needs to be altered.

The weather cannot be controlled, but it is possible to alter irrigation practices that may be contributing to the problem. The algal crusts can be broken by spiking, verticutting, aerifying, hand-raking or any other method that physically disturbs the crusts. Drying out the soil surface can be difficult, especially during a summer with above normal rainfall. Light topdressings

Research Table 1.

Fungicide use to control blue-green algae (cyanobacteria), before the algae develop (preventatively) and after they develop (curatively) on the FGCSA bermudagrass putting green at the Fort Lauderdale R.E.C

Treatment/Formulation*	Rate per 1000 sq ft	Spray Interval <sup>†</sup>	1993 Quality Scores <sup>‡</sup>		1994 Quality Scores <sup>‡</sup>	
			Preventative	Curative	Preventative	Curative
Daconil 2787 90% WDG	3.5 oz	14 days	5.9 bc	5.2 b	5.5 a	3.2 b
Daconil 2787 90% WDG	3.5 oz	28 days	5.3 d	NT	4.4 c	NT
Daconil 2787 90% WDG	6.5 oz	14 days	6.5 a	5.5 ab	5.5 a	3.8 a
Daconil 2787 90% WDG	6.5 oz	28 days	5.6 cd	NT	4.6 bc	NT
Dithane T/O 75% DF	6.0 oz	14 days	6.6 a	6.0 a	5.5 a	3.6 ab
Dithane T/O 75% DF	6.0 oz	28 days	6.1 b	NT	4.9 b	NT
Dithane + CS-7 75% DF	6.0 oz	14 days	6.5 a	5.8 ab	NT	NT
Dithane + CS-7 75% DF	6.0 oz	28 days	6.5 a	NT	NT	NT
Dithane M-45 80% WP	6.0 oz	14 days	NT	NT	5.5 a	3.6 ab
Fore 80% WP	6.0 oz	14 days	NT	NT	5.5 a	3.5 ab
Algaen-X 20% L	4.2 fl oz	14 days	4.3 e	NT	3.9 d	NT
Algaen-X 20% L	12.5 fl oz	14 days	3.9 f	NT	NT	NT
Control			4.3 ef	4.1 c	3.9 d	2.6 c

\*Daconil 2787: ISK Biosciences chlorothalonil; Dithane T/O, Dithane M-45, Fore: Rohm & Haas formulations of mancozeb; Algaen-X: Scotts (Grace-Sierra) quaternary ammonium salts; CS-7 is a spreader-binder surfactant that was added at the recommended rate of 1 pint per 100 gallons of fungicide solution.

<sup>†</sup>Preventative applications were made on 14 day and 28 day intervals and were initiated before algae had developed. Curative applications, applications made after the blue-green algae were already present, were made on 14-day intervals only.

<sup>‡</sup>Quality scores are based on color and density using a scale of 1 to 10 (best quality). Values are means of four replicate plots for nine (1993) or 10 (1994) rating dates. Means in the same column followed by the same letter were not significantly different (P=0.05) according to Waller-Duncan k-ratio t-test. NT=not tested.

of root-zone mix will help absorb some of the moisture and will physically mask the problem areas. However, during the rainy season, it may require more than cultural controls to prevent a blue-green algae problem.

While some superintendents have used a dilute bleach solution (ex: Chlorox) for algal control, the product does not have a legal turf site pesticide label. Certain fertilizer materials, copper sulfate and hydrated lime, have been recommended for algal control. Copper sulfate has a tendency to burn closely mowed turfgrass - especially if the wrong formulation is used. Hydrated lime is effective in initially desiccating (drying out) the algal layer, but it also helps to create an ideal environment for blue-green algae because it increases the soil surface pH. If the root rot disease Bermudagrass Decline is active, the hydrated lime will increase disease activity as this fungus prefers the high soil pH also.

In 1991 when preliminary studies on

blue-green algal control were initiated, the only legal chemical control for algae was the fungicide mancozeb (ex: Rohm & Haas Fore, DuPont Manzate 200, LESCO 4 Flowable Mancozeb). While there is nothing wrong with having only one chemical for control of an organism, it makes plant pathologists like myself very nervous since this would be an ideal situation for chemical resistance to develop within a population. There was also some concern at that time that this fungicide chemical group would be removed from the market. That would have left the golf course community without a legal chemical control product. This was the motivation for initiating this research project.

The two most likely fungicide candidates for algal control at that time were anilazine (ex: Dyrene) and chlorothalonil (ex: Daconil 2787) because these products had been or were still being used in paints for inhibiting algal growth. Anilazine was not included in the study

since it was not expected to be manufactured, and so available for use, in the near future. Quaternary ammonium salts (e.g. Algaen-X) were evaluated beginning in 1993.

### Materials and methods

The experimental site for the chemical trials in 1991-1994 was the FGCSA Research Green located at the Fort Lauderdale Research and Education Center. The 'Tifdwarf' was cut 6 days a week at 3/16 inch. During the summer cycle (May through October), the green was fertilized with 1/2 lb. nitrogen per 1000 sq. ft. every two weeks using a slow release fertilizer source. No other chemical pesticides were applied as a broadcast treatment. If necessary, *Bacillus thuringiensis* (Bt formulated as DiPel) was used to control sod webworms, and spot applications of Dursban insecticidal bait were used for mole cricket control.

To increase algal development, the experimental site was irrigated twice each

## Over the four-year period, various fungicide treatments were examined...

day (10 AM and 4 PM) for 7 minutes, even on days when it rained. No cultural controls were utilized in these studies, so the area was not verticut, aerified or topdressed during the study periods (July-October) each year. This method was utilized to insure that the fungicides were being evaluated under moderate to severe environmental conditions for algal development. In other words, a superintendent's worst nightmare come true!

Over the four-year period, various fungicide treatments were examined but Dithane (75% mancozeb DF; the labeled rate) and Daconil 2787 (90% chlorothalonil WDG; two rates) were always examined each year as preventative treatments applied at 14 day intervals. In other words, after the first application of fungicides was made, the twice daily irrigation cycles were initiated. In 1993 and 1994, these fungicides were also examined as: (a) preventative treatments applied at 28 day intervals, and (b) curative treatments applied at 14 day intervals in

which the fungicides were not applied until the algal population had increased substantially. Other treatments during the four year period included quaternary ammonium salts (ex: Algaen-X), maneb, Dithane (mancozeb) with a surfactant, and three different formulations of mancozeb.

The fungicides were applied in 3-5 gallons water per 1000 sq. ft. using a backpack CO2 sprayer and boom with 40 psi pressure at the handle and stainless steel 8002 Tee Jet nozzles. Deionized water was used to eliminate any interaction due to water source, especially pH effects. The 3 gallon rate was used for all fungicide treatments except the Algaen-X which was applied in 5 gallons water according to the label and information supplied by the company (Grace-Sierra) in 1993. Each plot was 25 sq. ft., and each treatment was replicated four times.

### Results and discussion

The rate at which the algae developed differed between years, primarily due to

the precipitation received during each study period. Plots were rated for quality (color and density) and amount of algae present in each plot. Only quality scores are presented as they are the most accurate assessment of fungicide effect since some products were phytotoxic. The area used for this experiment was of limited size, so it was impossible to use every treatment every year. Results for 1993 and 1994 are presented in Table 1. During July through September, the FLREC received 25 inches of rain in 1993 and 39 inches of rain in 1994. Except for Dithane M-45, only data for products with legal turf site labels are presented. One note on interpreting the table, values in a single column followed by the same letter are NOT statistically different.

Prevention or reduction in algal development was consistently observed when chlorothalonil (both rates) and mancozeb (the labeled rate) were applied at 14 day intervals AND treatments were initiated at the time the environmental pressure for algal development increased



**SERVICES**

DESIGN, SALES, INSTALLATION AND SERVICE OF ALL TYPES OF GOLF COURSE AND COMMERCIAL IRRIGATION PUMP SYSTEMS.

FULLY STOCKED WAREHOUSE WITH SPECIFIC EQUIPMENT TO MEET YOUR IRRIGATION NEEDS.

FULLY EQUIPPED, RADIO DISPATCHED TRUCKS PROVIDE 24 HOUR EMERGENCY SERVICE.

COMPLETE CRANE SERVICE.

INHOUSE PUMP REPAIR FACILITY.

MOTOR REPAIR AND REWIND FACILITY.

PREVENTATIVE MAINTENANCE PROGRAMS.

TOTAL LIGHTNING PROTECTION AND INTERFACE SYSTEMS. UL LISTED INSTALLER.

UNDER WATER WETWELL AND INTAKE CLEANING AND REPAIR.

LICENSED & INSURED ELECTRICAL CONTRACTORS.

**PRODUCTS**

- SYNCROFLO IRRIGATION PUMPING SYSTEMS
- OTTERBINE AERATORS
- EPS LANDSCAPE PUMP STATIONS.
- EPS HYDRAULIC PUMP STATIONS.
- WESCO FOUNTAINS.
- CLAYTON VALVES & PARTS.
- WESSELS HYDROPNEUMATIC TANKS.
- SELF CLEANING INTAKE SCREENS.
- THOMPSON LIGHTNING PROTECTION EQUIPMENT.
- VERTICAL TURBINE PUMPS.
- CENTRIFUGAL PUMPS.
- REPLACEMENT MOTORS.
- CUSTOM CONTROL PANELS, ETL LISTED

**"Servicing South Florida"**

## Prevention or reduction of algal growth possible with timed application of fungicide...

(i.e., initiation of twice daily irrigation). In both years, these treatments were significantly better than the control treatment (no fungicides). The use of a surfactant with mancozeb and the different mancozeb formulations did not increase or decrease its overall effectiveness.

When these products were used at 28 day application intervals, the fungicide treatments still provided better quality than the control but usually had significantly less quality compared to the 14 day application treatments. As can be seen in the graph, the turfgrass quality would increase after a fungicide application and then slowly decrease over the 28 day period. With the 14 day application interval, the quality was more uniform over time.

When the fungicide applications were initiated after algae had developed (curative treatments), the quality of the treated plots was still better than the control but did not equal the preventative fungicide treatments. This was especially evident in

1994 when the environmental conditions were quite conducive for algal growth.

The 90% WDG formulation of Daconil 2787 was utilized throughout the four year period for consistency during the overall study. When you are working with small plots, dry formulations are much easier to measure than flowable materials. This formulation has since been replaced with an 82.5% SDG formulation. Check the Daconil 2787 labels for the current legal use rates.

Mancozeb fungicides are one of the very few fungicides that have a label stating that use of a surfactant may increase efficacy. There did not appear to be a strong advantage in this particular study. Daconil 2787 should not be mixed with a surfactant as one is already part of the formulation. A minimum of 2 gallons water per 1000 sq. ft. should be used to mix all fungicides. (Yes, the labels do state this!!) I used 3 gallons in this study because I wanted to be sure the fungicides came in contact with the algae. Re-

member, you want to stop algal growth. The algae are not growing on the leaf surface but on the soil surface!

Different mancozeb formulations were evaluated as some superintendents believed they observed better efficacy with mancozeb formulations containing the blue dye. Dithane T/O and Dithane M-45 (which is not labeled for turf) do not contain the blue dye that Fore contains. Dithane M-45 was used because its formulation is more similar to Fore than Dithane T/O. No differences were observed between these formulations.

There has been some concern about using mancozeb products, which contain zinc, due to soil test reports that indicate the soil contains an "excessive" amount of zinc. The word "excessive" is misleading. Usually, soil zinc levels over 20-60 mg/kg soil are considered excessive, especially when you consider that turf normally only needs a few mg to satisfy its nutritional requirements. However, turfgrasses can tolerate much higher soil zinc levels. A study by Spear and Christians (1991) on bentgrass demonstrated that even at zinc soil levels of 4000 mg/kg, there was no consistent plant damage. I am not aware of any reports of zinc toxicity of bermudagrasses used on golf courses.

Quaternary ammonium salts for use on turfgrass sites are sold under the trade names Algaen-X and Consan Triple Action 20. There has been considerable confusion regarding the appropriate use of these products, primarily because the label itself is confusing and information provided by one company was inconsistent with the label. In 1993, two rates



### TESTING

- Soils
- Water
- Physical
- Tissue
- Environmental

**Tom Burrows**  
 Certified Agronomist  
 Turfgrass Specialist

Phone: 407-692-1221  
 1741 San Souci, Stuart, FL 34994

CGCS  
 Golf Course Supt. - 29 years  
 FGCSA Past President  
 Penn State - 1960  
 Musser Int'l Turf Foundation - VP

(305) 581-0444

**SPREAD-RITE, INC.**  
 CUSTOM SPREADING  
 FERTILIZER • DOLOMITE • ARAGONITE

LAMAR SAPP

6001 S.W. 19th STREET  
 PLANTATION, FL 33317  
 MOBILE: (407) 479-9417

YOUR "ONE-STOP" SOURCE  
 FOR AMERICA'S LEADING  
 GOLF SURFACES & SUPPLIES!

**PARTAC®**  
 GOLF COURSE  
 TOP-DRESSING



HEAT TREATED  
 PLUS CONSTRUCTION,  
 CART PATH, SAND TRAP,  
 AND DIVOT REPAIR MIXES

**TYPAR®**  
 GEOTEXTILES & TURF BLANKETS

**Terra-Green®**

SOIL CONDITIONER

NEW **PARTAC®** NEW  
**GREEN SAND**  
 DIVOT REPAIR MIX

A UNIQUE SOLUTION TO UNSIGHTLY DIVOTS!  
 PARTAC® GREEN SAND BLENDED  
 WITH PARTAC® PREMIUM  
 TOP-DRESSING FOR EXCELLENT  
 GERMINATION AND  
 DEEP GREEN COLOR!

THE BEST DIVOT REPAIR MIX AVAILABLE!

AVAILABLE NATIONWIDE  
**800-247-2326**  
 IN N.J. 908-637-4191

**PARTAC PEAT CORPORATION**  
 KELSEY PARK, GREAT MEADOWS, N.J. 07838  
 ORLANDO (FL) MEMPHIS (TN) RENO (NV)

HIGHLIGHT GOLF HOLES,  
 IMPROVE VISIBILITY,  
 & MAINTAIN CONSISTENCY

U.S. GOLF HOLE  
**TARGETS™**



**TerraFlow™**  
 Drainage Systems

FLEXIBLE GOLF COURSE MARKERS  
 SAND TRAP PUMPS  
 RAKES, PLUG PUSHERS, SQUEEGEES  
 AND MANY MORE  
 GOLF SPECIALTY PRODUCTS




Prevention of blue-green algae with fungicide treatments. Mancozeb fungicide (left) vs. control with no fungicide (right).

were used in the minimum amount of water permissible by the label - 5 gallons per 1000 sq. ft.

The high rate was distinctly phytotoxic to the bermudagrass. The low rate also caused some phytotoxicity but was not consistent in this characteristic. The product did not prevent the development of blue-green algae. This was not expected as it is strictly a curative product. However, even after the blue-green algae developed, there was no observable algal control in either 1993 or 1994.

### Summary

1. Both chlorothalonil and mancozeb were effective for controlling blue-green algae on bermudagrass putting greens.
2. The chlorothalonil rate required for control will depend on the level of algae present and environmental conditions.



## We understand the importance of a good mix.

That's why it's important that you get to know Atlantic FEC. Because we provide a variety of specialty mixes and related products specifically formulated for the South Florida turfgrass industry.

Each turfgrass mix is delivered by Spyder equipped trucks in water-resistant paper or plastic bags. Bulk material is available in either normal size Killebrews or "small door" (24") Killebrews to accommodate your turf spreader. Drop trailers, sulfur coated potash, and new Micro Prill Greenskote, a quality putting green blend, also available.

Call us toll free at 1-800-432-3413 or visit us at 18375 S.W. 260th Street in Homestead to find out how you can benefit from quality mix from the Atlantic FEC turfgrass professionals.


**Atlantic FEC**  
Better Mixes Since 1923.



**"The Most Versatile  
Service Lift  
Available In The  
Turf Industry"**

Wheeltronic's "VGL"  
Multipurpose Above Ground  
Lifts Were Specifically Designed  
To Provide Flexibility When  
Servicing Turf Equipment,  
Golf Carts And Road Vehicles.

**No Other Lift Offers  
So Many Features  
"Just Ask Those That  
Have Them"**



**For More Information Call 1-800-382-6735**

Distributed by: Du Cor International • 1011 W. Lancaster Rd., Orlando, FL 32809  
Tel: 407/859-4390 • Fax 407/851-8430

## ALGAE ON TURFGRASS A PROBLEM?



**E.P.A. REGISTERED ALGAEICIDE  
AND FUNGICIDE FOR TURFGRASS**

- EFFECTIVE ● ECONOMICAL
- CURATIVE ● BIODEGRADABLE

For a Free Sample Write or Call:



**PARKWAY  
RESEARCH**

**PARKWAY RESEARCH CORPORATION**  
1-800-256-FOOT

13802 Chrisman Rd. 713-442-9821  
Houston, Texas 77039 FAX 713-590-3353

3. The application intervals used for chlorothalonil and mancozeb will vary with environmental conditions. During periods of high stress, the 14-day interval will probably be most effective.
4. There were no differences in turfgrass quality between different formulations of mancozeb or in the use of a surfactant with mancozeb.
5. Quaternary ammonium salts were not effective under the conditions used for evaluation in this study. Research at Texas A&M indicates these products are only effective when they are used as drenches.
6. Initiate use of the fungicides before or shortly after a weather pattern is established that is conducive for algal development. Do not wait until there is a significant algal population on the turf area to initiate treatments.
7. Never, ever start a fungicide program without initiating cultural control practices also! Use good integrated pest management strategies and safe chemical practices for controlling blue-green algae on the golf course.

### Literature Cited

- Maddox, V., and J. V. Krans. 1991. The effect of algae and cyanobacterium composition and density on putting greens in Mississippi. Agron. Abstr. p. 179.
- Spear, G. T., and N. E. Christians. 1991. Creeping bentgrass response to zinc in modified soil. Commun. Soil. Sci. Plant Anal. 22:2005-2016.

### Acknowledgements

We thank Dr. Carol Bourne for identification of the algae samples and information on maintaining algae in culture. We also thank those suppliers who participate in the South Florida Turfgrass Exposition as the funds raised are used to pay the salary of Marcus Prevatte who maintains the FGCSA research greens.