

# Fairy Ring Identification, Symptoms, Development And Cultural and Chemical Control

Courtesy North Carolina State University

## SYMPTOMS

The symptoms of fairy ring (*Basidiomycetes*) appear in patches, rings, or arcs that are initially 1 foot or less in diameter, but expand in size year after year, reaching up to several hundred feet in diameter in old turf stands. Most fairy ring fungi do not infect or parasitize the turf.

Instead, growth of these fungi in the soil can indirectly affect, or even kill, the turfgrass above. Three types of fairy ring symptoms are observed in turfgrasses: Type I, Type II, and Type III. A Type I fairy ring causes the soil and thatch to become hydrophobic, killing the turf in patches, rings, or arcs. In areas affected by a Type I fairy ring, the thatch and soil are extremely dry and repel water. Type II fairy rings appear as rings or arcs of turf that are dark green and growing more quickly than the surrounding turf. In a Type III fairy ring, mushrooms or puffballs are produced in a ring or arc. The type of symptom expressed by a fairy ring may change during the year according to weather conditions. Type III fairy ring symptoms are more prevalent during extended periods of wet weather. Type I and Type II fairy ring symptoms are most common during hot, dry weather in the summer.

Characteristic	Description
Host Grass Species	all
Month(s) with symptoms	all
Stand Symptoms	patches (1-3 feet), rings

**Note:** Still not sure if this is the right disease? The Turfgrass Disease Identification program may be helpful. Or consult the experts at the Turf Diagnostics Lab. Check the TurfFiles glossary for definitions of unfamiliar terms.

## FACTORS AFFECTING DISEASE DEVELOPMENT

Fungi that cause fairy rings are common inhabitants of forested areas. These fungi begin growing on a source of organic matter, such as an old stump, waste lumber, or dead tree roots in the soil. Once established, the turf produces thatch and organic matter, which provides a source of food for continued expansion of the fairy ring.

On golf course putting greens, fairy rings are often observed spreading into the root zone mix from the surrounding native soil. Mushroom fungi are also prolific spore producers and may be spread into putting greens by wind, water, or other means.

Type I fairy rings are most damaging to turf and are encouraged by drought stress, inadequate irrigation, and infrequent aeration.

Type II fairy ring symptoms are a result of the release of nitrogen and other nutrients into the soil as organic matter is degraded by the fairy ring fungi. These symptoms are most evident in turf that is deficient of nutrients, especially nitrogen and iron.

Type III fairy rings are encouraged by over-irrigation or poor soil drainage.

## CULTURAL CONTROL

The most effective means for control is to prevent the fairy ring fungi from becoming established in the turf. Remove large pieces of woody material (stumps, waste lumber, and dead tree roots) before turf is planted to prevent the establishment of fairy rings. Landscape contractors should remove this debris around new construction sites before seeding or sodding.

Installation of a plastic barrier between the root zone mix and surrounding native soil may limit the spread of fairy ring into

<b>Foliar Symptoms Location/Shape</b>	dieback from leaf tip, no distinct leaf symptoms
<b>Foliar Symptoms Color</b>	tan, yellow, orange
<b>Root/Crown Symptoms</b>	none
<b>Fungal Signs</b>	mushrooms, puffballs, none

golf course putting greens.

Power raking or vertical mowing to remove excessive thatch will help to minimize fairy ring problems. Golf course superintendents should regularly aerify and topdress putting greens to prevent thatch buildup and maintain soil aeration. Avoid extremes in soil moisture (too wet, too dry), apply nitrogen based on local University recommendations, and ensure balanced fertility through regular soil testing.

Once a fairy ring appears, the best management practices depend on the type of symptom that is observed. To control a Type I fairy ring, the water-repellent thatch and soil beneath the affected turf must be re-wet. Hollow-tine aerification, spiking, water-injection, application of soil surfactants, and heavy irrigation are effective strategies for re-wetting this hydrophobic layer. Affected areas should be hand-watered to prevent over-watering of the unaffected turf.

Symptoms of a Type II fairy ring can be masked with an application of nitrogen or iron. This will cause the surrounding turf to green-up, making the affected turf less evident. Collect soil or tissue samples for nutrient analysis from the turf immediately surrounding the Type II fairy rings, and correct any nutrient imbalances as recommended. Use caution when applying nitrogen to mask Type II fairy ring symptoms on cool-season grasses during the summer. Too much nitrogen may

(Continued on Page 10)

## MGCSA FALL MIXER AT THE HORSE AND HUNT CLUB

"Trading Sticks for Guns in Prior Lake"

OCTOBER 4, 2010 - GO TO [WWW.MGCSA.ORG](http://WWW.MGCSA.ORG) FOR DETAILS





# Fairy Ring-

(Continued from Page 9)

over-stimulate the grass and lead to the development of more serious diseases. In this case, iron should be used to increase turf color without causing excessive foliar growth.

Drastic methods for control of fairy rings, such as soil fumigation, removal of infested soil, or turf renovation by tilling and mixing the soil may be effective in the short-term, but the fairy rings usually become re-established over a period of years.

## CHEMICAL CONTROL

Over 60 species of fungi have been associated with fairy ring symptoms in turfgrasses, and these species likely vary in their sensitivity to fungicides. Control of fairy rings with fungicides is a site-specific venture for this reason. Turfgrass managers should experiment with different products to identify those that will control the disease in their location.

Fungicides are most effective for fairy ring control when used on a preventative basis. Curative applications have little effect because the symptoms are caused by a change in the soil environment, and fungicides do nothing to change the soil. A preventative fungicide program should be initiated in the spring when mean daily soil temperatures are consistently above 55°F. Regular use of soil surfactants will help to maintain uniform soil moisture and may reduce the appearance of Type I fairy ring symptoms.

Because fairy ring fungi are in the thatch and soil, fungicides must be watered-in or applied in large volumes of water for best results. Applications in 2 gallons H<sub>2</sub>O per 1000 ft<sup>2</sup> followed by 0.25" of irrigation have provided excellent results in research trials at NC State University. Irrigation must be applied immediately before the spray begins to dry on the turfgrass foliage. Tank-mixing some fungicides with a soil penetrant may also enhance movement into the soil and improve fairy ring control.

The DMI fungicides provide excellent preventative control of the puffball fungi *Lycoperdon perlatum* and *Vascellum pratense* in golf course putting greens. Two applications on a 30 day interval, beginning in spring when 5-day average soil temperatures reach 55°F, have provided season-long control in many cases. The DMI fungicides, however, should not be tank-mixed with soil surfactants as this may reduce their efficacy and increase the potential for phytotoxicity.

(Editor's Note: © North Carolina State University. This information sheet was prepared by Lane P. Tredway, Gail G. Wilkerson, Bridget R. Lassiter, Jenifer J. Reynolds, and Gregory S. Buol. Departments of Plant Pathology and Crop Science, College of Agriculture & Life Sciences, North Carolina State University. Prepared March 23, 2009.

Fungicide	Efficacy (1)	Resistance/Risk (2)	Class (3)	Products (4)
triadimefon	++++	2	DMI	Bayleton
metconazole	+++	2	DMI	Tourney
azoxystrobin	+++	3	QoI	Heritage
pyraclostrobin	+++	3	QoI	Insignia
azoxystrobin	+++	3	QoI + DMI	Headway
+ propiconazole				
flutolanil	++	2	carboxamide	ProStar
flutolanil +	++	2	carboxamide +	SysStar
thiophanate-methyl			benzimidazole	
polyoxin D	++	2	polyoxins	Endorse

### Efficacy Codes:

++++	excellent control when conditions are highly favorable for disease development
+++	good control when disease pressure is high, excellent control when disease pressure is moderate
++	good control when disease pressure is moderate, excellent control when disease pressure is low
+	good control when disease pressure is low
0	does not provide adequate control under any conditions
?	cannot be rated due to insufficient data

### Resistance Risk:

- Rotating and tank-mixing not necessary, but recommended to avoid potential side effects from continuous use of same chemical class.
- Rotate to different chemical class after 3-4 applications; tank-mixing not necessary.
- Rotate to different chemical class after 2-3 applications; tank-mixing not necessary.
- Rotate to different chemical class after 1-2 applications; tank-mixing not necessary.
- Rotate to different chemical class after 1-2 applications; tank-mixing with low or moderate risk product recommended.
- Rotate to different chemical class after EVERY application; tank-mix with low or moderate risk product for EVERY application.

Continual use of fungicides with similar control mechanisms (modes of action) can result in fungi that are resistant to some chemicals. Poor or ineffective disease control can be expected when this occurs. Managers can reduce the chances of this happening by mixing or alternating fungicides belonging to different chemical classes.

Recommendations of specific chemicals are based upon information on the manufacturer's label and performance in a limited number of trials. Because environmental conditions and methods of application may vary widely, performance of the chemical will not always conform to the safety and pest control standards indicated by experimental data. When more than one brand name exists for an agricultural chemical, the name of brand that first came onto the market is listed first. Otherwise, brand names are listed in alphabetical order. The order in which brand names are given is not an indication of a recommendation or criticism.

Recommendations for the use of agricultural chemicals are included in this publication as a convenience to the reader. The use of brand names and any mention or listing of commercial products or services does not imply endorsement by North Carolina State University or discrimination against similar products or services not mentioned. Other brand names may be labeled for use on turfgrasses. Individuals who use agricultural chemicals are responsible for ensuring that the intended use complies with current regulations and conforms to the product label. Be sure to obtain current information about usage regulations and examine a current product label before applying any chemical. For assistance, contact your county's Cooperative Extension agent.

## STATEMENT OF OWNERSHIP MANAGEMENT & CIRCULATION

- Title of publication: *Hole Notes*.
- Publication No.: 1082-7994.
- Date of Filing: October 1, 2010
- Frequency of issue: monthly, except bi-monthly December-January and February-March.
- No. of issues published annually: 10.
- Annual subscription price: \$20.00.
- Mailing address of known office of publication: 217-D Minnetonka Ave. S., Wayzata, Minnesota 55391
- Mailing address of headquarters or general business office of publisher: Same as No. 7.
- Names and address of publisher, editor and managing editor: Publisher and Managing Editor: Scott Turtinen - same as #7. Editor: Jack MacKenzie - same as #7.
- Owner: Minnesota Golf Course Superintendents' Association, 217-D Minnetonka Ave. S., Wayzata, Minnesota 55391
- Known bondholders, mortgages, and other security holders owning or holding 1% or more of total amount of bonds, mortgages, or other securities: None.
- For completion by nonprofit organizations authorized to mail at special rates: Not applicable.
- Publication title: *Hole Notes*.
- Issue date for circulation data below: 9-15-10
- Extent and nature of circulation:

Column 1: (Below) Actual number of copies of each issue. Column 2: (Below) Average number of copies during preceding 20 months of single issue published nearest to filing date.

	C 1	C 2
a. Total copies printed	850	850
b. Paid circulation:		
1. Mail subscriptions	794	794
2. Sales through dealers and carriers, street vendors and counter sales	0	0
c. Total paid circulation	794	794
d. Free distribution/ mail	0	0
e. Free distribution/ other	0	0
f. Total free distribution	0	0
g. Total distribution	794	794
h. Copies not distributed	56	56
i. Total	850	850
j. Percent paid and/or requested circulation	100	100

- I certify that all information on this form is true and complete.

- Signed by Scott Turtinen, Publisher