Results of the Superintendent-Pro Tourney

LOW PRO Larry Ringer 75 Charlie Bassler 76 Lumbart 77 Quinn Sullivan 78 Romans 78 Greg Shrews 79 Paul Hauiland 80 Willis Johnson 80)
SUPERINTENDENT LOW GROSS Herb Hienlien 90 Harry Allen 80	1
SUPERINTENDENT LOW NET Ralph Strouth 100 36 64 John Tutich 100 26 74	
TEAM NET Ralph Strouth and Chuck Walker	

GUEST LOW NET Mike Hannah
GUEST LOW GROSS Tom Comalli
LONGEST DRIVE Greg ShrewsPro Columbia
#3 Ed Wilson 5'8" #7 Dennis O'Leary 9'11" #17 Larry Ringer 7'11"
50-50 DRAWING WINNER Mark Shields\$64.00





Everything you need to control diseases and weeds in turf and ornamentals.

Ten turf diseases and almost 100 kinds of weeds yield to the Diamond Shamrock Pro-4 turf system. One broad-spectrum fungicide. One preemergence herbicide. One postemergence broadleaf herbicide. One postemergence grassy weed herbicide. That's Diamond Shamrock Pro-4.

DACONIL 2787°. The new chemistry fungicide. And now it's a flowable, as well as a wettable powder.

DACTHAL® preemergence herbicide controls *Poa annua*, crabgrass and 15 other weeds. Available in two formulations: wettable powder and granules.

DACAMINE® postemergence herbicide. Non-volatile 2,4-D. Kills more than 70 broadleaf weeds.

DACONATE® 6 postemergence herbicide. The organic arsonate with a high-quality surfactant already mixed in.

Regional Office

Three Commerce Park Square 23200 Chagrin Boulevard Beachwood, Ohio 44122 216-464-5620

Diamond Shamrock Ask for our Product AGRICULTURAL CHEMICALS DIVISION Information sheets.

More Terminal Bermuda News

Dr. Doug Hawes reports the following observations and feedback on bermudagrass winterkill. On the University of Maryland demonstration plots, only Midiron (P-16) survived the winter to any extent. Tufcote appears lost in Dr. Hawes's Tee Turf Research Plots and in a new area he overseeded last fall to Penncross. The overseeding was done to continue studies on bermuda-bentgrass turf. A similar overseeding done at the same time of Kentucky bluegrass into Meyer zoysiagrass was made without any apparent loss of Meyer.

A large plot area of Tufcote not currently in a research project on the farm was lost except for a strip within 20 feet of a snow fence on its north and west sides. Protection from northwest winds appeared to be a factor in survival of Tufcote lawns in the Bowie area also. Older (thatchy) lawns appeared to suffer more damage than newly established lawns. Traffic also increased the chances of winterkill.

Research plots at USDA-Beltsville, Md. suffered severe damage as observed by Doug on May 5th. Essentially all Tufcote was killed. Midiron looked a little better. One as yet unreleased Kansas variety and one from Missouri came through in good shape, also a coarse bladed selection from Columbia Country Club. Dr. Hawes suggests you come to the Beltsville Field Day to hear a report on this by Jack Murray. Field Day is scheduled for August 24th.