

# GUIDELINES FOR CONTROLLING MOSS IN GREENS

The quality of golf course greens by present day standards is often determined by greens' speed. Golf course superintendents are mowing greens shorter and keeping the nitrogen fertility lower than ever before to obtain faster speeds. A consequence of these practices has been a reduction in turfgrass vigor to a point whereby the greens are much more prone to weed encroachment. One of the more troublesome weeds to have become a problem is moss.

Until recently, the only known means of controlling moss was through the use of mercury products. With the support of the Metropolitan Golf Association, research was conducted to look at means of controlling this serious weed. This research identified both chemical and cultural tools that could be used in a moss eradicating program.

## CHEMICAL CONTROL

Herbicides and other materials offer hope in controlling moss on bentgrass greens. In the early spring, moss commences its growth much earlier than bentgrass, giving it an early competitive advantage. Hydrated lime applied in late March at 3 to 5 pounds per 1000 square feet will burn back the moss during this period. The lime can be spread easily if mixed with a dry sand topdressing.

An effective treatment for moss control would be the Scotts Goosegrass Control; a betasaronstar combination. Labeled for use on bentgrass greens, this product provided 83% control from only a single application. While this product will cause some discoloration, it appears to be one of the more promising moss control products.

Siduron (Tupersan) and bentazon (Basagran) provided from 53 to 74% control of moss. While they were not quite as effective as the Scotts products, both siduron and bentazon were much safer since no injury occurred for either product.

You should note that with the exception of bentazon the most effective treatments are preemergence herbicides. While it can't be determined from these trials whether the effect is pre- or post-emergent, it should be mentioned that the herbicidal activity of these materials on moss was chronic. It was several weeks before we noticed any significant decrease in moss populations.

## CULTURAL CONTROL

Chemicals only offer a partial solution to the moss problem. Unless cultural steps are taken to increase turfgrass vigor, chemical control of moss will be ongoing battle. We designed studies to look at the effects of cultivation techniques and fertility on moss eradication. The results clearly demonstrated that culture can be changed to the detriment of moss.

While silvery thread moss will tolerate dry conditions, it is favored by an abundance of free water. Core cultivation immediately followed by sand topdressing would create a system of "vertical drains" that would facilitate a rapid water removal of the surface. We found that moss removal was hastened where this practice was followed compared to core cultivation alone. Deep spiking was also beneficial compared to core cultivation alone.

Nitrogen and iron are the most important tools in a moss eradication program. Moss control improved as the rate of nitrogen was increased. Moss was eliminated over two


growing seasons from plots that were initially 40% moss by increasing nitrogen rates to about 0.8 lbs. per 1000 square feet per growing month (6 lbs. N/year). Iron applications at a rate of 6 ounces per 100 sq. ft. per month were beneficial during the first year, especially at the higher rates of nitrogen. Iron had no effect on moss in the second year.

While we didn't measure greens' speeds, these high nitrogen treatments no doubt resulted in slower speeds. The bottom line though, is if you have moss, you are going to have to at least temporarily increase nitrogen rates. Effects on greens' speeds can be minimized by careful control of water, double cutting, or increasing potassium levels.

Moss control research has until now looked at fertility and herbicides independently. Studies will be conducted this year to look at combinations and nitrogen fertility in moss eradication "programs". Perhaps this research will identify more reasonable nitrogen rates to use in conjunction with a herbicide program to eliminate moss from greens.

In summary enough information is known for a superintendent to develop a legal moss control program. Early spring applications of hydrated lime, followed about a month later and in the early fall with a herbicide are the first steps in controlling moss. Increasing your nitrogen levels during this period will no doubt improve the competitive advantage of desirable grasses at the expense of moss. Furthermore, control your soil moisture levels through careful irrigation and by providing good drainage throughout the soil profile.

Credit: Our Collaborator, Northeastern GCSA, Sept. 1990



**COUNTRY CLUB**  
*Sales, Inc.*

830 LATOUR CT. (707) 255-2828  
NAPA, CA 94558 (800) 252-6660



**JENKINS MACHINERY CO.**  
Agricultural Machinery  
Construction Machinery  
Turf Machinery

I. F. "GEORGE" MOORE  
CRAIG WATTERS  
KEVIN MILLER

Bus. (510) 685-6685

JENKINS MACHINERY CO.  
1848 Arnold Industrial Place, Concord, CA 94520  
CA/NV (800) 642-2456 FAX 510-685-9303

*Christensen Irrigation Co., Inc.*  
*Golf Course Irrigation Systems*

*W. Jack Christensen*  
*President*

785 Golden Gate Avenue, #401 1820 E. Garry Ave. #116  
San Francisco, CA 94102 Santa Ana, CA 92705-5804  
(415) 255-3155 • Fax (415) 255-3155 (714) 261-6076 • Fax (714) 756-0663