

# NEW TURFGRASS DISEASE CULPRITS

By Todd Burkdoll, market development specialist, BASF Turf and Ornamentals

The presence of several new turfgrass diseases has increased in recent years. Three new diseases in particular – Pythium root dysfunction, brown ring patch and rapid blight – are a challenge for turf management and maintenance professionals. Fortunately, researchers have made headway regarding how to detect and treat these destructive new diseases.

## Tackling New Diseases

New turfgrass diseases can evolve for a number of reasons and several factors contribute to the prevalence of disease such as geography, moisture and temperature.

Stress caused by heat, drought and excess moisture can weaken turf and make it more prone to disease, whereas healthy turf is less susceptible to disease.

A specific maintenance challenge facing golf courses is keeping turf short and healthy, particularly on greens, because when golf course superintendents use plant growth regulators and/or frequent mowing practices they are stressing the vitality of the plant.


It is a difficult balance, especially when new diseases emerge and superintendents do not know how to treat them.

University and industry researchers are addressing these three emerging problem diseases. To avoid being caught off-guard, turf professional and course superintendents should take time to learn more about these diseases. Doing so will help identify and treat the diseases, and in some cases, avoid them altogether.

## Pythium Root Dysfunction

Discovered in North Carolina in 1994, Pythium root dysfunction attacks putting greens and is limited to newly constructed greens less than eight years old. It is most commonly found in the Southeast, but also occurs in Midwestern areas with harsh summers. Bentgrass is most susceptible to the disease, which occurs on turf stressed from one or more of the following factors: high heat, repeated close mowing, low fertility schedules and drought.

*(Continued on page 11)*



Whether it's removing hazardous trees, pruning for views of a green, or opening up a fairway; we will work with you to design a tree care management plan specific to your golf course.

Serving the Greater Bay Area from 15 service centers.



**cagwin & norward**

800-891-7710

[www.cagwin.com](http://www.cagwin.com)




Cut water use by 15 to 25 percent



Aqua-PhyD's in-line water and soil treatment system uses resonant frequency technology to reduce soil compaction and harmful Na levels

Works with all water sources including brackish and reclaimed water

Increase root growth and nutrient uptake



Installation at a Southern California Golf Course

888-288-AQUA (2782)

Call toll-free for more information or email [info@aqua-phyd.com](mailto:info@aqua-phyd.com)



**Satisfaction Guaranteed!**

[www.aqua-phyd.com](http://www.aqua-phyd.com)

*(Continued from page 7)*

## NEW TURFGRASS DISEASE CULPRITS

Pythium root dysfunction causes the roots and crown of turfgrass to turn brown or black. The symptoms are most visible during the summer, but the disease actually spreads during spring and fall, when it is cool and wet.

Because symptoms are less prevalent on plants with a strong root system, there are several cultural practices superintendents can undertake to minimize damage caused by Pythium root dysfunction. Root enhancement techniques – specifically aerification, nutrition supplements, verticutting and reduced irrigation – are helpful in counteracting symptoms of the disease.

Irrigation management is also extremely important. Clay and compacted soils are more likely to harbor Pythium root dysfunction because of reduced drainage.

It is less difficult and less expensive to prevent Pythium root dysfunction than it is to try to cure it. Fungicides such as pyraclostrobin and triticonazole are two of the most effective at preventing the disease.

Dr. Lane Tredway, turfgrass pathologist at North Carolina State University, is one of the foremost experts on Pythium root dysfunction. To learn more about his research and information on NC State's Center for Turfgrass Environmental Research & Education, visit [www.turffiles.ncsu.edu](http://www.turffiles.ncsu.edu).

### Brown Ring Patch

Formerly known as waitea patch, brown ring patch has been reported sporadically throughout the Midwest and is a mounting problem in Southern California. Occurring primarily on greens with high annual bluegrass (*poa annua*) populations, the disease is prevalent in warm and moist conditions.

Initial symptoms of brown ring patch are thin, yellow, concentric rings several inches in diameter that turn brown under hot or wet conditions. Once established, brown ring patch can quickly damage turfgrass. Temperature plays a significant role in regards to whether or not brown ring patch occurs. The disease does not spread in hot or cold conditions, but rather during times of mild (mid-60s to low-80s F) temperature.

Cultural control options of aerification and higher mowing heights are sometimes used to combat brown ring patch. Alternating among several fungicides – pyraclostrobin, propiconazole and triticonazole – has been an effective treatment.

Dr. Frank Wong, assistant plant pathologist at the University of California-Riverside, is considered one of the top brown ring patch researchers. For more information, visit UC Riverside's Department of Plant Pathology and Microbiology at [www.plantpathology.ucr.edu](http://www.plantpathology.ucr.edu).

*(Continued on page 14)*



(Continued from page 11)

## NEW TURFGRASS DISEASE CULPRITS

### Rapid Blight

Rapid blight occurs in the fall and winter, affecting several annual winter grasses used to overseed Bermudagrass. Affected species include ryegrass, annual bluegrass and poa trivialis. It is primarily seen in the Southwest, including Nevada, Arizona and Southern California, as well as on coastal areas in the Southeast and Northeast. Perennial grasses are not affected by rapid blight.

The disease is associated with a marine organism and cases of rapid blight rise significantly in areas where superintendents use reclaimed water or water with high salinity for irrigation. The disease can occur on any area that has been overseeded, but is usually treated only on putting greens.

Dr. Mary Olsen, plant pathology specialist for the University of Arizona-Tucson, has confirmed that rapid blight is caused by an obscure microorganism that prior to its discovery in turf was known to infect in marine plants such as seagrass, diatoms and algae.

Rapid blight shows itself as water-soaked, slightly sunken and darker looking turf. It turns yellow and dies in patches.

The primary cultural control option is to use better quality irrigation water, avoiding reclaimed water, if possible. Pyraclostrobin provides the most effective preventative control, with mancozeb as a less effective alternative.

Olsen is a leading rapid blight researcher. The University of Arizona's Division of Plant Pathology and Microbiology is available on the Web at <http://ag.arizona.edu/PLP>.

### Prevention, Education Key

To avoid being caught off guard by new diseases, it is important to stay educated on new diseases, be consistent with preventative tactics and devote time to detection efforts.

Part of being proactive is keeping up with the latest research and information about turfgrass disease. Superintendents who collect and absorb background information are better prepared when they encounter a problem – they know what they are dealing with and who can help them.

Some superintendents are quick to write off an undiagnosed problem as being untreatable by a particular fungicide they have already applied, so they simply retreat with a different product. Instead, they should take a turf sample and send it in to a diagnostics lab.

Fungicide manufacturer representatives, university extension personnel and other course superintendents are also good sources of information. It is wise to seek the help of others if they encounter an abnormality they do not recognize.

It may seem like common sense, but it is important for superintendents to walk their courses every day to keep an eye out for abnormalities. New diseases such as pythium root dysfunction, brown ring patch and rapid blight can cause problems quickly if undetected.

### Sidebar: Getting to the Root of the Southeast's Pythium Disease

The good news for Charles Sheffield, superintendent at Croasdaile C.C. in Durham, N.C., is that only one disease significantly disturbs his turf. The bad news is that it is Pythium root dysfunction caused by Pythium volutum. Superintendents and pathologists living in the Southeast are learning that Pythium root dysfunction on creeping bentgrass greens is a plague.



**PROFESSIONAL GRADE FERTILIZER**  
**Spread The Word**

**N** Micro Green®  
Greens King®  
Short Kut

**P** Best® Phos  
6-20-20XB®  
9-Iron Mini

**K** Greenskote®  
Mini Pro®  
Triple Pro®

For More Information, Contact Your BEST®  
Distributor, or Call 800-832-8891



**Quality sand based greens sod:**

- Penn A4
- Dominant X-treme
- Northwest Poa Greens sod

1-800-267-7763 [www.bossod.com](http://www.bossod.com)