Annual Bluegrass Control Progress Report

TITLE:

Selective Control of Annual Bluegrass (Poa annua L.) in Creeping Bentgrass Putting Greens

INVESTIGATORS:

Jim Baird, Brent Barnes, Alea Miehls, and Vanessa Ferrel Department of Botany and Plant Sciences 2137 Batchelor Hall University of California, Riverside, CA 92521 (951) 827-5630 Office • (951) 333-9052 Cell jbaird@ucr.edu

SPONSORS:

Golf Course Superintendents Association of Northern California Northern California Golf Association San Diego Golf Course Superintendents Association Sierra Nevada Golf Course Superintendents Association Southern California Golf Association Southern California Golf Course Superintendents Association Southern California Section, PGA

COOPERATORS:

Gary Custis, PBI Gordon
Doug Houseworth, Arysta LifeScience Corporation
Suk-Jin Koo, Moghu Research Center
Bob Mack, Helena Chemical
Todd Mayhew, Valent Corporation
Dean Mosdell, Syngenta
Chris Olsen, Bayer Environmental Science

OBJECTIVES:

Evaluate existing and experimental herbicides for selective removal of annual bluegrass that persists in creeping bentgrass putting green turf. Evaluate herbicides and herbicide combinations for potential bentgrass injury prior to inclusion in experiments on golf courses throughout the State.

LOCATION:

UCR Turfgrass Research Facility, Riverside CA

SOIL:

Loamy sand amended with sand topdressing

EXPERIMENTAL DESIGN:

Randomized complete block; three replications

PLOT SIZE:

3 ft x 6 ft

SPECIES/CULTIVARS:

Creeping Bentgrass (Agrostis stolonifera L.) 'Cobra'

MOWING HEIGHT:

0.180 inches; 3 days/wk

IRRIGATION:

80% ETo (historical from previous week)/DU

CULTIVATION:

alternate bi-monthly verticutting/solid-tine aeration; topdressing monthly

SPRAYER:

Bicycle with two 8003VS nozzles • 20-inch spacing • 35 psi • 510 ml/nozzle/30 sec = 80 GPA • Metronome = 50 beats = 1 mph

APPLICATION DATES:

A = May 13 and June 10,2010

B = May 13, 20, 28 and June 4, 2010

C = Bi-weekly beginning May 13, 2010

D = June 10, 2010

E = Scheduled for September 1, 2010

DATA COLLECTED:

Poa annua control (0-100%) based on initial Poa cover in each plot Bentgrass quality (1-9, 1 = dead; 6 = minimally acceptable)

RESULTS:

- From May 13 to July 14, average minimum and maximum air temperatures were 59.1 F (45.7 F low) and 85.0 F (99.2 F high), respectively; average minimum and maximum relative humidity was 35% (8% low) and 80% (85% high), respectively; average soil temperatures ranged from 62.5 F to 76.0 F (73.3 F average); total CIMIS ETo was 2.78 inches; no precipitation.
- More than one application of amicarbazone applied at 2.0 oz/A or greater caused severe injury or death to bentgrass turf (Figure 1; Table 1). Preliminary results from studies in Northern California indicate that these rates can be safely applied in cooler climates or during cooler periods of the year, but they are too high for late spring/early summer in Riverside.
- Maximum air temperatures occurred between the July 6 and July 14 rating dates. Furthermore, the green was vertical mowed on July 7. The combination of both stresses resulted in the lowest overall bentgrass quality ratings on July 14, and plots treated with higher rates of HM 0814 began showing significant turf injury relative to the control.
- MRC-01 provided the best combination of Poa control and bentgrass safety during this phase of the experiment (Table 2); however, there appeared to a minimum total dosage required to achieve optimal control. Hence, the 3.0 oz/1,000 ft2 rate (surfactant vs. no surfactant) was added on to the treatment list later during the experiment.
- The Riverside study will continue along with ongoing studies on golf courses Northern California, and new studies to be initiated on golf courses in Southern California. Focus will be on refining application rates and frequencies of MRC-01, evaluation of tankmix partners with MRC-01 and with amicarbazone at 1.0 oz/A, and evaluation of higher rates and/or more frequent applications of other herbicides in this study to achieve maximum Poa control with minimal bentgrass injury.



FIGURE 1.
Bentgrass injury on
July 6 following four
weekly applications of
amicarbazone at 2 oz/A
from May 13 to June 4,
2010. Riverside, CA.

TABLE 1. Creeping bentgrass quality (1-9, 1=dead, 6=minimally acceptable) following application of herbicides, Riverside, CA, 2010.

	_								
Trt	Rate	Timing	5/20	5/28	6/6	6/19	6/25	7/6	7/14
Velocity	4 oz/A	Α	7.0	7.7	7.7	8.0	9.0	9.0	8.0
Velocity +	2 oz/A +	Α	7.0	8.0	8.0	8.0	9.0	9.0	8.0
Trimmit	8 oz/A	Α							
Amicarbazone	1 oz/A	Α	8.0	8.0	8.0	8.0	8.0	9.0	7.3
Amicarbazone	2 oz/A	Α	7.7	7.7	8.0	6.7	6.0	8.3	8.0
Amicarbazone	4 oz/A	Α	7.0	6.0	6.3	2.7	2.3	2.7	3.0
Amicarbazone +	2 oz/A +	Α	7.0	6.7	7.3	5.0	4.7	6.0	7.7
Trimmit	8 oz/A	Α							
MRC-01	1.25 oz/1,000ft ²	Α	8.0	8.0	8.0	9.0	9.0	8.7	7.0
Prograss	8 oz/A	Α	8.0	8.0	7.7	8.7	9.0	9.0	7.7
Prograss +	6 oz/A +	Α	8.0	8.0	8.0	7.7	7.0	8.7	7.7
Amicarbazone	2 oz/A	Α							
HM 0814	3 oz/1,000ft ²	Α	8.0	8.0	8.0	8.3	8.7	8.0	6.7
HM 0814	6 oz/1,000ft ²	Α	8.0	8.0	8.0	8.3	8.0	8.0	6.7
HM 0814 +	2 oz/1000ft ² +	Α	8.0	8.0	8.0	8.3	8.3	8.0	7.3
Trimmit	8 oz/A	Α							
Trimmit	10 oz/A	Α	8.0	8.0	8.0	8.0	8.3	9.0	8.3
Trimmit	16 oz/A	Α	8.0	8.0	8.3	9.0	8.3	9.0	8.7
Bensumec 4 LF	9.4 oz/1000ft ²	Е	8.0	8.0	8.0	8.7	9.0	9.0	7.0
SureGuard	6 oz/A	Е	8.0	8.0	8.0	9.0	9.0	9.0	7.7
Amicarbazone	1 oz/A	В	7.7	7.7	6.0	6.7	6.7	8.3	8.0
Amicarbazone	2 oz/A	В	7.3	6.0	2.0	1.7	1.3	2.7	3.7
Amicarbazone +	1 oz/A +	В	8.0	7.7	5.0	6.3	6.7	8.0	8.0
Trimmit	2 oz/A	В							
FeSO ₄ ¹	16 oz/1000ft ²	С	9.0	8.0	8.0	9.0	9.0	9.0	9.0
MRC-01	0.75 oz/1000ft ²	В	8.0	8.0	7.7	9.0	9.0	8.7	7.7
MRC-01	3 oz/1000ft ²	D	8.0	8.0	8.0	8.7	8.7	9.0	7.3
MRC-012	3 oz/1000ft ²	D	8.0	8.0	8.0	8.3	9.0	9.0	7.7
Untreated Control	-	-	8.0	8.0	8.0	9.0	9.0	9.0	8.0
LSD (0.05)*	0.3	0.4	0.6	0.8	1.0	1.0	1.1		

^{*} Treatment mean differences in columns greater than or equal to LSD are significantly different, Fisher's Protected LSD, P = 0.05.

TABLE 2. Annual bluegrass control (0-100%) following application of herbicides. Riverside, CA. 2010.

Trt	Rate	Timing	5/28	6/6	6/19	6/25	7/6	7/14	
Velocity	4 oz/A	Α	28	33	48	57	70	65	
Velocity + Trimmit	2 oz/A + 8 oz/A	A A	31	25	40	34	73	48	
Amicarbazone	1 oz/A	Α	29	44	57	60	47	64	
Amicarbazone	2 oz/A	Α	37	52	96	99	80	92	
Amicarbazone	4 oz/A	Α	70	87	100	100	100	100	
Amicarbazone + Trimmit	2 oz/A + 8 oz/A	A A	40	71	100	100	94	89	
MRC-01	1.25 oz/1,000ft ²	Α	20	61	84	75	82	87	
Prograss	8 oz/A	Α	12	32	52	44	36	37	
Prograss + Amicarbazone	6 oz/A + 2 oz/A	A A	38	45	90	88	78	68	
HM 0814	3 oz/1,000ft ²	Α	10	20	25	21	23	30	
HM 0814	6 oz/1,000ft ²	Α	21	18	54	48	51	54	
HM 0814 + Trimmit	2 oz/1,000ft ² + 8 oz/A	A A	19	9	31	22	52	63	
Trimmit	10 oz/A	Α	10	14	24	19	47	33	
Trimmit	16 oz/A	Α	18	24	32	28	51	62	
Bensumec 4 LF	9.4 oz/1,000ft ²	Е	4	18	51	30	39	54	
SureGuard	6 oz/A	Е	17	10	34	27	21	17	
Amicarbazone	1 oz/A	В	32	84	87	78	61	74	
Amicarbazone	2 oz/A	В	69	98	100	100	100	100	
Amicarbazone + Trimmit	1 oz/A + 2 oz/A	B B	21	82	100	97	87	76	
FeSO ₄ ¹	16 oz/1,000ft ²	С	20	23	26	28	49	40	
MRC-01	0.75 oz/1,000ft ²	В	32	86	97	98	76	74	
MRC-01	3 oz/1,000ft ²	D	10	20	50	45	56	98	
MRC-01 ²	3 oz/1,000ft ²	D	23	25	34	27	62	91	
Untreated Control	-	-	17	11	22	19	11	30	
LSD (0.05)*			21	23	21	24	29	29	

^{*} Treatment mean differences in columns greater than or equal to LSD are significantly different, Fisher's Protected LSD, P = 0.05.

ENTERPRISES, INC.

Phone: 408-432-9040

IN BUSINESS SINCE 1961

erving California Golf Courses For More Than Forty Years **GCSANC Members Since 1976**

USGA Root Zone Sand or Mixes **Green & Fairway Top Dressing Sand Divot Mixes & Planter Mixes** Bark, Compost, Peat, Coir, Sawdust, Mulch Gypsum & Lime in Bulk, Bags or Bulk Sacks

Bunker Sand of any Type, Color, Origin Kiln Dried Sand in Bulk or Bags Gravel, Drain Rock, Road Base, DG **Stabilized Cart Path Aggregates NEXPAVE Wax Coated Aggregates**

Proud Sponsor



Nationwide Tour

Manufacturer Direct Distributors for



Web: www.tmtenterprises.net Email: info@tmtenterprises.net

¹ FeSO₄ applied in 320 GPA of water; all other treatments applied in 80 GPA.

² No surfactant added; all other treatments applied with 0.25% non-ionic surfactant.

¹ FeSO₄ applied in 320 GPA of water; all other treatments applied in 80 GPA.

² No surfactant added; all other treatments applied with 0.25% non-ionic surfactant.