

Prepare for the worst

Steve and Suz Trusty, partners in Trusty & Associates based in Council bluffs, IA

Fields that enter winter in top shape have the best chance of emerging from winter stress in playable condition. Survival strategies include a well-managed, year-round maintenance program, and as much fall preparation as field use schedules, weather, time, and funding allow.

Ideally, baseball and softball fields should be completely ready for play when they are put to bed for the winter. Repair lip areas and basepaths; prep batter's boxes and pitching mound areas; and where turf covers are available, blanket skinned areas prior to heavy snow cover. When basepath covers aren't in the budget, it may help to erect an upright screen of landscape-type fabric to reduce wind and storm damage.

Whenever possible, aerate turf to reduce compaction and improve water infiltration and percolation. Timing of core aeration is critical. This is especially true for uncovered fields with high exposure to winter winds because of the desiccation factor. Take into consideration existing field conditions, soil profile, turfgrass varieties and cultivars, traditional winter and early-spring weather conditions, and reliability of long-range weather forecasts.

Remove excess thatch prior to the onset of winter to avoid disease infestation. But be careful, too little thatch can expose turfgrass crowns to desiccation, especially in areas with multiple freeze-thaw cycles. On fields subjected to winter play or other traffic, some thatch is needed to cushion turfgrass crowns from wear.

Lower mowing heights late in the season to reduce surface injury from traffic, and to reduce disease susceptibility. But again, be careful. Dropping the height of cut too early may reduce leaf surface enough to drop carbohydrate production to an unacceptable level. The height of cut and timing of height reduction depends on the turfgrasses involved, soil profile, field use, and off-season traffic. For late-season overseeding of cool-season turf, time your program based on whether you want germination during the current

season or early the following spring. Some sports turf managers make one late-season overseeding with a mix of turfgrasses to cover both needs. High-use, multi-sport, low- to medium-maintenance fields can be overseeded with turf-type tall fescues, perennial ryegrasses, and bluegrasses. The fescues and ryegrasses will germinate late, while bluegrass germination will follow in the spring.

Evaluate late-season turf fertility before setting a fertilization strategy. Provide adequate phosphorus for strong root development and support of new seedlings. Adequate potash will help overall turf vigor and provide resistance to stress.

Late-season nitrogen (N) applications depend on existing levels, turf development, types of turfgrasses, type of N, and weather conditions. Cool-season turf needs a period of lower N levels to harden off, but levels must be sufficient to support fall growth and to start spring green-up.

Apply preventive fungicide for snow molds on high-profile, high-maintenance fields where air movement is restricted by a stadium enclosure, snow cover, or tarping. Moisture and temperature conditions leading into winter dictate the number and timing of applications.

Tarps protect turfgrasses from snow and ice buildup, excess moisture, extreme cold, and desiccation. They retain warmth, keeping turf growing

longer into the fall and giving it an earlier start in the spring.

Sustained periods of above-normal temperatures can combine with lack of air movement to produce excess heat beneath a tarp. This promotes inappropriate turf growth and anaerobic decomposition in the thatch layer, and it creates favorable conditions for disease.

Temporarily increase air circulation across the turf to prevent these problems. You can lift a portion of the tarp to introduce air flow beneath it, or temporarily remove the tarp if conditions are severe enough to warrant it.

Drying winds moving across open fields may combine with lack of precipitation to produce damaging conditions. If temperatures are high enough to allow acceptable water penetration, irrigate to counteract the lack of moisture.

Check turf and soil conditions frequently as temperatures warm. If conditions are favorable for disease, consider alternatives to change one or more of the contributing factors. Increase air movement, reduce temperatures, or reduce moisture. If these conditions can't be changed and signs of disease are present, a preventive spray may be cost effective.

Consider overseeding these areas in spring to counteract turf loss. Match spring tarp removal to appropriate weather conditions: cool, cloudy, and still. Too much sun, heat, or wind can desiccate newly uncovered turf, especially when soil temperatures are too cool to allow roots to replace surface moisture loss. Be prepared to syringe the field lightly, but frequently, to avoid desiccation.

A competent, vigilant sports turf manager is the greatest factor in winter field survival. ♦



Count on it.

*Toro Irrigation
Athletic Field Specialists*

For sales and design assistance, contact:

Storr Tractor Co.
Mike Pastori
908-722-9830

Phila. Turf Co.
Rich Toleno
267-266-0389