
WEEDWATCH

STANDING SENTINEL TO PROTECT PLANT HEALTH



CAROLINA GERANIUM

Geranium carolinianum

IDENTIFICATION TIPS

- > This winter annual or biennial is often called crane's bill because of the shape of the fruit.
- > Leaves are deeply dissected and divided into five to nine toothed segments.
- As it matures, the basal rosette experiences stem elongation and branching.
- > The densely hairy stems are erect, usually pink to red in color.
- > Whitish-pink to purple flowers help differentiate this weed from smaller geranium species.

CONTROL TIPS

- > Fall applications are recommended. Hand-pull any existing geraniums.
- Apply a pre-emergent broadleaf herbicide labeled for use on your ornamental beds. Products containing isoxaben are effective, and are labeled for ornamental plant tolerance on more than 630 ornamental species.



BUCKHORN PLANTAIN

Plantago lanceolata

IDENTIFICATION TIPS

- > This perennial turfgrass weed is commonly found in maintained turf, agricultural land and other disturbed sites.
- > Leaves are footballshaped and spiral around a short stem, clustering around the base of the plant.
- It has narrower leaves and shorter flower stalks than broadleaf plantain (*Plantago major*).
- > Flower heads consist of leafless, slender spikes of inconspicuous flowers clustered densely along

the upright flowering head. Open flowers have protruding white stamens.

CONTROL TIPS

- In spring or fall, apply a post-emergent broadleaf herbicide with good translocation properties to thoroughly control the root system of this plant. Two- and three-way auxinic herbicide products provide thorough control.
- > Properly fertilize, mow and water to maintain lush turf growth and minimize the potential for buckhorn plantain to establish itself.
- * Confront is not for sale or use in Nassau and Suffolk counties, New York.

For more information regarding these and other turf weeds — and related control technologies and tips — please visit www.DowProvesIt.com or call 800/255-3726.

