

## Submergent Plants (Continued)




Floating pondweed (Potamogeton natans)—Has two types of leaves. Underwater leaves are narrow, grasslike, and appear as stalks. Floating leaves are oval to heart-shaped, each with notched base. Flowers and seeds on a spike.




Leafy pondweed (Potamogeton foliosus)—Leaves ribbon-like, about 1/16-inch wide, lack sheath at base.




Curly pondweed (Potamogeton crispus)—Leaves alternate, have finely toothed, crinkled, or puckered edges. No floating leaves. Flowers and seeds in spike at tip extending above water for fertilization. Grows in fertile hard water. Introduced from Europe.



Water-stargrass (Heteranthera dubia) —Looks like some narrow-leaved pondweeds (Potomogeton), but leaves lack a midvein. Flower yellow, star-like.




Sago pondweed (Potamogeton ped atus)—Leaves fine, thread-like, spread as a fan, have sheathed base. floating leaves. Stems usually m branched. Tubers grow from horizo roots.




Wild celery (Vallisneria americ—Light green, ribbon-like leaves ma as long as 6 feet (but usually much I with tips floating on surface.


## **Floating Plants**



White waterlily (Nymphaea odorata) —Round floating leaves grow to 10 inches diameter, split to stem at center, often purple on underside. Flowers showy, usually white but sometimes pink. Flowers open from morning until shortly after midday.



Duckweed (Lemna species)—Tiny, free-floating, bodies are flat and round or lobed, oatmeal-sized or smaller, often mistaken for algae. Barely-visible roots dangle thread-like. Sometimes several plants attached. Masses of this plant accumulate as a scum blanketing quiet shallows. As scum dies, it turns yellowish or whitish.



Lotus (Nelumbo lutea)—Round fing leaves grow to 24 inches widepression in center where stem atted. Leaf veins radiate from cellowers yellow, large, showy.