

1986 Michigan Soybean Performance Report

Extension Bulletin E-1206 January 1987

By

**M.L. Vitosh, T.G. Isleib, R.H. Leep,
J.L. Lockwood, D.E. Wolfe and L. Rood-Kao**

Department of Crop and Soil Sciences
Department of Botany and Plant Pathology

This bulletin provides information on the performance of soybean varieties available in Michigan.

Comprehensive variety yield trials were conducted in Southeastern Michigan (Lenawee County), Southwestern Michigan (St. Joseph County), Far Southwestern Michigan (Berrien County), South Central Michigan (Ingham County), Central Michigan (Saginaw County), and East Central Michigan (Sanilac County). Smaller trials were conducted in Huron and Alger Counties.

Testing Procedures

Commercial varieties voluntarily entered were obtained from seed companies. Public varieties were supplied by the Michigan Foundation Seed Association.

Cooperators, planting and harvest dates, fertilizer practices, previous crops, and soil management groups at the seven locations are listed in Table 1.

Maturity groups of all varieties tested are listed in tables 2 and 5. Seed of entries was planted in plots 20 feet long with a 20-inch row spacing. Seeds were planted 1½ inches deep at 4.5 seeds per foot of row. Each plot was randomized in the field and replicated 3 times. Fourteen feet of the center two rows were harvested for yield.

Evaluation of Characteristics

YIELD—Yield is expressed in bushels per acre at 13% moisture.

MATURITY DATE—Entries were considered mature when 95% of the pods had attained their final color and would crack under finger pressure. Additional field drying was required before the plants were ready to harvest. Dates were recorded by month and day.

HEIGHT—Plant height, in inches, was measured at maturity from the soil surface to the tip of the main stem.


LODGING—Lodging rates reflect the erectness of the plants before harvest. Ratings are based on the following scale:

1. Almost all plants erect

2. All plants leaning slightly, or fewer than 25% of the plants down
3. All plants leaning moderately (45%), or 25% to 50% of the plants down
4. All plants leaning considerably, or 50% to 80% of the plants down
5. Almost all plants down

Results

Tables 3-6 show results of 1986 soybean variety trials. Table 7 is 1985 data from Alger County in the Upper Peninsula. Values given are the averages of all replications harvested at each location. Extremely rainy weather in September adversely affected the plots at Saginaw, Sanilac and Huron counties.



Soybean Maturity Zones for Full-Season Varieties in Michigan, and Locations (*) of Trials.

The LSD (least significant difference) value is useful when comparing two varieties in the same table. Two varieties with the same genetic potential for yield may have different yields due to variation in soil fertility, compaction, and other environmental factors. If the difference is less than the LSD value, the difference between the varieties may be due to chance or minor environmental differences. However, if the difference between two varieties is greater than the LSD, there is a 95%, or better, probability that the performance is actually different. The CV value is an indicator of the degree of precision for a particular test. The lower the CV value, the more discerning the test.

Selecting a Variety

The primary consideration in selecting a variety is yield. When evaluating a variety, consider yield performance over several years, if available. Give preference to data obtained in the nearest variety trial. Use all trials in determining a variety's performance under various environmental conditions.

Considerations other than yield are important in selecting a variety, and in some cases result in choosing a variety with only moderate performance. It is especially important to select a variety with proper maturity. From past weather data, farmers can determine the percent probability of the first fall frost. A general rule of thumb is to choose a variety that will mature (see maturity date definition) before the average date for 25% chance of the first killing frost in the fall. Farmers growing soybeans for the first time may wish to contact neighbors to determine what varieties mature before frost in their area. When large acreages of soybeans are planted, varieties of different maturities provide staggered maturity dates for a longer harvest season.

The degree of lodging varies among varieties. Lodged plants in variety trials are manually picked up and threshed, thus yield losses from lodging are not reflected in the yields reported. Lodging ratings should be used to evaluate potential losses. Farmers who have experienced lodging in the past and have had harvest problems may select a more lodging-resistant variety. Alternately, a variety susceptible to lodging may be planted at a slightly lower population to increase standability. Evaluate lodging data over all locations to determine a variety's lodging characteristics.

Note seed size when selecting planting rates. Planting rates should be based on number of seeds per foot of row and not on pounds per acre.

Many diseases occur in soybean fields in Michigan. The diseases which contribute most significantly to yield reduction are seed and seedling diseases and those causing root and stem rot. Root rots of soybeans are generally recognized when plants turn yellow prematurely, wilt, or die. Less noticeable is the yield reduction that occurs when root rot destroys part of the root system, but causes no visible symptoms to above-ground parts. The fungi

that cause root rots often survive in the soil for several years, even in the absence of a host plant. Once root rot fungi are established in a field, control is difficult, even with crop rotation.

New varieties with resistance to one or more diseases are being developed, particularly varieties resistant to *Phytophthora* root rot. Disease resistance characteristics to *Phytophthora* root rot are noted in Table 2, but the large number of races of this fungus complicates variety selection.

Sclerotinia stem rot (white mold) studies were conducted at East Lansing in 1984 and 1985 with 20 and 16 varieties, respectively. The disease was encouraged by irrigating with 1.5 to 2 inches of water weekly beginning at flowering until the end of August. The disease ratings, expressed as percentage of plants infected, are shown in Table 2. The disease ratings for 13 of the varieties grown both years were positively correlated. Yields were measured in 1985 and the data showed that for every 10 percent increase in disease, yield was reduced 3.5 bushels per acre. The maximum yield in the study was 45 bushels per acre.

It is often beneficial for growers to select a few good varieties for planting each year. Yield determination and careful field evaluation during the growing season will add to the grower's knowledge of varietal performance and allow better selection.

More information about variety selection and cultural practices can be found in Extension Bulletin E-1549, "Soybean Production in Michigan."

Use of Data

Table 3 presents multiple-environment averages from all tests in the Southern and Central Michigan regions since 1975. The column labeled N refers to the number of tests in which each variety was included. The column labeled DEV. refers to the difference (in bushels per acre) between the mean yield of the variety over N tests and the mean yield of all varieties in those tests. The maturity checks used for tests of Group I and Group II varieties were "Hodgson 78" (H78) and "Corsoy 79" (C79), respectively. A positive relative maturity value means that the variety matured later than the check and a negative value means that the variety matured earlier than the check. The value is the actual number of days in either direction.

Data presented in Tables 4 through 6 are from both regional and site-specific performance trials. Both 1986 yields and multiple-year average yields from all tests since 1975 are given. Maturity, height (in inches), and lodging scores are the 1986 regional averages. Maturity is expressed as + or - days when compared with the check variety. For 1986 yield data, all starred entries designate yields not significantly different from the highest yield for that location. Multiple-environment and multiple-year averages comprised of a greater number of tests (greater N) should be considered more reliable.

The presentation of data for the entries tested does not

suggest approval or endorsement of varieties by the authors or by those responsible for conducting the performance trials.

The MSU Cooperative Extension Service approves the reproduction of the information in this publication only if

no portion is deleted, if the data are not rearranged or otherwise manipulated, and if appropriate credit is given.

TABLE 1. Variety Trial Information.

| County | Lenawee | St. Joseph | Ingham | Saginaw | St. Clair | Sanilac | Huron |
|----------------------------------|-------------------------------|------------------------------|--|--------------------------------|--|----------------------------------|---|
| CES Director/ Agent | N.H. Bless G.A. Wuethrich | F.J. Henningsen D. Bowen | M.M. Preston R.A. Morrison | H.R. Ferris S.S. Poindexter | M.F. Hansen T.R. Johnson | A.R. Sieting R.C. Weber | R.A. Johnson J.L. LeCureux |
| Farmer Cooperator | D. Woods | B. Marantette J. Sheppard | | C. Gosen | R.A. Greenia | K. Pritchett | H. Hass |
| Address | 10992 Holloway Britton, MI | 25660 Simpson Mendon, MI | MSU Campus East Lansing, MI | 8735 Swan Creek Saginaw, MI | 1395 Kronner Richmond, MI | 2985 S. Sandusky Sandusky, MI | 1001 Learman Bad Axe, MI |
| Soil Type | Lenawee silty clay loam | Elston sandy loam | Aubbeenaubbee- Capac and Riddles-Hillsdale sandy loam | Colwood silt loam | Wasepi sandy loam and Bayer loamy sand | Capac and Parkhill loam | Corunna sandy loam and Kilmanagh loam |
| Soil Management Group | 1.5 c | 4 a | 3.2 b/2.5 b and 2.5 a/3 a | 2.5 c-s | 4 a and 4 b | 2.5 b and 2.5 c | 3/2 c and 2.5 c |
| Previous Crop | Sod | Corn | Corn | Soybeans | Wheat | Soybeans | Corn |
| Fertilizer | 100# K ₂ O | | 200# 6-24-24 | 200# 8-18-36 | 200# K ₂ O | 290# 8-23-10 | 250# 10-26-26 |
| Planting Date | 5/10/86 | 5/29/86 | 6/18/86 | 5/30/86 | 6/2/86 | 6/2/86 | 5/27/86 |
| Harvest Date | 10/11/86 | 10/9/86 | 10/31/86 | 10/23/86 | 10/10/86 | 11/13/86 | 10/21/86 |

TABLE 2. Reactions of Soybean Cultivars to Sclerotinia Stem Rot (White Mold) in 1984-85 Field Tests.

| Cultivar | Maturity Group | Disease Incidence (%) | |
|------------|----------------|-----------------------|------|
| | | 1984 | 1985 |
| GNOME | II | 39.5 | 52.4 |
| WEBER 84 | I | 31.1 | 48.5 |
| SPRITE | III | — | 28.8 |
| CENTURY | II | 19.1 | 28.1 |
| ELGIN | II | 15.5 | 21.2 |
| VICKERY | II | 13.8 | — |
| AMSOY 71 | II | 13.7 | — |
| AMCOR | II | 13.4 | — |
| CORSOY | II | — | 7.9 |
| CORSOY 79 | II | 13.3 | 5.3 |
| LAKOTA | I | 13.0 | — |
| BEESON 80 | II | 12.6 | 18.0 |
| HARCOR | II | 12.4 | — |
| WELLS II | II | 11.9 | 23.1 |
| NEBSOY | II | 10.0 | 29.3 |
| HODGSON 78 | I | 6.8 | 2.8 |
| PELLA | III | 6.7 | 3.8 |
| SIMPSON | 0 | 6.1 | — |
| HARDIN | I | 5.7 | 7.1 |
| HOBBIT | III | 4.1 | 40.4 |
| DAWSON | 0 | 2.8 | — |
| EVANS | 0 | 2.2 | 12.0 |
| OZZIE | 0 | — | 0.0 |
| LSD (.05) | | 12.4 | 13.6 |

*Percentage of diseased plants in two rows 14 feet long.

TABLE 3. Performance Summary for Varieties Entered in the Michigan Trials in 1986. Phytophthora Resistance Designations Denote the Following: Type 1A Resistant to Races 1, 2, and 10; Type 1B Resistant to Races 1 and 3-9; Type 1C Resistant to Races 1-3 and 6-10; Type 1K Resistant to Races 1-10; Type 3 Resistant to Races 1-5, 8, and 9; Type 6 Resistant to Races 1-4 and 10.

| BRAND | ENTRY | MG | PHYT. RES. | YIELD (BU/A) WITH DEVIATION FROM MEAN | | | | MATURITY RELATIVE TO CHECKS | | | | | | |
|--------------------|-----------|------|---------------|---------------------------------------|--------------|---------|--------------|-----------------------------|------|-------|---------|------|---------------|-----|
| | | | | SOUTHERN | | CENTRAL | | SOUTHERN | | | CENTRAL | | LODGING SCORE | |
| | | | | TYPE | YIELD (N) | DEV. | YIELD (N) | DEV. | DATE | H78 | C79 | DATE | H78 | C79 |
| PUBLIC | | | | | | | | | | | | | | |
| DASSEL | O 6 | -- | - | - | 36.3 | (7) | -2.3 | -- | - | 9-29 | 0 | -7 | - | 1.3 |
| DAWSON | O 1A | -- | - | - | 40.5 | (19) | -0.7 | -- | - | 9-23 | -5 | -12 | - | 1.6 |
| EVANS | O 1A | -- | - | - | 36.8 | (31) | -2.5 * | -- | - | 9-21 | -4 | -12 | - | 1.5 |
| OZZIE | O 1A | -- | - | - | 39.0 | (17) | -3.7 * | -- | - | 9-19 | -8 | -15 | - | 1.2 |
| SIMPSON | O 1A | -- | - | - | 38.5 | (15) | -3.5 * | -- | - | 9-23 | -4 | -11 | - | 1.3 |
| BSR 101 | I 1A | 46.4 | (11) | 2.0 * | 41.4 | (13) | 2.0 * | 9-22 | 3 | -4 | 10-5 | 7 | -1 | 1.6 |
| HARDIN | I 1A | 42.7 | (17) | 1.1 | 45.2 | (22) | 3.8 * | 9-22 | 3 | -3 | 10-2 | 4 | -3 | 2.2 |
| HODGSON 78 | I 1A | 40.3 | (26) | -1.0 | 41.3 | (33) | 0.0 | 9-19 | 0 | -6 | 9-28 | 0 | -6 | 1.9 |
| SIBLEY | I 1A | 49.3 | (8) | 4.4 * | 40.8 | (10) | 1.2 | 9-18 | 0 | -6 | 9-30 | 0 | -7 | 2.1 |
| WEBER 84 | I 1A | 40.2 | (13) | -1.8 | 38.8 | (15) | -0.5 | 9-23 | 4 | -3 | 10-2 | 4 | -3 | 2.7 |
| AMCOR | II 1A | 41.8 | (17) | -0.6 | 42.3 | (17) | 0.0 | 9-29 | 10 | 3 | 10-9 | 11 | 4 | 2.9 |
| BEESON 80 | II 1C | 38.1 | (17) | -4.3 * | 39.2 | (19) | -2.6 | 9-27 | 8 | 2 | 10-9 | 10 | 3 | 1.9 |
| BSR 201 | II 1B | 47.4 | (14) | 1.3 | 42.2 | (12) | 0.3 | 9-26 | 8 | 1 | 10-8 | 11 | 2 | 2.6 |
| CENTURY | II 1A | 44.2 | (18) | 1.4 | 42.3 | (20) | 0.8 | 9-29 | 10 | 3 | 10-9 | 10 | 3 | 1.8 |
| CENTURY 84 | II 1K | 42.1 | (9) | -0.5 | 41.4 | (9) | 1.4 | 9-29 | 9 | 2 | 10-10 | 12 | 3 | 1.5 |
| CORSOY | II NONE | 41.4 | (26) | 0.7 | 39.4 | (29) | 0.3 | 9-24 | 5 | -1 | 10-2 | 6 | 0 | 2.3 |
| CORSOY 79 | II 1C | 42.7 | (24) | 1.0 | 43.4 | (30) | 2.2 * | 9-25 | 6 | 0 | 10-5 | 7 | 0 | 2.5 |
| ELGIN | II NONE | 44.5 | (15) | 1.4 | 43.8 | (16) | 2.7 * | 9-25 | 6 | -1 | 10-5 | 6 | -1 | 2.0 |
| HACK | II 1A | 45.3 | (10) | 1.5 | 41.7 | (10) | 1.6 | 9-28 | 8 | 2 | 10-8 | 10 | 2 | 1.4 |
| HOYT | II 1A | 48.1 | (7) | 2.4 | 39.6 | (8) | -0.1 | 9-27 | 9 | 4 | 10-9 | 11 | 3 | 1.6 |
| KELLER | II 1C, 3 | 39.5 | (8) | -2.1 | 40.0 | (8) | -0.4 | 9-29 | 8 | 1 | 10-10 | 11 | 2 | 2.2 |
| MIAMI | II 1C, 3 | 38.7 | (9) | -3.9 * | 38.5 | (9) | -1.6 | 9-24 | 4 | -3 | 10-6 | 7 | -1 | 1.8 |
| NEBSOY | II 1A | 41.8 | (18) | -1.1 | 40.6 | (20) | -1.0 | 9-24 | 5 | -1 | 10-6 | 7 | 0 | 1.6 |
| PRESTON | II NONE | 47.0 | (6) | 1.6 | 38.5 | (7) | 0.0 | 9-28 | 10 | 4 | 10-9 | 11 | 3 | 2.5 |
| VICKERY | II 1C | 42.6 | (18) | -0.2 | 42.1 | (21) | 0.9 | 9-24 | 5 | -2 | 10-4 | 6 | -1 | 2.9 |
| WELLS II | II 1C | 40.7 | (20) | -2.1 * | 40.2 | (22) | -0.7 | 9-24 | 4 | -2 | 10-4 | 5 | -1 | 1.6 |
| CUMBERLAND | III NONE | 36.8 | (13) | -3.2 | -- | - | - | 10-3 | 13 | 7 | -- | - | - | 2.5 |
| GNOme 85 | III 1K | 47.8 | (10) | -0.3 | -- | - | - | 9-25 | 10 | 4 | -- | - | - | 1.4 |
| HOBBIT | III NONE | 40.0 | (15) | 0.8 | -- | - | - | 9-30 | 12 | 5 | -- | - | - | 1.3 |
| PELLA | III 1A | 40.2 | (13) | 0.3 | 43.4 | (13) | 2.2 | 10-1 | 11 | 4 | 10-9 | 11 | 4 | 1.7 |
| SPRITE | III NONE | 39.9 | (16) | -0.1 | -- | - | - | 10-1 | 13 | 7 | -- | - | - | 1.7 |
| WILLIAMS 82 | III 1K | 39.0 | (10) | -1.6 | -- | - | - | 10-5 | 14 | 8 | -- | - | - | 2.8 |
| WINCHESTER | III 1B, 3 | 38.9 | (7) | -2.2 | -- | - | - | 10-6 | 14 | 6 | -- | - | - | 2.9 |
| ZANE | III NONE | 41.5 | (7) | 0.4 | 39.5 | (7) | 0.6 | 10-2 | 10 | 3 | 10-11 | 12 | 4 | 2.1 |
| AGRIPRO | | | | | | | | | | | | | | |
| AP200 | II 1A | 42.6 | (16) | -0.3 | 43.4 | (18) | 1.1 H | 9-23 | 2 | -4 | 10-2 | 4 | -3 | 2.2 |
| AP240 | II 1A | 41.0 | (15) | 0.2 | 42.0 | (8) | -1.3 | 9-27 | 8 | 1 | 10-6 | 6 | 0 | 1.7 |
| AP2190 | II 1A | 40.9 | (7) | -0.2 H | 40.0 | (7) | 1.0 | 9-30 | 8 | 0 | 10-8 | 9 | 1 | 2.1 |
| AP3023 | III 1A | 41.1 | (3) | 0.7 | -- | - | - | 10-3 | 11 | 5 | -- | - | - | 2.5 |
| AP3132 | III NONE | 40.8 | (3) | 0.4 | -- | - | - | 10-2 | 10 | 4 | -- | - | - | 1.9 |
| HP20-20 | II 1A | 43.4 | (11) | -1.1 | 40.6 | (8) | -2.9 | 9-23 | 3 | -4 | 10-4 | 5 | -3 | 2.2 |
| HP2530 | II 1A | 44.0 | (14) | 1.4 | 44.5 | (9) | 1.6 | 9-28 | 7 | 1 | 10-8 | 7 | 2 | 2.1 |
| EX 2021 | II 1A | -- | - | - | 38.4 | (4) | 1.7 | -- | - | 10-8 | 6 | -1 | - | 2.1 |
| ASGROW | | | | | | | | | | | | | | |
| A0949 | O 1C | -- | - | - | 32.9 | (5) | -2.5 | -- | - | 10-3 | 1 | -6 | - | 1.6 |
| A1525 | I 1A | -- | - | - | 37.0 | (9) | -0.9 | -- | - | 10-1 | 1 | -6 | - | 1.3 |
| A1937 | I 1A | 42.6 | (16) | 1.5 | 44.7 | (19) | 3.4 * | 9-21 | 2 | -4 | 10-1 | 3 | -4 | 2.0 |
| A2187 | II 1A | 40.9 | (7) | -0.2 | 41.2 | (4) | 0.8 | 9-26 | 3 | -4 | 10-6 | 5 | -4 | 1.6 |
| A2522 | II NONE | 41.7 | (11) | 0.8 H | 44.1 | (6) | 2.3 * | 9-29 | 8 | 1 | 10-8 | 7 | 2 | 2.5 |
| A2943 | II 1A | 43.8 | (11) | 2.9 * | 42.5 | (3) | -0.1 | 10-4 | 12 | 5 | 10-15 | 13 | 3 | 2.2 |
| CALLAHAN | | | | | | | | | | | | | | |
| 1250 BRAND | II NONE | 41.0 | (13) | 1.1 | 44.1 | (11) | 2.9 | 9-30 | 11 | 4 | 10-11 | 13 | 5 | 2.0 |
| 5300 BRAND | III NONE | 43.2 | (7) | 2.1 | -- | - | - | 10-6 | 14 | 6 | -- | - | - | 2.7 |
| 5350 BRAND | III NONE | 42.6 | (11) | 1.7 | 42.1 | (3) | -0.5 | 10-4 | 12 | 5 | 10-14 | 13 | 3 | 2.8 |
| 6180X BRAND | I NONE | -- | - | - | 42.0 | (8) | 3.1 | -- | - | 10-2 | 2 | -6 | - | 2.9 |
| 6220X BRAND | II 1A | -- | - | - | 41.4 | (7) | 2.5 * | -- | - | 10-10 | 12 | 3 | - | 1.4 |
| 6262X BRAND | II 1A | 41.4 | (7) | 0.3 | 42.2 | (7) | 3.3 * | 10-1 | 9 | 1 | 10-11 | 12 | 3 | 1.8 |
| 7244X BRAND | II NONE | 43.9 | (3) | 3.5 | 39.8 | (4) | 3.0 | 10-1 | 9 | 3 | 10-12 | 10 | 3 | 1.9 |
| 7260X BRAND | II NONE | 43.5 | (3) | 3.1 | 40.4 | (4) | 3.7 *H | 9-29 | 7 | 1 | 10-12 | 10 | 3 | 1.1 |
| 7272X BRAND | II NONE | 43.3 | (3) | 2.9 H | 37.2 | (4) | 0.4 | 10-1 | 9 | 3 | 10-12 | 10 | 3 | 2.1 |
| COUNTRYMARK | | | | | | | | | | | | | | |
| FFR 112 | II NONE | 40.2 | (3) | -0.2 | 35.7 | (4) | -1.0 | 9-28 | 6 | 0 | 10-10 | 9 | 2 | 1.4 |
| FFR 226 | II NONE | 35.4 | (5) | -2.6 * | 33.5 | (4) | -3.2 | 9-27 | 10 | 4 | 10-13 | 12 | 4 | 1.9 |
| FFR 241 | II NONE | 40.0 | (3) | -0.4 | 37.0 | (4) | 0.3 | 9-30 | 8 | 2 | 10-12 | 10 | 3 | 1.6 |
| DAIRYLAND | | | | | | | | | | | | | | |
| DSR-120 | I NONE | -- | - | - | 41.7 | (20) | -0.1 L | -- | - | 9-26 | -2 | -9 | - | 1.7 |
| DSR-128 | I 1C | -- | - | - | 39.8 | (8) | 1.0 | -- | - | 9-29 | -1 | -9 | - | 1.2 |
| DSR-135 | I 1A | -- | - | - | 37.2 | (10) | 0.0 | -- | - | 9-30 | 0 | -8 | - | 1.7 |
| DSR-171 | I NONE | 42.8 | (18) | 0.9 | 43.1 | (21) | 1.3 H | 9-23 | 4 | -2 | 10-3 | 5 | -2 | 2.1 |
| DSR-212 | II NONE | 39.2 | (15) | -1.6 | 41.9 | (12) | -1.0 | 9-26 | 6 | 0 | 10-7 | 7 | 1 | 1.4 |
| DSR-255 | II 1C | 37.9 | (7) | -3.2 * | 35.4 | (7) | -3.5 * | 9-29 | 7 | -1 | 10-8 | 9 | 1 | 1.9 |
| DSR-287 | II NONE | 42.3 | (7) | 1.2 | -- | - | - | 10-4 | 12 | 4 | -- | - | - | 2.2 |
| DSR-297 | II 1C | 40.4 | (7) | -0.7 | -- | - | - | 10-5 | 13 | 6 | -- | - | - | 1.9 |
| DSR-317 | III NONE | 40.5 | (7) | -0.6 | -- | - | - | 10-6 | 14 | 7 | -- | - | - | 2.7 |
| DST-1103 | I NONE | -- | - | - | 37.0 | (5) | 1.6 | -- | - | 10-1 | -1 | -7 | - | 2.1 |
| DST-1207 | I 1C | -- | - | - | 35.9 | (5) | 0.4 | -- | - | 10-4 | 2 | -5 | - | 2.2 |
| DST-2203 | II NONE | 37.6 | (3) | -2.8 L | 33.2 | (4) | -3.5 L | 10-2 | 10 | 4 | 10-12 | 11 | 4 | 1.4 |
| DST-2204 | II NONE | 33.3 | (3) | -7.1 | 32.4 | (4) | -4.3 | 10-2 | 10 | 4 | 10-13 | 12 | 4 | 1.7 |

(CONT'D.)

* STATISTICALLY SIGNIFICANT DEVIATION ($P<.05$)

H VARIETY EXHIBITS HIGHER THAN AVERAGE RESPONSE TO HIGHLY PRODUCTIVE ENVIRONMENTS.

L VARIETY EXHIBITS LOWER THAN AVERAGE RESPONSE TO HIGHLY PRODUCTIVE ENVIRONMENTS.

TABLE 3. (Continued) Performance Summary for Varieties Entered in the Michigan Trials in 1986. Phytophthora Resistance Designations Denote the Following: Type 1A Resistant to Races 1, 2, and 10; Type 1B Resistant to Races 1 and 3-9; Type 1C Resistant to Races 1-3 and 6-10; Type 1K Resistant to Races 1-10; Type 3 Resistant to Races 1-5, 8, and 9; Type 6 Resistant to Races 1-4 and 10.

| BRAND | ENTRY | MG | PHYT. RES. TYPE | YIELD (BU/A) WITH DEVIATION FROM MEAN | | | | | | MATURITY RELATIVE TO CHECKS | | | | | | LODGING SCORE | |
|---------------------------|---------------|-----|-----------------------|---------------------------------------|------|--------------|---------|------|--------|-----------------------------|------|-----|---------|------|-----|---------------|---------|
| | | | | SOUTHERN | | | CENTRAL | | | SOUTHERN | | | CENTRAL | | | SOUTH. | CENTRAL |
| | | | | YIELD (N) | DEV. | YIELD (N) | DEV. | DATE | H78 | C79 | DATE | H78 | C79 | DATE | H78 | C79 | |
| DEKALB-PFIZER | | | | -- | - | 39.4 | (13) | 0.5 | -- | - | 10-5 | 7 | 0 | - | - | 1.7 | |
| | CX174 | I | NONE | 40.7 | (5) | 0.2 | 38.6 | (5) | -0.2 | 10-6 | 9 | 1 | 10-11 | 12 | 3 | 2.1 | 1.9 |
| | CX265 | II | 1C | 42.3 | (11) | 1.4 | 45.4 | (3) | 2.8 L | 10-1 | 10 | 3 | 10-13 | 12 | 2 | 2.4 | 2.3 |
| | CX283 | II | NONE | 38.4 | (3) | -2.0 | -- | - | 10-5 | 13 | 7 | -- | - | - | 1.7 | - | |
| | CX326 | III | 1C | -- | - | - | -- | - | -- | -- | -- | -- | -- | -- | -- | -- | |
| DIEHL FIELDS | EXP 101 | I | | 45.6 | (3) | 5.2 | 42.2 | (4) | 5.4 L | 9-25 | 3 | -3 | 10-6 | 4 | -3 | 1.6 | 1.5 |
| FUNK | G3145 BLEND | I | 1A | 42.2 | (3) | -0.0 | 40.6 | (13) | 1.8 * | 10-8 | 5 | -2 | 10-4 | 5 | -2 | 1.9 | 2.0 |
| | G3180 | I | NONE | -- | - | - | 38.8 | (8) | 1.0 | -- | - | - | 10-2 | 2 | -6 | - | 2.1 |
| | G3232 | II | NONE | 42.7 | (3) | 2.3 | -- | - | 10-3 | 11 | 5 | -- | -- | -- | -- | 2.6 | - |
| | G3239 BLEND | II | NONE | 42.3 | (8) | 0.8 | 42.3 | (6) | 0.6 | 10-4 | 12 | 4 | 10-9 | 8 | 3 | 2.4 | 2.0 |
| | 12283 | I | NONE | -- | - | - | 35.2 | (5) | -0.2 | -- | - | - | 10-4 | 2 | -5 | - | 1.6 |
| GOLDEN HARVEST | H-1233 BRAND | II | NONE | 42.6 | (5) | 2.1 | 41.5 | (7) | 2.6 * | 10-4 | 7 | -1 | 10-7 | 9 | 0 | 2.1 | 1.8 |
| | H-1265 BRAND | II | 1B | -- | - | - | 37.3 | (4) | 0.6 | -- | - | - | 10-12 | 10 | 3 | - | 2.1 |
| | H-1285 BRAND | II | NONE | 41.5 | (5) | 0.9 | 42.4 | (7) | 3.5 | 10-8 | 11 | 4 | 10-12 | 13 | 4 | 2.1 | 1.9 |
| GREAT LAKES HYBRIDS (GLH) | GL1434 BRAND | I | 1A | -- | - | - | 33.8 | (9) | -4.1 * | -- | - | - | 10-2 | 2 | -5 | - | 1.7 |
| | GL1900 BRAND | I | 1A | -- | - | - | 37.6 | (9) | -0.3 | -- | - | - | 10-8 | 7 | 0 | - | 1.7 |
| | GL1937 BRAND | I | 1A | 41.5 | (3) | -0.7 | 41.4 | (16) | 1.0 | 10-5 | 2 | -4 | 10-3 | 5 | -2 | 2.2 | 2.0 |
| | GL2634 BRAND | II | NONE | 45.5 | (16) | 4.0 * | 44.6 | (13) | 3.3 *H | 9-29 | 10 | 4 | 10-7 | 9 | 2 | 2.0 | 2.0 |
| | GL2908 BRAND | II | NONE | 43.0 | (11) | 2.0 | 41.3 | (3) | -1.3 | 10-4 | 13 | 6 | 10-15 | 14 | 4 | 2.4 | 2.5 |
| | XP1909 | II | 1A | -- | - | - | 38.3 | (4) | 1.6 | -- | - | - | 10-9 | 8 | 0 | - | 2.2 |
| | XP2206 | II | 1A | -- | - | - | 40.4 | (4) | 3.7 | -- | - | - | 10-13 | 12 | 4 | - | 2.0 |
| | XP2478 BRAND | II | NONE | 42.9 | (3) | 2.6 | 39.0 | (4) | 2.3 | 9-30 | 8 | 2 | 10-11 | 10 | 2 | 3.0 | 2.1 |
| | XP2633 BRAND | II | NONE | -- | - | - | 35.4 | (4) | -1.4 | -- | - | - | 10-14 | 13 | 6 | - | 2.2 |
| | XP2737 BRAND | II | NONE | -- | - | - | 36.7 | (4) | 0.0 | 10-1 | 9 | 3 | 10-12 | 10 | 3 | 1.6 | 2.2 |
| JACQUES | J-201 | II | 1A | 41.6 | (3) | 1.3 | 38.8 | (4) | 2.1 H | 9-25 | 3 | -3 | 10-6 | 5 | -2 | 1.3 | 1.7 |
| | J-231 | II | 1A | 43.8 | (11) | 2.9 * | 46.9 | (3) | 4.3 | 9-29 | 8 | 1 | 10-12 | 11 | 1 | 2.0 | 2.4 |
| | E85092 | I | 1A | -- | - | - | 35.3 | (5) | -0.1 | -- | - | - | 10-2 | 0 | -7 | - | 1.8 |
| KING GRAIN | KG60 | O | 1A | 39.3 | (3) | -2.9 | 36.2 | (13) | -2.6 | 9-27 | -6 | -13 | 9-25 | -4 | -11 | 1.2 | 1.7 |
| | KG70 | I | 1A | 43.2 | (5) | -0.1 | 39.0 | (18) | -1.8 | 9-27 | -1 | -6 | 9-29 | 1 | -6 | 1.7 | 1.7 |
| | KG80 | II | 1A | -- | - | - | 38.7 | (7) | -0.2 | -- | - | - | 10-5 | 6 | -2 | - | 2.3 |
| | KG81 | II | 1C | -- | - | - | 40.1 | (4) | 3.4 | -- | - | - | 10-6 | 4 | -3 | - | 1.7 |
| | KG90 | II | 1A | 44.9 | (3) | 2.3 | 42.4 | (11) | 2.4 * | 10-9 | 8 | -2 | 10-7 | 8 | 1 | 2.2 | 2.2 |
| LAKESIDE STATES | MILLLER BRAND | III | NONE | 43.3 | (7) | 2.2 | -- | - | 10-6 | 13 | 6 | -- | - | - | 2.4 | - | |
| | 15 BRAND | I | NONE | -- | - | - | 39.0 | (5) | 3.6 | -- | - | - | 10-5 | 3 | -4 | - | 1.8 |
| | 21 BRAND | II | NONE | 42.5 | (3) | 2.1 | 39.9 | (4) | 3.2 | 10-3 | 11 | 5 | 10-13 | 12 | 4 | 1.8 | 1.6 |
| | 22 BRAND | II | 1A | -- | - | - | 38.0 | (4) | 1.2 | -- | - | - | 10-12 | 10 | 3 | - | 2.0 |
| | 24 BRAND | II | 1A | -- | - | - | 40.0 | (4) | 2.8 H | -- | - | - | 10-11 | 10 | 2 | - | 1.8 |
| | 30 BRAND | III | NONE | 37.5 | (3) | -2.9 | -- | - | 10-4 | 12 | 6 | -- | - | - | 3.0 | - | |
| | 36 BRAND | III | | 38.5 | (3) | -1.8 | -- | - | 10-5 | 13 | 7 | -- | - | - | 1.5 | - | |
| | 106 BRAND | II | 1A | -- | - | - | 35.0 | (4) | -1.7 | -- | - | - | 10-12 | 11 | 3 | - | 2.4 |
| | 116 | I | 1A | -- | - | - | 38.3 | (5) | 2.9 * | -- | - | - | 10-9 | 7 | 0 | - | 1.6 |
| | 125 | I | 1A | 41.5 | (3) | 1.1 | -- | - | 9-29 | 7 | 1 | -- | - | - | 1.8 | - | |
| MAUMEE VALLEY | CALIBER | I | 1A | 41.5 | (11) | 0.6 | -- | - | 9-27 | 5 | -1 | -- | - | - | 2.5 | - | |
| | ENTERPRISE | II | NONE | 39.2 | (7) | -1.9 | -- | - | 10-2 | 10 | 3 | -- | - | - | 2.1 | - | |
| | KODIAK | III | NONE | 41.2 | (7) | 0.1 | -- | - | 10-5 | 13 | 5 | -- | - | - | 2.9 | - | |
| | MV-2E1 | II | 1A | 41.9 | (11) | 1.0 | -- | - | 10-2 | 11 | 4 | -- | - | - | 2.1 | - | |
| | WARRIOR | II | 1A | 41.7 | (7) | 0.6 | -- | - | 10-3 | 11 | 4 | -- | - | - | 2.2 | - | |
| | WASHINGTON V | III | 1A | 40.6 | (11) | -0.4 | -- | - | 10-2 | 10 | 3 | -- | - | - | 3.0 | - | |
| NORTHRUP KING (NK) | S15-50 | I | 1C | -- | - | - | 37.9 | (9) | 0.0 | -- | - | - | 10-2 | 2 | -6 | - | 1.7 |
| | S18-84 | I | 1B,3 | 43.0 | (10) | 2.6 * | 44.1 | (18) | 3.4 * | 9-24 | 3 | -3 | 10-3 | 4 | -2 | 1.6 | 1.8 |
| | S23-03 | II | NONE | 41.6 | (11) | 0.6 | 40.9 | (8) | 1.2 | 9-27 | 5 | -2 | 10-6 | 7 | -1 | 2.2 | 2.0 |
| | S23-12 | II | NONE | 47.1 | (3) | 6.7 | 40.1 | (4) | 3.4 | 9-28 | 6 | 0 | 10-9 | 8 | 0 | 1.2 | 1.3 |
| | S2596 | II | 1A | 45.6 | (16) | 1.7 * | 44.9 | (8) | 0.8 | 9-28 | 7 | 1 | 10-9 | 10 | 2 | 1.8 | 1.7 |
| | S27-10 | II | 1C | 38.9 | (7) | -2.2 | 38.7 | (4) | -1.8 | 10-1 | 9 | 1 | 10-11 | 10 | 1 | 2.0 | 1.5 |
| PIONEER | 1981 | I | 1A | -- | - | - | 39.4 | (9) | 1.5 | -- | - | - | 10-5 | 5 | -2 | - | 1.7 |
| | 9251 | II | 1A | -- | - | - | 39.2 | (4) | 2.5 | -- | - | - | 10-12 | 10 | 3 | - | 1.3 |
| | 9271 | II | NONE | 43.2 | (11) | 2.3 * | 44.2 | (6) | 2.4 | 9-29 | 8 | 0 | 10-7 | 6 | 1 | 1.5 | 1.6 |
| | 9292 | II | NONE | 42.8 | (11) | 1.8 | 44.5 | (6) | 2.7 | 9-27 | 6 | -1 | 10-6 | 5 | -1 | 1.2 | 1.5 |
| PRIDE | B152 | I | 1C,NONE | -- | - | - | 39.2 | (9) | 1.4 | -- | - | - | 9-30 | 1 | -6 | - | 1.8 |
| | B203 | II | 1C | 42.9 | (5) | -2.1 | 40.6 | (11) | 0.1 H | 9-26 | 0 | -3 | 10-4 | 6 | -1 | 1.3 | 1.9 |
| | B242 | II | NONE | 43.4 | (7) | 2.8 * | -- | - | 10-1 | 11 | 4 | -- | - | - | 1.7 | - | |
| RUPP | RS2300 | II | 1A | 43.3 | (18) | 0.5 | 43.6 | (17) | 1.6 | 9-23 | 4 | -2 | 10-4 | 5 | -1 | 1.6 | 1.7 |
| | RS2460P | II | NONE | 44.7 | (11) | 3.8 * | 41.1 | (11) | 1.1 H | 10-2 | 10 | 3 | 10-9 | 10 | 4 | 2.1 | 2.0 |
| | RS2544 | III | NONE | 42.2 | (7) | 1.1 | -- | - | 10-6 | 14 | 7 | -- | - | - | 2.4 | - | |
| | EX 26905 | III | NONE | 43.3 | (3) | 3.0 | -- | - | 10-3 | 11 | 5 | -- | - | - | 1.7 | - | |

(CONT'D)

* STATISTICALLY SIGNIFICANT DEVIATION (P<.05)
H VARIETY EXHIBITS HIGHER THAN AVERAGE RESPONSE TO HIGHLY PRODUCTIVE ENVIRONMENTS.
L VARIETY EXHIBITS LOWER THAN AVERAGE RESPONSE TO HIGHLY PRODUCTIVE ENVIRONMENTS.

TABLE 3. (Continued) Performance Summary for Varieties Entered in the Michigan Trials in 1986. Phytophthora Resistance Designations Denote the Following: Type 1A Resistant to Races 1, 2, and 10; Type 1B Resistant to Races 1 and 3-9; Type 1C Resistant to Races 1-3 and 6-10; Type 1K Resistant to Races 1-10; Type 3 Resistant to Races 1-5, 8, and 9; Type 6 Resistant to Races 1-4 and 10.

| BRAND | ENTRY | PHYT. RES. TYPE | YIELD (BU/A) WITH DEVIATION FROM MEAN | | | | | | MATURITY RELATIVE TO CHECKS | | | | | | LODGING SCORE SOUTH. CENTRAL | |
|------------|----------------|-----------------------|---------------------------------------|-----------|------|-----------|------|-------|-----------------------------|-----|-------|---------|-----|--------|---------------------------------|-----|
| | | | SOUTHERN | | | CENTRAL | | | SOUTHERN | | | CENTRAL | | | | |
| | | | MG | YIELD (N) | DEV. | YIELD (N) | DEV. | DATE | H78 | C79 | DATE | H78 | C79 | SOUTH. | CENTRAL | |
| SCOTT FARM | L1808 | I 1A | -- | -- | | 32.9 | (5) | -2.5 | -- | -- | 10-6 | 4 | -3 | - | 2.0 | |
| STINE | L2456 | II 6 | 43.1 | (7) | 2.0 | -- | - | - | 10-1 | 9 | 1 | -- | -- | 2.7 | - | |
| | 2510 BRAND | II 1A | 42.1 | (11) | 1.2 | 43.2 | (3) | 0.6 | 9-29 | 8 | 1 | 10-11 | 10 | 0 | 2.2 | 1.9 |
| | 2530 BRAND | II 1A | 42.1 | (7) | 1.0 | -- | - | - | 10-4 | 12 | 5 | -- | - | 2.2 | - | |
| | 2710E BRAND | II 1A | -- | - | | 40.8 | (4) | 4.1 * | -- | - | 10-12 | 10 | 3 | - | 1.9 | |
| TERRA | 2820 BRAND | II NONE | -- | - | | 40.0 | (4) | 2.8 | -- | - | 10-7 | 6 | -2 | - | 2.1 | |
| | HURDLE BRAND | II 1A | -- | - | | 37.5 | (4) | 0.8 | -- | - | 10-10 | 9 | 2 | - | 1.5 | |
| | OLYMPIAN BRAND | II NONE | 39.7 | (3) | -0.7 | -- | - | - | 10-1 | 9 | 3 | -- | - | 1.7 | - | |
| | RUNNER BRAND | I 1A | -- | - | | 33.4 | (5) | -2.0 | -- | - | 10-9 | 7 | 0 | - | 1.8 | |
| VORIS | SPRINT BRAND | II NONE | 41.7 | (3) | 1.3 | -- | - | - | 10-3 | 11 | 5 | -- | - | 1.7 | - | |
| | V207 | II 1A | 44.3 | (15) | 0.8 | 44.8 | (19) | 2.5 * | 9-23 | 3 | -3 | 10-4 | 5 | -2 | 2.0 | 2.0 |
| | V235 | II 1A | 39.0 | (3) | -1.4 | 35.6 | (4) | -1.2 | 10-2 | 10 | 4 | 10-12 | 11 | 4 | 1.8 | 1.7 |
| | V311 | III NONE | 42.2 | (11) | 1.3 | 41.8 | (8) | 2.1 | 10-3 | 11 | 4 | 10-13 | 15 | 6 | 2.8 | 2.6 |

* STATISTICALLY SIGNIFICANT DEVIATION (P<.05)

H VARIETY EXHIBITS HIGHER THAN AVERAGE RESPONSE TO HIGHLY PRODUCTIVE ENVIRONMENTS.

L VARIETY EXHIBITS LOWER THAN AVERAGE RESPONSE TO HIGHLY PRODUCTIVE ENVIRONMENTS.

TABLE 4. Southern Michigan.

| BRAND | ENTRY | YIELD (BU/A) | | | | | | M A T H O U E D S R I G C I G I N R T H N R T Y T G E | | | | | | | | |
|----------|------------------|------------------------|------|------|-------------------------|------|------|---|------|-----|----------------------------|------|-----|------|----|-----|
| | | ENTIRE SOUTHERN REGION | | | SOUTHEAST (LENAWEE CO.) | | | SOUTHWEST (ST. JOSEPH CO.) | | | SOUTH CENTRAL (INGHAM CO.) | | | | | |
| | | 1986 | AVG. | (N) | 1986 | AVG. | (N) | 1986 | AVG. | (N) | 1986 | AVG. | (N) | | | |
| PUBLIC | BSR 101 (I) | 43.2* | 46.4 | (11) | 49.2 | 51.0 | (6) | 40.9* | 42.6 | (2) | 39.6* | 44.0 | (2) | -1 | 30 | 1.5 |
| PUBLIC | HARDIN | 41.5* | 42.7 | (17) | 48.1 | 51.5 | (7) | 36.1 | 35.0 | (4) | 40.2* | 42.7 | (3) | -2 | 31 | 2.3 |
| PUBLIC | HODGSON 78 | 36.1 | 40.3 | (26) | 46.3 | 48.2 | (10) | 31.6 | 31.7 | (7) | 30.5 | 41.9 | (4) | -6 | 30 | 1.6 |
| PUBLIC | SIBLEY | 43.3* | 49.3 | (8) | 49.1 | 52.3 | (6) | 33.4 | -- | - | 47.3* | -- | - | -5 | 29 | 2.1 |
| PUBLIC | WEBER 84 | 37.5 | 40.2 | (13) | 37.6 | 47.1 | (8) | 31.6 | 33.8 | (3) | 43.3* | 44.1 | (3) | -4 | 32 | 3.0 |
| PUBLIC | AMCOR (II) | 39.2 | 41.8 | (17) | 46.2 | 50.8 | (7) | 33.1 | 30.8 | (4) | 38.4 | 40.9 | (3) | 4 | 38 | 3.6 |
| PUBLIC | BEESON 80 | 26.3 | 38.1 | (17) | 30.3 | 45.0 | (7) | 28.1 | 27.2 | (4) | 20.4 | 36.6 | (3) | 5 | 27 | 1.4 |
| PUBLIC | BSR 201 | 41.5* | 47.4 | (14) | 47.6 | 51.2 | (9) | 38.8* | 39.7 | (2) | 38.2 | 46.0 | (2) | 3 | 31 | 2.2 |
| PUBLIC | CENTURY | 40.2 | 44.2 | (18) | 53.4* | 52.2 | (8) | 35.7 | 34.8 | (4) | 31.5 | 41.4 | (3) | 4 | 34 | 1.9 |
| PUBLIC | CENTURY 84 | 40.3 | 42.1 | (9) | 50.3 | 48.6 | (4) | 33.8 | 35.7 | (2) | 36.7 | 40.5 | (2) | 3 | 32 | 1.4 |
| PUBLIC | CORSOY | 39.8 | 41.4 | (26) | 44.7 | 47.4 | (11) | 35.0 | 33.1 | (6) | 39.7* | 41.7 | (5) | -1 | 33 | 3.1 |
| + PUBLIC | + CORSOY 79 | 38.5 | 42.7 | (24) | 46.7 | 52.1 | (10) | 31.4 | 30.3 | (6) | 37.3 | 43.5 | (4) | 9-28 | 34 | 3.1 |
| PUBLIC | ELGIN | 41.4* | 44.5 | (15) | 51.3 | 51.0 | (7) | 35.5 | 39.2 | (3) | 37.3 | 42.3 | (3) | 2 | 27 | 1.7 |
| PUBLIC | HACK | 42.7* | 45.3 | (10) | 51.0 | 50.8 | (5) | 41.2* | 43.0 | (2) | 36.0 | 41.5 | (2) | 2 | 30 | 1.1 |
| PUBLIC | HOYT | 40.6 | 48.1 | (7) | 52.5 | 53.5 | (5) | 39.6* | -- | - | 29.6 | -- | - | 4 | 22 | 1.2 |
| PUBLIC | KELLER | 38.5 | 39.5 | (8) | 49.3 | 45.4 | (3) | 32.5 | 32.4 | (2) | 33.7 | 41.4 | (2) | 1 | 31 | 1.8 |
| PUBLIC | MIAMI | 36.7 | 38.7 | (9) | 46.7 | 45.2 | (4) | 32.1 | 29.5 | (2) | 31.4 | 39.0 | (2) | -1 | 31 | 1.7 |
| PUBLIC | NEBSOY | 38.7 | 41.8 | (18) | 46.1 | 47.3 | (8) | 30.9 | 31.8 | (4) | 39.2* | 47.0 | (3) | -0 | 30 | 1.4 |
| PUBLIC | PRESTON | 42.0* | 47.0 | (6) | 50.6 | 51.6 | (4) | 35.9 | -- | - | 39.6* | -- | - | 5 | 34 | 2.4 |
| PUBLIC | VICKERY | 39.5 | 42.6 | (18) | 48.8 | 51.5 | (8) | 34.2 | 31.3 | (4) | 35.4 | 40.7 | (3) | 0 | 35 | 3.3 |
| PUBLIC | WELLS II | 37.4 | 40.7 | (20) | 44.8 | 46.7 | (8) | 30.8 | 32.5 | (5) | 36.7 | 40.8 | (3) | -1 | 33 | 2.0 |
| PUBLIC | CUMBERLAND (III) | 33.6 | 36.8 | (13) | 47.9 | 48.9 | (4) | 29.8 | 27.8 | (4) | 23.2 | 34.7 | (3) | 6 | 33 | 2.4 |
| PUBLIC | GNOME 85 | 37.4 | 47.8 | (10) | 45.7 | 49.3 | (9) | 33.9 | -- | - | -- | -- | - | 7 | 21 | 0.9 |
| PUBLIC | HOBBIT | 38.6 | 40.0 | (15) | 46.9 | 49.5 | (5) | 33.3 | 30.4 | (5) | 35.6 | 44.8 | (3) | 6 | 23 | 1.0 |
| PUBLIC | PELLA | 38.7 | 40.2 | (13) | 52.2 | 50.0 | (4) | 30.7 | 30.4 | (4) | 33.2 | 45.0 | (3) | 4 | 34 | 1.6 |
| PUBLIC | SPRITE | 40.3 | 39.9 | (16) | 51.5 | 51.1 | (6) | 30.9 | 28.9 | (5) | 38.5 | 41.9 | (3) | 7 | 26 | 1.8 |
| PUBLIC | WILLIAMS 82 | 35.1 | 39.0 | (10) | 51.8 | 50.7 | (3) | 31.6 | 34.7 | (3) | 21.9 | 34.3 | (2) | 8 | 36 | 2.1 |

(CONT'D)

LSD (.05) 6.22 6.79 4.22 8.90 2.3 2.6 0.80

* CHECK VARIETY USED TO CALCULATE DEVIATION FROM STANDARD MATURITY.

* NOT SIGNIFICANTLY DIFFERENT FROM HIGHEST YIELD WITHIN THAT COLUMN.

TABLE 4. (Continued) Southern Michigan.

| BRAND | ENTRY | YIELD (BU/A) | | | | | | | | | | M | L |
|---------------|--------------|------------------------|------|------|--------------------------|------|-----|----------------------------|------|-----|----------------------------|------|-----|
| | | ENTIRE SOUTHERN REGION | | | SOUTHEAST (LENAWEET CO.) | | | SOUTHWEST (ST. JOSEPH CO.) | | | SOUTH CENTRAL (INGHAM CO.) | | |
| | | 1986 | AVG. | (N) | 1986 | AVG. | (N) | 1986 | AVG. | (N) | 1986 | AVG. | (N) |
| PUBLIC | WINCHESTER | 34.9 | 38.9 | (7) | 46.2 | 47.8 | (2) | 31.4 | 31.9 | (2) | 27.0 | 39.4 | (2) |
| PUBLIC | ZANE | 39.9 | 41.5 | (7) | 50.8 | 49.7 | (2) | 37.2* | 39.3 | (2) | 31.7 | 37.1 | (2) |
| AGRIPRO | AP247 | 40.0 | 41.0 | (15) | 48.3 | 48.7 | (5) | 35.3 | 35.7 | (4) | 36.3 | 42.4 | (3) |
| AGRIPRO | AP2190 | 39.8 | 40.9 | (7) | 52.4 | 52.2 | (2) | 33.8 | 35.8 | (2) | 33.3 | 42.7 | (2) |
| AGRIPRO | AP3023 | 41.1* | -- | -- | 57.6* | -- | - | 37.9* | -- | - | 27.7 | -- | - |
| AGRIPRO | AP3132 | 40.8 | -- | -- | 52.2 | -- | - | 35.9 | -- | - | 34.2 | -- | - |
| AGRIPRO | HP20-20 | 40.0 | 43.4 | (11) | 48.5 | 50.8 | (5) | 36.0 | 35.9 | (2) | 35.6 | 38.4 | (2) |
| AGRIPRO | HP2530 | 41.6* | 44.0 | (14) | 49.8 | 50.7 | (5) | 34.5 | 36.1 | (3) | 40.6* | 47.7 | (3) |
| ASGROW | A1937 | 39.4 | 42.6 | (16) | 48.3 | 51.0 | (6) | 35.8 | 35.1 | (4) | 34.1 | 43.4 | (3) |
| ASGROW | A2187 | 40.8 | 40.9 | (7) | 49.3 | 46.9 | (2) | 38.1* | 38.2 | (2) | 35.1 | 41.0 | (2) |
| ASGROW | A2522 | 41.8* | 41.7 | (11) | 54.3* | 52.7 | (3) | 33.4 | 35.5 | (3) | 37.8 | 45.2 | (3) |
| ASGROW | A2943 | 41.2* | 43.8 | (11) | 53.2* | 52.0 | (3) | 36.6 | 40.4 | (3) | 33.9 | 42.5 | (3) |
| CALLAHAN | 1250 | 41.1* | 41.0 | (13) | 54.8* | 49.4 | (4) | 33.0 | 33.5 | (4) | 35.6 | 44.9 | (3) |
| CALLAHAN | 5300 | 41.6* | 43.2 | (7) | 59.6* | 56.6 | (2) | 36.2 | 38.4 | (2) | 29.0 | 38.8 | (2) |
| CALLAHAN | 5350 | 38.8 | 42.6 | (11) | 55.4* | 54.9 | (3) | 29.9 | 36.2 | (3) | 31.0 | 42.1 | (3) |
| CALLAHAN | 6262X | 38.7 | 41.4 | (7) | 49.5 | 51.1 | (2) | 31.0 | 34.0 | (2) | 35.6 | 43.8 | (2) |
| CALLAHAN | 7244X | 43.9* | -- | -- | 54.0* | -- | - | 35.9 | -- | - | 41.8* | -- | - |
| CALLAHAN | 7260X | 43.5* | -- | -- | 55.0* | -- | - | 35.3 | -- | - | 40.1* | -- | - |
| CALLAHAN | 7272X | 43.3* | -- | -- | 56.1* | -- | - | 36.0 | -- | - | 37.8 | -- | - |
| COUNTRYMARK | FFR 112 | 40.2 | -- | -- | 48.9 | -- | - | 36.3 | -- | - | 35.4 | -- | - |
| COUNTRYMARK | FFR 226 | 38.4 | 35.4 | (5) | 47.8 | 44.6 | (2) | 33.1 | 26.8 | (2) | 34.2 | -- | - |
| COUNTRYMARK | FFR 241 | 40.0 | -- | -- | 50.8 | -- | - | 34.4 | -- | - | 34.8 | -- | - |
| DAIRYLAND | DSR-171 | 38.1 | 42.8 | (18) | 42.7 | 49.8 | (8) | 37.1* | 36.7 | (4) | 34.4 | 40.6 | (3) |
| DAIRYLAND | DSR-212 | 38.0 | 39.2 | (15) | 52.6 | 46.6 | (5) | 38.4* | 35.1 | (4) | 23.1 | 38.0 | (3) |
| DAIRYLAND | DSR-255 | 39.3 | 37.9 | (7) | 50.0 | 44.0 | (2) | 34.5 | 34.4 | (2) | 33.3 | 38.8 | (2) |
| DAIRYLAND | DSR-287 | 39.7 | 42.3 | (7) | 50.3 | 48.8 | (2) | 32.9 | 39.4 | (2) | 36.0 | 42.9 | (2) |
| DAIRYLAND | DSR-297 | 39.9 | 40.4 | (7) | 52.3 | 49.9 | (2) | 32.3 | 34.0 | (2) | 35.1 | 40.8 | (2) |
| DAIRYLAND | DSR-317 | 36.9 | 40.5 | (7) | 51.3 | 52.4 | (2) | 36.9 | 39.9 | (2) | 22.6 | 30.2 | (2) |
| DAIRYLAND | DST-2203 | 37.6 | -- | -- | 43.1 | -- | - | 34.6 | -- | - | 35.0 | -- | - |
| DAIRYLAND | DST-2204 | 33.3 | -- | -- | 44.8 | -- | - | 32.4 | -- | - | 22.6 | -- | - |
| DEKALB-PFIZER | CX283 | 43.0* | 42.3 | (11) | 52.6 | 51.3 | (3) | 35.8 | 35.4 | (3) | 40.5* | 45.4 | (3) |
| DEKALB-PFIZER | CX326 | 38.4 | -- | -- | 51.7 | -- | - | 34.7 | -- | - | 28.7 | -- | - |
| DIEHL FIELDS | EXP 101 | 45.6* | -- | -- | 57.2* | -- | - | 37.2* | -- | - | 42.4* | -- | - |
| FUNK | G3232 | 42.7* | -- | -- | 53.8* | -- | - | 36.8 | -- | - | 37.5 | -- | - |
| FUNK | G3239 | 41.9* | 42.3 | (8) | 54.6* | 49.4 | (2) | 33.8 | 38.1 | (2) | 37.4 | 43.3 | (3) |
| GLH | GL2634 | 44.6* | 45.5 | (16) | 57.9* | 53.8 | (6) | 34.5 | 34.9 | (4) | 41.5* | 46.0 | (3) |
| GLH | GL2908 | 41.3* | 43.0 | (11) | 54.8* | 55.5 | (3) | 35.2 | 37.5 | (3) | 33.8 | 41.3 | (3) |
| GLH | XP2478 | 42.9* | -- | -- | 50.5 | -- | - | 35.8 | -- | - | 42.5* | -- | - |
| GLH | XP2737 | 42.4* | -- | -- | 52.5 | -- | - | 35.5 | -- | - | 39.2* | -- | - |
| JACQUES | J-201 | 41.6* | -- | -- | 51.1 | -- | - | 36.3 | -- | - | 37.5 | -- | - |
| JACQUES | J-231 | 43.9* | 43.8 | (11) | 54.5* | 54.0 | (3) | 36.7 | 37.5 | (3) | 40.5* | 46.9 | (3) |
| LAKESIDE | 21 | 42.5* | -- | -- | 54.2* | -- | - | 38.8* | -- | - | 34.4 | -- | - |
| LAKESIDE | 30 | 37.5 | -- | -- | 49.4 | -- | - | 27.2 | -- | - | 35.9 | -- | - |
| LAKESIDE | 36 | 38.5 | -- | -- | 51.7 | -- | - | 32.8 | -- | - | 31.1 | -- | - |
| LAKESIDE | 125 | 41.5* | -- | -- | 50.4 | -- | - | 30.5 | -- | - | 43.5* | -- | - |
| LAKESIDE | MILLER | 39.0 | 43.3 | (7) | 49.5 | 51.2 | (2) | 39.8* | 44.5 | (2) | 27.6 | 37.2 | (2) |
| MAUMEE VALLEY | CALIBER | 41.3* | 41.5 | (11) | 47.9 | 48.4 | (3) | 36.8 | 38.3 | (3) | 39.2* | 44.8 | (3) |
| MAUMEE VALLEY | ENTERPRISE | 39.7 | 39.2 | (7) | 53.4* | 48.9 | (2) | 35.3 | 35.5 | (2) | 30.5 | 34.8 | (2) |
| MAUMEE VALLEY | KODIAK | 38.2 | 41.2 | (7) | 49.0 | 49.4 | (2) | 30.3 | 34.8 | (2) | 35.3 | 42.2 | (2) |
| MAUMEE VALLEY | MV-2E1 | 39.8 | 41.9 | (11) | 49.7 | 50.0 | (3) | 34.6 | 37.6 | (3) | 35.0 | 42.7 | (3) |
| MAUMEE VALLEY | WARRIOR | 40.0 | 41.7 | (7) | 51.6 | 51.0 | (2) | 32.1 | 34.7 | (2) | 36.4 | 42.0 | (2) |
| MAUMEE VALLEY | WASHINGTON V | 33.2 | 40.6 | (11) | 45.5 | 51.8 | (3) | 31.5 | 37.7 | (3) | 22.6 | 36.8 | (3) |
| NK | S23-03 | 41.9* | 41.6 | (11) | 51.4 | 49.4 | (3) | 39.9* | 40.8 | (3) | 34.5 | 40.3 | (3) |
| NK | S23-12 | 47.1* | -- | -- | 54.0* | -- | - | 39.8* | -- | - | 47.5* | -- | - |
| NK | S2596 | 41.3* | 45.6 | (16) | 51.8 | 53.3 | (7) | 37.4* | 38.8 | (3) | 34.7 | 41.7 | (3) |
| NK | S27-10 | 37.5 | 38.9 | (7) | 46.0 | 46.2 | (2) | 33.6 | 36.2 | (2) | 32.9 | 40.2 | (2) |
| PIONEER | 9271 | 44.5* | 43.2 | (11) | 53.4* | 52.9 | (3) | 38.7* | 39.7 | (3) | 41.4* | 46.5 | (3) |
| PIONEER | 9292 | 41.0* | 42.8 | (11) | 50.3 | 51.7 | (3) | 32.1 | 38.3 | (3) | 40.6* | 46.7 | (3) |
| PRIDE | B242 | 42.4* | 43.4 | (7) | 52.5 | 53.9 | (2) | 35.3 | 36.9 | (2) | 39.4* | 44.4 | (2) |
| RUPP | RS2300 | 38.9 | 43.3 | (18) | 47.6 | 51.6 | (8) | 33.5 | 33.1 | (4) | 35.6 | 44.1 | (3) |
| RUPP | RS2460P | 42.8* | 44.7 | (11) | 54.2* | 55.7 | (3) | 38.0* | 41.1 | (3) | 36.2 | 44.7 | (3) |
| RUPP | RS2544 | 39.3 | 42.2 | (7) | 51.0 | 47.8 | (2) | 37.9* | 43.1 | (2) | 29.1 | 36.2 | (2) |
| RUPP | EX 26905 | 43.3* | -- | -- | 53.5* | -- | - | 36.2 | -- | - | 40.3* | -- | - |

(CONT'D)

LSD(.05) 6.22 6.79 4.22 8.90 2.3 2.6 0.80

* NOT SIGNIFICANTLY DIFFERENT FROM HIGHEST YIELD WITHIN THAT COLUMN.

TABLE 4. (Continued) Southern Michigan.

| BRAND | ENTRY | YIELD (BU/A) | | | | | | | | | | | | M A T U E R I G I T H N R O D S C I O G E | | |
|-------|-----------|------------------------|------|------|--------------------------|------|-----|----------------------------|------|-----|----------------------------|------|-----|---|------|------|
| | | ENTIRE SOUTHERN REGION | | | SOUTHEAST (LENAWEET CO.) | | | SOUTHWEST (ST. JOSEPH CO.) | | | SOUTH CENTRAL (INGHAM CO.) | | | | | |
| | | 1986 | Avg. | (N) | 1986 | Avg. | (N) | 1986 | Avg. | (N) | 1986 | Avg. | (N) | | | |
| SCOTT | L2456 | 42.7* | 43.1 | (7) | 50.7 | 49.2 | (2) | 35.0 | 36.1 | (2) | 42.5* | 48.8 | (2) | 3 | 32 | 2.5 |
| STINE | 2510 | 39.6 | 42.1 | (11) | 51.1 | 53.2 | (3) | 31.6 | 37.3 | (3) | 36.1 | 43.2 | (3) | 2 | 26 | 1.7 |
| STINE | 2530 | 39.3 | 42.1 | (7) | 51.9 | 52.6 | (2) | 32.3 | 35.5 | (2) | 33.6 | 41.1 | (2) | 5 | 34 | 1.8 |
| TERRA | OLYMPIAN | 39.7 | -- | -- | 53.2* | -- | - | 35.8 | -- | - | 30.1 | -- | - | 3 | 30 | 1.7 |
| TERRA | SPRINT | 41.7* | -- | -- | 52.2 | -- | - | 37.7* | -- | - | 35.2 | -- | - | 5 | 33 | 1.7 |
| VORIS | V207 | 41.5* | 44.3 | (15) | 45.8 | 49.5 | (7) | 34.2 | 32.9 | (3) | 44.5* | 48.4 | (3) | -2 | 36 | 2.1 |
| VORIS | V235 | 39.0 | -- | -- | 46.6 | -- | - | 33.2 | -- | - | 37.1 | -- | - | 4 | 32 | 1.8 |
| VORIS | V311 | 39.0 | 42.2 | (11) | 54.6* | 51.3 | (3) | 29.4 | 36.0 | (3) | 33.0 | 43.4 | (3) | 5 | 34 | 2.5 |
| | LSD(.05) | 6.22 | | | 6.79 | | | 4.22 | | | 8.90 | | | 2.3 | 2.6 | 0.80 |
| | TEST MEAN | 40.0 | | | 50.3 | | | 34.6 | | | 35.2 | | | 2.8 | 32.1 | 1.94 |
| | CV | 9.7 | | | 7.9 | | | 7.1 | | | 14.5 | | | 5.1 | 25.7 | |

+ CHECK VARIETY USED TO CALCULATE DEVIATION FROM STANDARD MATURITY.

* NOT SIGNIFICANTLY DIFFERENT FROM HIGHEST YIELD WITHIN THAT COLUMN.

TABLE 5. Central Michigan.

| BRAND | ENTRY | YIELD (BU/A) | | | | | | | | | | | | M A T U E R I G I T H N R O D S C I O G E | | | | | |
|--------|-------------|-----------------------|------|------|----------------------------|------|-----|-----------------------|------|------|----------------------------|------|------|---|------|------|----|----|-----|
| | | ENTIRE CENTRAL REGION | | | SOUTH CENTRAL (INGHAM CO.) | | | CENTRAL (SAGINAW CO.) | | | EAST CENTRAL (SANILAC CO.) | | | | | | | | |
| | | 1986 | Avg. | (N) | 1986 | Avg. | (N) | 1986 | Avg. | (N) | 1986 | Avg. | (N) | | | | | | |
| PUBLIC | DASSEL (O) | 28.2 | 36.3 | (7) | -- | -- | - | 31.5 | 43.4 | (2) | 26.7 | 26.4 | (2) | 26.2 | 26.4 | (2) | -6 | 25 | 1.2 |
| PUBLIC | DAWSON | 36.3 | 40.5 | (19) | -- | -- | - | 30.6 | 39.3 | (7) | 38.5* | 41.5 | (5) | 39.7 | 41.5 | (5) | -8 | 28 | 1.8 |
| PUBLIC | EVANS | 31.6 | 36.8 | (31) | -- | -- | - | 29.5 | 37.5 | (14) | 32.0 | 36.2 | (9) | 33.2 | 36.2 | (9) | -7 | 27 | 1.6 |
| PUBLIC | BSR 101 (I) | 39.1* | 41.4 | (13) | 39.6* | 44.0 | (2) | 31.4 | 43.0 | (6) | 41.3* | 40.1 | (3) | 44.2* | 40.1 | (3) | 1 | 32 | 1.8 |
| PUBLIC | HARDIN | 38.2* | 45.2 | (22) | 40.2* | 42.7 | (3) | 39.1 | 48.7 | (9) | 37.8 | 44.8 | (7) | 35.6 | 44.8 | (7) | -3 | 32 | 2.2 |
| PUBLIC | HODGSON 78 | 29.4 | 41.3 | (33) | 30.5 | 41.9 | (4) | 28.0 | 41.3 | (15) | 28.9 | 41.9 | (10) | 30.1 | 41.9 | (10) | -8 | 29 | 2.0 |
| PUBLIC | SIBLEY | 36.5 | 40.8 | (10) | 47.3* | -- | - | 28.3 | 43.0 | (6) | 36.3 | 35.2 | (2) | 34.2 | 35.2 | (2) | -8 | 30 | 2.1 |
| PUBLIC | WEBER 84 | 37.3 | 38.8 | (15) | 43.3* | 44.1 | (3) | 27.3 | 39.9 | (6) | 39.5* | 36.3 | (4) | 39.0 | 36.3 | (4) | -5 | 33 | 2.4 |
| PUBLIC | AMCOR (II) | 38.4* | 42.3 | (17) | 38.4 | 40.9 | (3) | 32.3 | 42.5 | (8) | 40.0* | 42.9 | (6) | 43.0* | 42.9 | (6) | 4 | 36 | 2.7 |
| PUBLIC | BEESON 80 | 29.1 | 39.2 | (19) | 20.4 | 36.6 | (3) | 32.3 | 43.3 | (9) | 31.1 | 35.2 | (7) | 32.4 | 35.2 | (7) | 6 | 28 | 1.7 |
| PUBLIC | BSR 201 | 35.4 | 42.2 | (12) | 38.2 | 46.0 | (2) | 22.2 | 42.1 | (7) | 39.6* | 39.9 | (3) | 41.5 | 39.9 | (3) | 3 | 32 | 2.5 |
| PUBLIC | CENTURY | 36.1 | 42.3 | (20) | 31.5 | 41.4 | (3) | 29.9 | 43.3 | (10) | 39.2* | 41.3 | (7) | 43.9* | 41.3 | (7) | 4 | 32 | 1.7 |
| PUBLIC | CENTURY 84 | 39.0* | 41.4 | (9) | 36.7 | 40.5 | (2) | 36.6 | 42.2 | (4) | 42.3* | 41.1 | (3) | 40.3 | 41.1 | (3) | 4 | 32 | 1.5 |
| PUBLIC | + CORSOY 79 | 36.2 | 39.4 | (29) | 39.7* | 41.7 | (5) | 26.0 | 40.2 | (14) | 39.3* | 36.6 | (8) | 39.9 | 36.6 | (8) | -1 | 34 | 2.5 |
| PUBLIC | ELGIN | 37.3 | 43.8 | (16) | 37.3 | 42.3 | (3) | 30.9 | 46.6 | (8) | 41.0* | 42.2 | (4) | 40.0 | 42.2 | (4) | 1 | 26 | 1.8 |
| PUBLIC | HACK | 39.2* | 41.7 | (10) | 36.0 | 41.5 | (2) | 37.4 | 42.1 | (5) | 44.5* | 41.2 | (3) | 38.9 | 41.2 | (3) | 2 | 30 | 1.5 |
| PUBLIC | HOYT | 34.0 | 39.6 | (8) | 29.6 | -- | - | 28.4 | 41.8 | (5) | 36.3 | 39.0 | (2) | 41.7 | 39.0 | (2) | 2 | 27 | 1.9 |
| PUBLIC | KELLER | 34.6 | 40.0 | (8) | 33.7 | 41.4 | (2) | 29.2 | 40.0 | (3) | 36.9 | 39.1 | (3) | 38.7 | 39.1 | (3) | 3 | 30 | 1.8 |
| PUBLIC | MIAMI | 33.3 | 38.5 | (9) | 31.4 | 39.0 | (2) | 25.8 | 39.3 | (4) | 35.1 | 37.1 | (3) | 40.8 | 37.1 | (3) | 1 | 31 | 1.6 |
| PUBLIC | NEBSOY | 36.6 | 40.6 | (20) | 39.2* | 47.0 | (3) | 34.6 | 42.2 | (10) | 38.7* | 35.6 | (7) | 33.9 | 35.6 | (7) | 1 | 29 | 1.8 |
| PUBLIC | PRESTON | 36.4 | 38.5 | (7) | 39.6* | -- | - | 38.4 | 40.6 | (4) | 38.4* | 33.7 | (2) | 29.0 | 33.7 | (2) | 4 | 33 | 2.1 |
| PUBLIC | VICKERY | 36.2 | 42.1 | (21) | 35.4 | 40.7 | (3) | 31.3 | 42.6 | (11) | 41.3* | 41.8 | (7) | 36.7 | 41.8 | (7) | 0 | 34 | 2.9 |
| PUBLIC | WELLS II | 33.9 | 40.2 | (22) | 36.7 | 40.8 | (3) | 31.9 | 42.7 | (11) | 34.1 | 36.4 | (8) | 32.9 | 36.4 | (8) | -1 | 32 | 1.6 |
| PUBLIC | PELLA (III) | 37.4 | 43.4 | (13) | 33.2 | 45.0 | (3) | 29.7 | 39.3 | (5) | 40.6* | 46.7 | (5) | 45.9* | 46.7 | (5) | 6 | 33 | 1.8 |

(CONT'D)

LSD(.05) 5.71 8.90 6.49 6.96 10.65 2.4 2.8 0.55

+ CHECK VARIETY USED TO CALCULATE DEVIATION FROM STANDARD MATURITY.

* NOT SIGNIFICANTLY DIFFERENT FROM HIGHEST YIELD IN THAT COLUMN.

TABLE 5. (Continued) Central Michigan.

| BRAND | ENTRY | YIELD (BU/A) | | | | | | | | | | M | A | L | | | | | |
|-----------|--------|-----------------------|------|------|----------------------------|------|-----|-----------------------|------|-----|----------------------------|------|------------------------------|-------|------|-----|------|------|------|
| | | ENTIRE CENTRAL REGION | | | SOUTH CENTRAL (INGHAM CO.) | | | CENTRAL (SAGINAW CO.) | | | EAST CENTRAL (SANILAC CO.) | | EAST CENTRAL (ST. CLAIR CO.) | | T | H | D | | |
| | | 1986 | AVG. | (N) | 1986 | AVG. | (N) | 1986 | AVG. | (N) | 1986 | AVG. | (N) | Y | T | G | | | |
| SCOTT | L1808 | 34.0 | 32.9 | (5) | 31.5 | -- | - | 30.9 | -- | - | 39.6* | 36.8 | (2) | 33.9 | 36.8 | (2) | -2 | 29 | 2.1 |
| STINE | 2710E | 40.8* | -- | -- | 40.7* | -- | - | 35.4 | -- | - | 44.5* | 43.6 | (2) | 42.7* | 43.6 | (2) | 3 | 34 | 1.9 |
| STINE | 2820 | 39.6* | -- | -- | 43.0* | -- | - | 35.2 | -- | - | 37.6 | 40.0 | (2) | 42.4 | 40.0 | (2) | -2 | 34 | 2.1 |
| TERRA | HURDLE | 37.5* | -- | -- | 37.6 | -- | - | 36.0 | -- | - | 39.6* | 38.2 | (2) | 36.8 | 38.2 | (2) | 2 | 31 | 1.5 |
| TERRA | RUNNER | 34.7 | 33.4 | (5) | 38.3 | -- | - | 23.4 | -- | - | 36.7 | 38.4 | (2) | 40.2 | 38.4 | (2) | 0 | 32 | 2.0 |
| VORIS | V207 | 39.5* | 44.8 | (19) | 44.5* | 48.4 | (3) | 31.6 | 45.1 | (9) | 36.4 | 42.8 | (7) | 45.4* | 42.8 | (7) | -1 | 36 | 1.7 |
| VORIS | V235 | 35.6 | -- | -- | 37.1 | -- | - | 31.3 | -- | - | 40.7* | 37.0 | (2) | 33.2 | 37.0 | (2) | 4 | 30 | 1.7 |
| VORIS | V311 | 37.5* | 41.8 | (8) | 33.0 | 43.4 | (3) | 27.5 | 37.6 | (2) | 43.3* | 43.0 | (3) | 46.2* | 43.0 | (3) | 7 | 34 | 2.5 |
| LSD(.05) | | 5.71 | | | 8.90 | | | 6.49 | | | 6.96 | | | 10.65 | | | 2.4 | 2.8 | 0.55 |
| TEST MEAN | | 36.9 | | | 36.11 | | | 32.01 | | | 38.50 | | | 40.26 | | | 31.0 | 1.88 | |
| CV | | 11.2 | | | 14.5 | | | 11.8 | | | 10.5 | | | 15.4 | | | 6.0 | 21.1 | |

* NOT SIGNIFICANTLY DIFFERENT FROM HIGHEST YIELD IN THAT COLUMN.

TABLE 6. Saginaw Bay Area (Huron Co.).

| BRAND | ENTRY | YIELD | | | MATURITY | | | LODGING SCORE | | |
|---------------|--------------|-------|------|-----|----------|------|--------|---------------|--|-----|
| | | 1985 | AVG. | (N) | DATE | DEV | HEIGHT | | | |
| PUBLIC | DASSEL | 23.9 | 38.3 | (3) | 10-2 | -3 | 24 | | | 1.6 |
| PUBLIC | DAWSON | 32.4 | 40.2 | (5) | 9-25 | -10 | 31 | | | 2.2 |
| PUBLIC | EVANS | 29.0 | 35.0 | (4) | 9-27 | -8 | 30 | | | 1.8 |
| PUBLIC | OZZIE | 28.0 | 34.9 | (4) | 9-26 | -8 | 29 | | | 1.3 |
| PUBLIC | SIMPSON | 34.4 | 39.2 | (4) | 9-28 | -7 | 28 | | | 1.6 |
| PUBLIC | BSR 101 | 29.4 | 36.2 | (2) | 10-7 | 3 | 30 | | | 1.8 |
| PUBLIC | HARDIN | 28.0 | 37.8 | (3) | 10-5 | 1 | 29 | | | 2.0 |
| PUBLIC | + HODGSON 78 | 24.8 | 39.5 | (4) | 10-5 | 10-5 | 27 | | | 1.9 |
| PUBLIC | SIBLEY | 31.9 | -- | - | 10-5 | 0 | 34 | | | 2.5 |
| PUBLIC | WEBER 84 | 28.2 | 32.4 | (2) | 10-6 | 1 | 38 | | | 2.2 |
| PUBLIC | CORSOY 79 | 30.6 | 37.2 | (3) | 10-9 | 4 | 38 | | | 2.3 |
| PUBLIC | ELGIN | 33.1 | -- | - | 10-7 | 3 | 28 | | | 2.3 |
| ASGROW | A0949 | 28.6 | -- | - | 10-2 | -3 | 36 | | | 1.2 |
| ASGROW | A1525 | 30.8 | 35.0 | (2) | 10-5 | 1 | 31 | | | 1.6 |
| ASGROW | A1937 | 31.8 | 43.5 | (3) | 10-3 | -2 | 29 | | | 2.1 |
| DAIRYLAND | DSR-120 | 33.6 | 39.3 | (3) | 9-29 | -6 | 30 | | | 1.9 |
| DAIRYLAND | DSR-128 | 32.5 | 35.0 | (3) | 10-1 | -4 | 29 | | | 0.8 |
| DAIRYLAND | DSR-135 | 29.7 | 35.0 | (3) | 9-30 | -5 | 33 | | | 2.0 |
| DAIRYLAND | DSR-171 | 28.4 | 38.3 | (3) | 10-6 | 2 | 35 | | | 2.3 |
| DAIRYLAND | DST-1103 | 28.3 | -- | - | 9-30 | -5 | 29 | | | 2.1 |
| DAIRYLAND | DST-1207 | 30.5 | -- | - | 10-5 | 0 | 32 | | | 2.3 |
| DEKALB-PFIZER | CX174 | 31.4 | 36.7 | (2) | 10-7 | 3 | 31 | | | 1.5 |
| FUNK | G3145 | 33.0 | 38.0 | (2) | 10-7 | 2 | 35 | | | 2.3 |
| FUNK | G3180 | 32.3 | -- | - | 10-4 | 0 | 30 | | | 2.0 |
| FUNK | 12283 | 32.5 | -- | - | 10-6 | 1 | 28 | | | 1.7 |
| GLH | GL 1434 | 28.0 | 30.6 | (2) | 10-5 | 1 | 35 | | | 2.3 |
| GLH | GL 1900 | 31.4 | 34.5 | (2) | 10-8 | 3 | 27 | | | 2.0 |
| GLH | GL 1937 | 32.0 | 38.5 | (3) | 10-6 | 1 | 29 | | | 2.0 |
| JACQUES | E85092 | 29.3 | -- | - | 10-1 | -3 | 33 | | | 1.4 |
| KING GRAIN | KG60 | 30.4 | 31.6 | (2) | 9-28 | -6 | 24 | | | 1.3 |
| (CONT'D) | | | | | | | | | | |
| LSD(.05) | | NS | | | 3.4 | 4.9 | 0.99 | | | |

+ CHECK VARIETY USED TO CALCULATE DEVIATION FROM STANDARD MATURITY.

TABLE 6. (Continued) Saginaw Bay Area (Huron Co.).

| BRAND | ENTRY | YIELD | | | MATURITY | | | LODGING SCORE |
|------------|--------|-------|------|-----|----------|------|--------|------------------|
| | | 1985 | Avg. | (N) | DATE | DEV | HEIGHT | |
| KING GRAIN | KG70 | 31.1 | 37.3 | (3) | 10-4 | -1 | 36 | 2.3 |
| LAKESIDE | 15 | 32.2 | -- | - | 10-4 | -1 | 37 | 2.7 |
| LAKESIDE | 116 | 33.8 | -- | - | 10-8 | 3 | 35 | 2.5 |
| NK | S15-50 | 32.2 | 34.2 | (2) | 10-6 | 1 | 35 | 1.7 |
| NK | S18-84 | 34.2 | 41.9 | (3) | 10-7 | 2 | 34 | 2.7 |
| PIONEER | 1981 | 27.8 | 34.6 | (7) | 10-8 | 3 | 32 | 2.1 |
| PRIDE | B152 | 31.5 | -- | - | 10-4 | -1 | 32 | 1.7 |
| SCOTT | L1808 | 28.7 | -- | - | 10-4 | -1 | 28 | 1.9 |
| TERRA | RUNNER | 28.4 | -- | - | 10-8 | 3 | 29 | 1.3 |
| LSD(.05) | | NS | | | | 3.4 | 4.9 | 0.99 |
| TEST MEAN | | 30.35 | | | 10-3 | -1.4 | 30.8 | 1.86 |
| CV | | 14.6 | | | | 5.5 | 8.6 | 28.7 |

+ CHECK VARIETY USED TO CALCULATE DEVIATION FROM STANDARD MATURITY.

TABLE 7. Upper Peninsula (Alger County).

| Brand | Entry | Yield (Bu/A) | | | Maturity | Height | Lodging Score |
|-----------|-------------------|--------------|------|-----|----------|--------|------------------|
| | | 1985 | Avg. | (N) | | | |
| PUBLIC | BICENTENNIAL (00) | 27.3* | 28.7 | (2) | +5 | 18 | 1.5 |
| PUBLIC | CHICO (0) | 11.5 | — | — | +25 | 19 | 1.6 |
| PUBLIC | CLAY (0) | 15.8 | 17.0 | (3) | +12 | 15 | 1.8 |
| PUBLIC | MAPLE AMBER (00) | 26.0* | 31.0 | (3) | -1 | 17 | 1.0 |
| PUBLIC | MAPLE RIDGE (00) | 22.3 | 22.9 | (2) | -3 | 15 | 1.0 |
| PUBLIC | MCCALL (00) | 26.0* | 27.0 | (3) | 0 | 16 | 1.1 |
| LSD (.05) | | 3.3 | | | | | |
| TEST MEAN | | 21.5 | | | | | |

*Not significantly different from highest yield in that column.



MSU is an Affirmative Action/Equal Opportunity Institution. Cooperative Extension Service programs are open to all without regard to race, color, national origin, sex, or handicap.

Issued in furtherance of Cooperative Extension work in agriculture and home economics, acts of May 8, and June 30, 1914, in cooperation with the U.S. Department of Agriculture. W.J. Moline, Director, Cooperative Extension Service, Michigan State University, E. Lansing, MI 48824.

This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by the Cooperative Extension Service or bias against those not mentioned. This bulletin becomes public property upon publication and may be reprinted verbatim as a separate or within another publication with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company.

Revised (destroy previous edition) 1:87-3.5M-KMF-SP, 55¢, For Sale Only