

## **MSU Extension Publication Archive**

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Turf Tips for the Homeowner: Dandelion Control in Lawns

Michigan State University

Cooperative Extension Service

Michael Barrett, Weed Control Specialist, Department of Crop and Soil Sciences

May 1983

2 pages

The PDF file was provided courtesy of the Michigan State University Library

**Scroll down to view the publication.**





# Turf Tips

for the homeowner

Extension Bulletin E-1452

May 1983

## Dandelion Control in Lawns

Michael Barrett  
Weed Control Specialist  
Department of Crop and Soil Sciences  
Michigan State University



Dandelion in bloom (left), and in seed production stage (right).

Dandelion is a very common and familiar weed found in Michigan lawns. It is a cool season perennial, flowering from early spring to fall, although the yellow dandelion flowers are most abundant in April and May. The plants have a thick, fleshy taproot, often branched, which can penetrate several feet

into the soil. New shoots can sprout from this taproot.

Hundreds of seeds, each with a white, hairlike tuft of fibers borne on a minute stalk, are also produced by the dandelion plant. Dandelion reproduces and spreads by these seeds which are easily windborne. Seedlings appear in the spring and

summer, but are most numerous in the fall.

### Cultural Control

Instead of using herbicides, many people prefer to dig or cut the plants for control. For digging to be suc-



cessful, it must be performed in the spring (April or May). Dandelion roots have the lowest amount of stored food in the spring and are less able to resprout following cutting. Fall digging will not give good control, as root food storage is at its peak. The dandelion roots should be cut as deeply as possible, preferably 4 to 5 inches. Shallow cutting does not provide reliable dandelion kill because sufficient root material remains for resprouting. Dandelion plants not killed by deep, early cutting can produce numerous new shoots and may result in an increased weed problem.

Once dandelions are under control, maintain a thick, well fertilized, and managed lawn. A healthy lawn will best resist the invasion of dandelion and many other weeds.

## Chemical Control

Dandelions and many other broadleaf weeds (not grasses) are readily controlled by applications of 2,4-D. The herbicide should be applied to the foliage of the weeds during periods of active weed growth. That is, when there is ample soil moisture and moderate temperatures. Do not spray when rain is predicted within 12 hours.

Fall applications are best for

dandelion control because lawn grasses can fill spaces left from removal of the dandelions. Spring applications will control dandelions, but annual weeds, such as crabgrass, can fill the bare areas. If treating in the spring, spray before dandelion flowers appear. Fall applications also present less hazard of 2,4-D injury to desirable plants than spring treatments. Desirable trees and shrubs are growing less in the fall than in the spring and are less sensitive to 2,4-D. Also, many sensitive home vegetable plants are not present in the fall. However, care must be taken not to apply 2,4-D to desirable vegetation to avoid any possible injury to these plants.

Use only "2,4-D amine" formulations, not "2,4-D ester" products for spraying. The formulation is indicated on the label. Ester products are more volatile and the 2,4-D ester vapors can injure nearby vegetation. Spraying should only be done on days without wind. Avoid spray contact with desirable vegetation.

Sprayers used for 2,4-D applications should not be used for other purposes. Complete removal of 2,4-D from a sprayer is difficult, and minute amounts of 2,4-D can injure many plants. If you must use the sprayer for other purposes, best sprayer cleaning is obtained by rins-

ing twice with ammonia, including spraying through the nozzle. Follow with three rinses of water.

Check the herbicide label for the amount of product to use and be sure to follow the directions. Spray a mixture of herbicide and water onto the weed leaves until the solution begins to run-off. Do not apply excessive amounts of herbicide, as high rates will burn off only the top growth and will not kill the whole plant. Correct rates will move throughout the plant killing roots as well as shoots. For products containing 4 lb/gal 2,4-D (active ingredient) apply 1 qt per acre or 4½ teaspoons per 1000 square feet.


Fertilizer and other granular materials are also available mixed with 2,4-D. Follow the product directions for proper rates. Good control is obtainable when these products are properly used.

Other broadleaf weeds will be killed by 2,4-D treatment. Check the herbicide label for the exact species.

Herbicides are very effective tools for dandelion control. However, they must be used with respect. All agricultural chemicals should only be used in accordance with the instructions and limitations found on the label.

See Extension Bulletin E-653 for further information.

MICHIGAN STATE UNIVERSITY



COOPERATIVE  
EXTENSION  
SERVICE

MSU is an Affirmative Action/Equal Opportunity Institution. Cooperative Extension Service programs are open to all without regard to race, color, national origin, or sex.

Issued in furtherance of cooperative extension work in agriculture and home economics, acts of May 8, and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Gordon E. Guyer, Director, Cooperative Extension Service, Michigan State University, E. Lansing, MI 48824.

This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by the Cooperative Extension Service or bias against those not mentioned. This bulletin becomes public property upon publication and may be reprinted verbatim as a separate or within another publication with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company.

2P-10M-5:83-MP, Price 10 cents. Single copy free to Michigan residents.

File 22.41, 34.23