## **MSU Extension Publication Archive**

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Baby Food From Your Kitchen Michigan State University Extension Service Bureau of Health Services, Michigan Department of Public Health; Michigan State Unviersity Cooperative Extension Issued February 1978 4 pages

The PDF file was provided courtesy of the Michigan State University Library

Scroll down to view the publication.



Breast milk is the natural food for a young baby. If you are not nursing then several types of commercial formulas or an evaporated milk formula are the main food for baby's first five months. Whole milk can be introduced in the sixth month if you don't wish to continue formula or breast milk.

As baby grows and develops foods other than breast milk or formula are added to the diet. Your decision to feed "solid" food should be based upon the ability of your baby to swallow. Generally, most babies are ready to take solid foods from a spoon, in addition to liquids, sometime between the fourth and sixth month. Solids are fed by a small spoon and only a little will be eaten at first. As baby eats more solid food then the amount of formula or milk will usually decrease to about 16 to 24 ounces by the first birthday.

Many commercial baby foods are available. You may choose to use only commercial products or you may choose to make some or all of your own baby foods. Home prepared baby foods allow you to add a personal touch and help prepare your baby for the taste of foods that he will be eating later. Special modification in texture lasts only a short time since most babies can handle chopped or cooked finger foods by 10 months of age.

As you make baby food you will change the foods in consistency, texture and taste to suit your baby's

needs. The texture and consistency of foods for baby are altered by pureeing in a blender, or mashing in a sieve or food grinder, or with a fork or potato masher. Foods for a young baby should be *bland* and not seasoned by added salt, sugar, herbs and spices.

Electric blenders reduce fruit, vegetables and cooked meat to a puree. Initially you LIQUEFY and blend to desired consistency by adding liquid and lengthening blending time. Follow the directions outlined in your blender book. As baby develops more chewing ability you can reduce blending time or speed to produce desired consistency or you can use a grinder or mash.

When fixing home prepared baby foods use your common sense and remember the following:

use clean hands and equipment

use fresh food that is washed well prior to preparation

or meat that is freshly purchased

or frozen foods (held at or below 0°F)

or home or commercially canned fruit which is packed in water or light syrup

use clean storage containers (covered and suitable for refrigerator or freezer)

have appropriate refrigerator and freezer temperatures (37 to 39°F in refrigerator and 0°F or below in freezer)

Prepared by the

BUREAU OF PERSONAL HEALTH SERVICES, Department of Public Health, in Association with Michigan State University, Cooperative Extension Service



# FOOD CHOICE

#### CEREAL

Use single grains first, before combinations. Rice, barley and oat are less allergenic than wheat and corn.



#### FRUIT AND FRUIT JUICES

Use single cooked fruits first, before combinations. Frozen, fresh or canned juices should be diluted with equal parts of water when first introduced and then the water gradually decreased.



#### **VEGETABLES**

Use single cooked vegetables before combinations. Avoid using canned vegetables or those frozen vegetables with added salt.



### MEAT AND OTHER PLANT PROTEIN SOURCES

Introduce lamb, veal, beef, pork and poultry and then meat substitutes such as cooked dried peas, beans and lentils. Offer cooked egg yolk before whole egg.



### BREAD AND OTHER PRODUCTS

Use products that are made with enriched or whole grains. Limit the use of products containing sugar or other sweeteners.

## S FOR BABY

Iron is the nutrient most needed by infants and toddlers. Commercially prepared dry infant cereals are fortified with iron to a greater amount than hot cereals available for the family. Some brands of cooked cereal have more iron fortification. These are noted below with an asterisk(\*). If you don't use special infant cereals then suitable ones for infants include:

Cream of Rice (rice), Oatmeal (oat), Cream of Wheat\* (wheat), Malt-O-Meal \* (wheat and malt) and Corn Meal (corn).

When making home prepared hot cereals, omit salt and make with milk or formula. You will need to add more liquid to the cereals than box directions say when you first begin to feed them to your baby.

Fruits provide vitamins and some minerals to the baby's diet. Commercial infant juices are fortified with vitamin C so when you choose regular juice for your baby you need to select either those with natural vitamin C (such as orange, grapefruit or tomato) or those that are fortified with vitamin C. Read the juice container label.

Commercial infant fruit is available in unsweetened or sweetened varieties. Choose the fruits without added sugar, corn syrup or starch. Fruits are usually introduced after cereal because their texture is finer than the other food groups.

Ripe banana can be well mashed with a fork. Home prepared or canned applesauce can be used as is, or diluted slightly with warm water. Cooked or canned fruits can be drained and run through a sieve, food mill, masher or blender. Ripe, fresh fruit without skins, pits or seeds can be used if blended.

Many vitamins and minerals are found in vegetables; baby may need several opportunities to accept the different flavors and textures. Commercial infant vegetables may have other ingredients added to them. All cooked vegetables can be modified in texture and consistency. The stronger flavored ones (such as broccoli and cauliflower), the fibrous ones (such as celery and rutabaga) and white potatoes should be delayed until the baby has good chew and swallow reflexes. Sweet potatoes, winter squash, carrots, green beans, asparagus tips and spinach are easy to modify in consistency. Baby can progress to peas, beets, broccoli, lima beans, cauliflower and many others. Cooked vegetables can be pureed, chopped or mashed or used as finger foods.

Meat provides protein, vitamins and minerals. Babies consuming 24 ounces of milk or formula will not need more than a tablespoon or two of meat products each day to meet recommended protein levels. Commercial infant meats are available singly or combined with vegetables or with starch products. Meat chosen for baby should be tender, juicy inside pieces, free of gristle, fat and bone.

Preparing meat for baby usually requires a blender in order to produce a suitable consistency. Good protein sources that are easy to fix include cottage cheese, hard or soft cooked egg yolk, moist cooked ground beef, lamb or veal loaf, liver spread or cooked boneless fish.

Teething babies may like dried, toasted bread slices to chew. Crackers can be given after baby can bite foods off and chew well. Other bread and bakery items, and pasta products are not needed by baby in his first year.

Sweets and desserts are not appropriate for babies because their limited stomach capacity should be filled only by foods contributing to nutritional well-being and development.

Making your own baby food is not always less expensive than buying prepared baby food. It is satisfying to know what ingredients are being used. You can take food out of the amount cooked for the family and prepare for baby as you serve the family meal or you can batch prepare baby food.

If you prepare more than will be used in 24 hours then quickly freeze it. Spread it out on a tray in a thin layer or in ice cube trays or in small jars. As soon as food in trays is partially solid, slice or section into portions and place in freezer bags or freezer paper, date and label. Use within two months. To preserve nutrients defrost foods in refrigerator and then warm on the stove or in a microwave oven.

Amounts of food produced by blending varies depending on the amount of liquid added and the natural water present in the food. As your baby grows and develops more feeding skill you will vary the consistency by adding less water and/or decreasing the blending time or you can mash or finely chop the food. A typical serving of home prepared baby food for babies 6-9 months is usually ¼ cup cooked cereal, 2 to 4 tablespoons (½-¼ cup) fruit and vegetable, 1 to 2 tablespoons meat and 4 ounces (½ cup) juice.

One 4-ounce banana mashed, will yield about ½ baby food jar portion — at 26 cents/pound, a serving costs 6½ cents.

Sample Recipe Yields:

Eight ounces of frozen peas, cooked and blended with 3 tablespoons of water yields almost 2 baby food jars — at 69 cents/24 ounce package, a serving costs 6 cents.

One pound of raw carrots, washed, peeled, cooked and blended with three tablespoons of water, yields about 2½ baby food jars — at 39 cents/pound, a serving costs about 8 cents.

A 29-ounce can of pears in light syrup (10 halves) drained and blended, yields 2¾ baby food jars of pureed pears — at 69 cents/can, a serving costs 13 cents.

Making your own baby food is a satisfying experience that need not require a lot of time and effort.







Cooperative Extension Service Programs are open to all without regard to race, color, creed, or national origin. Issued in furtherance of cooperative extension work in agriculture and home economics, acts of May 8, and June 30, 1914, in cooperation with the U. S. Department of Agriculture. Gordon E. Guyer, Director, Cooperative Extension Service, Michigan State University, E. Lansing, MI 48824. IP-8M-2:78-St. Price 10 cents. Single copy free to Michigan residents.