

MSU Extension Publication Archive

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Chrysanthemum Pot Plant Production

Michigan State University Cooperative Extension Service

Meriam G. Karisson, Royal D. Heins, William H. Carlson, John A. Biernbaum,
Horticulture

Issued July 1987

12 pages

The PDF file was provided courtesy of the Michigan State University Library

Scroll down to view the publication.

Cooperative Extension Service
Michigan State University
Extension Bulletin E-2069 (New), July 1987

Chrysanthemum

Pot Plant

Production

by Meriam G. Karlsson, Royal D. Heins,
William H. Carlson and John A. Biernbaum,
Department of Horticulture

I. Introduction

- A. *Chrysanthemum morifolium* Ramat. is a complex hybrid of several species that grow wild in China and Japan. Two species are important contributors.
1. *C. indicum*, a native of China and Southern Japan with small yellow ray inflorescences.
 2. *C. sinense*, a species with white ray and orange disc flowers.
- B. Other species such as *C. coreanum*, *C. articum*, and *C. nipponicum* have also contributed flowering types and extended flowering time.
- C. In 1986, the pot chrysanthemum was the number two flowering pot crop in total sales, second only to Poinsettia.
1. In 1986, 33.4 million pots of chrysanthemums were produced with a value of \$87.3 million.
 2. Michigan chrysanthemum production for 1986 was valued at \$5.7 million.
 3. Based on dollar value of the crop, the leading producers in 1986 were California (27%), New York (10%), Ohio (9%), Texas (9%), Florida (9%), Michigan (6%), Illinois (5%), and North Carolina (4%).
- D. Gus Poesch, Alex Laurie and Ken Post, while at Michigan State University, made the first contributions in photoperiodic response of chrysanthemums by observing flower delay due to light.
- E. Year round production started in the late 1940s because of:
1. Daylength—lighting research work.
 2. Breeding of new varieties and increased commercial cutting production.

- F. The pot chrysanthemum is a popular crop all year round but also has large holiday sales.
- G. Pot production primarily includes pinched plants, but some single flower plants are also grown.
1. Production by pot size nationally is estimated at:

a. 4 in. pots	10%
b. 5 in. pots	5%
c. 6 in. pots	80%
d. Miscellaneous sizes	5%

II. Cultivars

- A. Chrysanthemum cultivar development in the U.S. initially centered around Elmer D. Smith of Michigan who exhibited his first seedling in 1889. Eventually, he introduced over 500 varieties.
- B. Development continued by many breeders over the years. Currently, Yoder Brothers and Pan American Plant Company have active breeding programs in the U.S. Breeding programs in The Netherlands and Japan are oriented primarily towards developing cut flower varieties.
- C. Current breeding objectives include:
1. Uniform development under a wide temperature range (high and low temperature tolerance).
 2. Faster flowering from the start of short days.
 3. New colors and flower types.
 4. Resistance to insects and diseases.
- D. Summary of cultivar characteristics:
1. Greatest production in the U.S. is of the following cultivars.
 - a. Bright Golden Anne (yellow decorative)
 - b. Surf (white daisy)
 - c. Circus (pink daisy)
 - d. Yellow and Sunny Mandalay (yellow decorative)
 2. Color

a. yellow	45%
b. white	25%
c. pink	18%
d. bronze	12%
 3. Flower type

a. decorative	73%
b. daisy	20%
c. other	7%
 4. Response group

a. 8 week	30%
b. 9 week	40%
c. 10 week	30%

- E. Flower shapes vary. Flowers can be classified as single (daisy), anemone, pompon, decorative, or under some miscellaneous classification based on petal shape (Spider, Fuji, Quill, or Spoon).

III. Flowering Requirements

- A. The chrysanthemum is classified as a short day plant, i.e., plants require a long dark period for both induction and development.
- B. The chrysanthemum has two critical photoperiods, one for flower initiation and one for flower development. Flower development requires a longer night period than initiation. The exact length of critical photoperiods for initiation and development depends on cultivar and temperature. Figure 1 shows the relationship between the natural photoperiod at 42° N latitude and the critical photoperiods for certain chrysanthemum cultivars (7).
- C. In general, the longer the response group, the longer the required dark span (night length) for both initiation and development. Some short response group varieties will initiate and develop a flower under a photoperiod unsuitable for initiation in a long response group variety. Examples of how critical photoperiods vary for chrysanthemum varieties grown at 60°F night temperature are given in the following table (1).


Variety	Response Group (weeks)	Initiation		Development	
		hr. light	hr. dark	hr. light	hr. dark
White Wonder	6	16.00	8.00	13.75	10.25
Pristine	8	15.25	8.75	12.00	12.00
Encore	10	14.50	9.50	12.00	12.00
Fortune	12	13.00	11.00	12.00	12.00
Snow	15	11.00	13.00	10.00	14.00

- D. The critical photoperiods are also temperature dependent. In general, as temperature increases or decreases from an optimum, the number of dark hours (night length) increases for initiation and further development. The effect of temperature on critical photoperiods in chrysanthemum is shown below (1).

Variety	Response Group (weeks)	Night Temp (°F)	Initiation		Development	
			hr. light	hr. dark	hr. light	hr. dark
White Wonder	6	50	13.75	10.25	*	*
		60	16.00	8.00	13.75	10.25
		80	16.00	8.00	12.00	12.00
Encore	10	50	13.75	10.25	13.75	10.25
		60	14.50	9.50	13.00	11.00
		80	15.25	8.75	12.00	12.00
Snow	15	50	12.00	12.00	12.00	12.00
		60	11.00	13.00	10.00	14.00
		80	10.00	14.00	9.00	15.00

*No flower development occurred at this temperature.

Figure 1.



E. Under short day conditions, temperature and light levels influence time to flower.


1. Three groups have been observed based on their temperature response during short day conditions. All three groups initiate and develop flowers most rapidly near 60°F night temperature (Figure 2).
 - a. Thermozero: Defined as cultivars that will flower earliest at 60°F night temperature with only a slight delay in flowering as temperatures decrease to 50°F or increase to 80°F.
 - b. Thermopositive: Defined as cultivars that will initiate and develop flower buds rapidly from 60° to 80°F night temperature. Temperatures below 60°F significantly delay development.
 - c. Thermonegative: Defined as cultivars that are delayed by temperatures above 60°F. Flower initiation and development occur most rapidly between 50° and 60°F.
2. Day temperature, night temperature and light level interact to influence the time to flower of a chrysanthemum cultivar.
 - a. Under the same temperature regime, the required time for flowering will decrease as the light level increases.

b. The optimum day and night temperature combination for quickest development is dependent on light level. As the light level increases, the optimum day temperature slowly increases while the optimum night temperature for development decreases.

F. When the flower buds show color, short day treatment can be discontinued. From this stage of development, the flowers will open normally under either long or short days.

G. Regardless of photoperiod and temperature, a chrysanthemum shoot will eventually initiate a flower. The longer the photoperiod or higher the temperature, the longer it will take to initiate the flower (i.e., more leaves will be formed before the flower initiates). If conditions are not proper for development after initiation, the bud will develop into a crown bud. Unintentional initiation under long days is normally not a problem under commercial conditions except in certain cultivars. Shown on the next page is the number of leaves initiated before the flower on five varieties of chrysanthemum grown under long days (natural daylength plus 5 hr night break) (3).

Figure 2. The number of days from start of short days to flower for a thermopositive, thermonegative and thermozero variety planted in early January from stock plants kept at 60° F. The plants were grown in a night temperature range from 50° to 80° F (2).



Cultivar	Number of Leaves on Main Plant Shoot		
	Date of Planting		
	6-13-73	10-10-73	5-29-74
Tuneful	45.3	90.3	56.9
Gold Crystal	44.0	69.2	49.5
Polaris	33.5	56.1	40.8
Bluechip	29.9	48.4	33.8
Bright Golden Anne	20.3	34.3	18.4

H. During the long nights of fall, winter, and early spring, plants can be maintained in a vegetative state through the use of low intensity night lighting.

1. Traditionally, incandescent lights have been used as a night interruption for 4 hr from 2200-0200 at 10 fc (foot candles). Light from fluorescent and high pressure sodium lamps at the same intensity and duration can also be used.
2. Cyclic lighting is effective as well. Six minutes of lighting per 30 minutes during the 4 hr period from 2200-0200 is as effective as continuous lighting. Cyclic lighting would normally not be used with high pressure sodium lamps due to the long "warm up" times of the lamps and reduction in lamp life due to repeated starts.

IV. Environmental Requirements

A. Light Duration

The following recommendations apply to northern U.S. latitudes.

1. From the beginning of September to the middle of April, a 4 hr night interruption is necessary to maintain plants in the vegetative state.
2. From the middle of April to the beginning of September, plants must receive artificial short days to initiate and develop flowers.

B. Light Intensity

1. Full sunlight is required for maximum growth, especially during the winter.
 - a. Some shade may be used from May 15 to September 15—mainly to reduce temperature.
 - b. Shading of plants in flower is necessary during the summer to prevent petal burn, especially on disbud varieties.
2. Uniformity of flowering is greatly reduced under low light intensities.
3. Supplemental light (500 to 1,000 fc) under low natural light conditions results in a higher quality plant and reduced time to flower.

a. Supplemental lighting is most beneficial on crops started during the period from early October through early March. Lighting time and duration depends on stage of crop development.

- 1) Long day period—light 20 hr per day (0200-2200 hour).
- 2) Short day period (first 2 to 3 weeks)—light 10 hr per day (0700 to 1700 hour).
- 3) Short day period (last 5 to 8 weeks)—normally not lit due to spacing of plants and reduced plant response.

C. Temperature

1. During vegetative growth (long days), maintain 68°F night temperatures.
2. Maintain 62° to 65°F night temperatures during flower initiation up through visible bud. Lower temperatures affect bud set and delay the crop.
3. High temperatures, especially above 70°F, cause heat delay in flower bud development.
4. After plants have reached visible bud, force at 60° to 62°F until color. Night temperature can then be lowered to 56° to 58°F.
5. The relationship between day and night temperature is the primary factor influencing shoot length. A large difference between day and night temperatures, with day temperature being the higher temperature, will result in longer internodes. High day and night temperatures result in more leaves beneath the flower (delayed flower initiation), producing more internodes and taller plants. (See section IV.C.3. above.)
6. Figure 3 shows the influence of day and night temperatures in the range from 50° to 86°F on shoot length for the cultivar 'Bright Golden Anne'. Light intensity does not greatly influence stem elongation unless plants are crowded. Plants will then be taller (6).

D. Water

1. Chrysanthemums require large amounts of water, especially during the summer.
2. Avoid overwatering new plants during the winter.
3. Allow the soil to dry somewhat between waterings. Water adequately so some leaching occurs. This requires about 12 fl oz (350 ml) per 6 in. pot, 4 to 5 fl oz (120 to 150 ml) per 4 in. pot.




Figure 3.

E. Fertilizer

1. A constant liquid feed of 200 ppm of nitrogen and potassium is recommended for mineral soils and 250 to 300 ppm nitrogen and potassium in soilless mixes. Higher rates may be necessary during the winter when plants dry out slowly.
2. An application of 500 ppm nitrogen and potassium is often recommended immediately after planting rooted cuttings or 7 to 10 days after sticking unrooted cuttings.
3. Phosphorus is normally added in the form of superphosphate prior to planting, 2½ lb/cu yd. Phosphorus readily leaches in a soilless medium. Therefore, fertilize with phosphorus at 100 ppm at least every 4 to 5 weeks.
4. A mixture of ammonium and nitrate nitrogen is preferred during most growth. Use mainly nitrate nitrogen (60% or more) during late fall and winter due to slow conversion of ammonium to nitrate.
5. In soilless mixes, include a source of trace elements.
6. Finish off crops with KNO_3 —or terminate fertilization no later than first flower color to improve postharvest quality.

F. Carbon Dioxide

1. CO_2 is most valuable during vegetative and early reproductive growth.
2. CO_2 levels of about 1,000 ppm early in the crop can result in heavier stems, larger flowers, and quicker flowering

compared with plants not receiving additional CO_2 .

3. Increased CO_2 levels in combination with high temperatures can result in excessive stem elongation.

V. Cultivation

A. Propagation

1. Propagation is entirely by cuttings and is normally done by specialists.
2. These propagators use culture-indexed, disease-free stock plants. Culture indexing has been used to obtain plants free of *Verticillium* and bacterial wilt, stunt virus and other viruses.
3. Rooted and unrooted cuttings are available.
 - a. Unrooted cuttings are easier and faster to plant than rooted cuttings due to the lack of roots. They are also cheaper to buy.
 - b. Unrooted cuttings require a special environment and extra time as compared to rooted cuttings to allow for the formation of roots.
4. Propagation procedure
 - a. Terminal cuttings 2 to 3 in. in length are normally rooted under intermittent mist. Cuttings can also be rooted under plastic without mist as long as high temperatures are avoided under the plastic.
 - b. Long photoperiods are used to maintain cuttings in the vegetative state.
 - c. Under low light conditions, high intensity discharge (HID) lighting accelerates rooting and a sturdier rooted cutting will develop.
 - d. Rooting occurs in 7 to 10 days with moderate temperatures (70° to 75°F). Use bottom heat.
 - e. Cuttings can be rooted in a peat-lite medium and transplanted or rooted directly in the pot.
 - f. Rooted cuttings can be stored at 31° to 33°F for 2 to 3 weeks if kept moist, but not excessively wet.

B. Media and Planting

1. A coarse and porous medium is preferred.
2. A pH of 6.2 to 6.4 is ideal. Foliar problems can occur due to micronutrient deficiencies above pH 6.8.
 - a. Adjust pH if needed—2½ lb ground limestone/cu yd raises pH ½ to 1 unit.
 - b. Check pH every 2 weeks throughout the entire life of the plant.

3. Pasteurize soil-based media at 180°F for 30 minutes with steam or at 140° to 160°F for 30 minutes with aerated steam.
4. Sort cuttings by size. Plant shallow with cuttings slanted towards the outside of the pot.
5. Four to six cuttings are planted per 5½-in. or 6-in. pot, depending on the market.
6. Freely breaking cultivars may require only 4 plants per pot, especially from Easter to Thanksgiving.

C. Spacing

Several spacing techniques can be used. Listed below are spacings commonly used for chrysanthemum production.

1. Two climate zero move system (2C0M).
Climate 1: 65°F—long days
Climate 2: 62° to 65°F—short days
In this system, the plants are placed at final spacing immediately after potting. The temperature and photoperiod are changed in the production area to achieve the second climate. This system has a low labor requirement but uses space inefficiently.
2. Traditional—two climate one move system (2C1M).
Climate 1: 65°F—long days
Climate 2: 62° to 65°F—short days
One production area is used for the long day environment. The plants are then moved to final spacing in a second production area. This system uses space more efficiently than the 2C0M system.
3. Unit spacing—two climate, two move system (2C2M).
Climate 1: 65°F—long days
Climate 2: 62° to 65°F—short days
Each unit consists of 3 benches in the short day area kept at 62° to 65°F. At the first move, the plants are placed on the middle bench of each unit. After 5 weeks (for a 10 week variety), the plants are spaced to the 2 outer benches of each unit.
 - a. Compared to the 2C1M system, this system is more space efficient.
 - b. It requires one additional spacing compared to the 2C1M system.
 - c. The system is designed for uniform year-round production of plants.
 - d. It does not easily allow for production of a holiday crop.
4. Zone system—3 climate, 2 move system (3C2M).
Climate 1: 65°F—long days
Climate 2: 62° to 65°F—short days (first 5-6 weeks)

Climate 3: 56° to 58°F—short days (to finish)

In this system, the plants are finished at a lower temperature (56°F) to improve quality.

- a. Compared to the 2C1M system, this system is more space efficient.
 - b. It requires two separate production areas during short days.
 - c. It also requires one additional moving of plants compared to the 2C1M system.
 - d. It is easier to fit holiday scheduling into production.
 - e. It may improve quality.
 - f. It saves energy through lower fuel costs.
- 5 The Space Utilization Chart (page 7) shows the four spacing systems described for production of 6-in. florist quality, 6-in. mass market quality, and 4-in. mass market quality using rooted or unrooted cuttings.

D. Support

1. No support is required unless stems are weak. Plants are then often wrapped with a string or a plastic hoop.

E. Pinching

1. The main purpose of pinching is to increase the number of flowers per plant.
2. The pinching schedule depends on time of year, cultivar, pot size and final plant height desired.
3. Time to pinch is based on growth of the plant.
 - a. Before pinching, establish a good root system (roots grown to the bottom of the pot).
 - b. Allow 1½ to 1¾ in. of new top growth.
4. In order to achieve desired plant height, it is necessary to pinch in relation to the beginning of short days according to each variety.
 - a. The following pinching schedule may be used as a guide for rooted cuttings (5):

Season and treatment	Days from planting to pinching	
	NORTH	SOUTH
LATE SPRING—EARLY FALL PLANTINGS:		
Tall treatment	7-10	7-10
Medium treatment	8-11	8-11
Short treatment	10-14	10-14
EARLY SPRING-LATE FALL PLANTINGS:		
Tall treatment	10-14	10-14
Medium treatment	11-14	11-15
Short treatment	14-17	14-17
MIDWINTER PLANTINGS:		
Tall treatment	14-17	10-14
Medium treatment	15-18	11-15
Short treatment	17-21	14-17

- b. The above schedule is modified when HID lighting is used.
- A soft pinch produces the best results. Remove the top 1/2 in. of the growing point leaving 6 to 8 leaves.
 - With short cultivars, the pinch is sometimes given 7 days before the start of SD while medium and tall cultivars are normally pinched at the start of SD.
 - Avoid a delayed pinch. A delayed pinch is one made after the start of SD. Breaks from a delayed pinch come from reproductive buds and produce sparse and narrow foliage. The resulting plants will appear very loose and open.

F. Disbudding

- Disbudding is used to modify the number of flower buds per shoot.
- Standard Disbudding—All lateral flower buds are removed, leaving only the terminal bud. It is done as soon as lateral

- flower buds are large enough to handle without damage to the terminal flower bud.
- Center Bud Removal—The terminal flower bud is removed. It is done on many daisy varieties and on plants in small pots. Normally requires one additional week to flower from planting.
 - Multiple Bud Removal (MBR)—Tip pinch each branch that develops from the first pinch when a minimum of 4 to a maximum of 6 leaves have developed on the breaks, or when you can feel the bud, but not see it yet.
 - Delays flowering.
 - Benefits of MBR:
 - Eliminates "clubbiness" on winter grown plants.
 - Increases headsize.
 - Can grow disbud varieties as spray.
 - Reduces height and eliminates need for a B-Nine application.

Space Utilization

Space utilization comparison of the 2 climate, 0 move (2COM), the 2 climate, 1 move (2C1M), the 2 climate, 2 move (2C2M), and the 3 climate, 2 move (3C2M) systems for production of 9-week chrysanthemum cultivars.


System	Long Day			Short Day						Total ft ² wk	ft ² wk savings compared to 2COM system (%)
	Spacing (in.)	Time (wk)	ft ² wk	First climate			Second climate				
				Spacing (in.)	Time (wk)	ft ² wk	Spacing (in.)	Time (wk)	ft ² wk		
<i>6 in. traditional spacing—unrooted cutting</i>											
2COM	14 x 14	3	4.08	14 x 14	9	12.25	—	—	—	16.33	—
2C1M	6 x 6	3	0.75	14 x 14	9	12.25	—	—	—	13.00	20
2C2M	6 x 6	3	0.75	10 x 10	4.5	3.13	14 x 14	4.5	6.13	11.38	30
3C2M	6 x 6	3	0.75	10 x 10	4.5	3.13	14 x 14	4.5	6.13	11.38	30
<i>6 in. traditional spacing—rooted cutting</i>											
2COM	14 x 14	2	2.72	14 x 14	9	12.25	—	—	—	14.97	—
2C1M	6 x 6	2	0.50	14 x 14	9	12.25	—	—	—	12.75	15
2C2M	6 x 6	2	0.50	10 x 10	4.5	3.13	14 x 14	4.5	6.13	9.76	35
3C2M	6 x 6	2	0.50	10 x 10	4.5	3.13	14 x 14	4.5	6.13	9.76	35
<i>6 in. tight spacing—unrooted cutting</i>											
2COM	12 x 12	2.5	2.50	12 x 12	9	9.00	—	—	—	11.50	—
2C1M	6 x 6	2.5	0.63	12 x 12	9	9.00	—	—	—	9.63	16
2C2M	6 x 6	2.5	0.63	8.5 x 8.5	4.5	2.26	12 x 12	4.5	6.13	7.39	36
3C2M	6 x 6	2.5	0.63	8.5 x 8.5	4.5	2.26	12 x 12	4.5	6.13	7.39	36
<i>6 in. tight spacing—rooted cutting</i>											
2COM	12 x 12	1.5	1.50	12 x 12	9	9.00	—	—	—	10.50	—
2C1M	6 x 6	1.5	0.38	12 x 12	9	9.00	—	—	—	9.38	11
2C2M	6 x 6	1.5	0.38	8.5 x 8.5	4.5	2.26	12 x 12	4.5	6.13	7.14	32
3C2M	6 x 6	1.5	0.38	8.5 x 8.5	4.5	2.26	12 x 12	4.5	6.13	7.14	32
<i>4 in.—unrooted cutting</i>											
2COM	7 x 7	2	0.68	7 x 7	9	3.06	—	—	—	3.74	—
2C1M	4 x 4	2	0.22	7 x 7	9	3.06	—	—	—	3.28	12
2C2M	4 x 4	2	0.22	5 x 5	4.5	0.78	7 x 7	4.5	1.53	2.53	32
3C2M	4 x 4	2	0.22	5 x 5	4.5	0.78	7 x 7	4.5	1.53	2.53	32
<i>4 in.—rooted cutting</i>											
2COM	7 x 7	1	0.34	7 x 7	9	3.06	—	—	—	3.40	—
2C1M	4 x 4	1	0.11	7 x 7	9	3.06	—	—	—	3.17	7
2C2M	4 x 4	1	0.11	5 x 5	4.5	0.78	7 x 7	4.5	1.53	2.42	29
3C2M	4 x 4	1	0.11	5 x 5	4.5	0.78	7 x 7	4.5	1.53	2.42	29

The 3C2M system differs from the 2C2M system by a change in temperature during the last climate.

G. Growth regulators

1. The quantity and timing of growth regulators is dependent on the season, the cultivar, the desired final plant height, and the greenhouse structure itself. General guidelines are listed below.
2. Spray 10 to 14 days after pinch or when shoots are 1 to 1½ in. long. A second application may be needed 2 to 3 weeks later.
3. The rate of stem elongation varies with stage of plant development. Apply growth regulators before the stem elongates most rapidly, within 4 weeks after start of short days. Figure 4 shows rate of stem elongation for the cultivar Bright Golden Anne at four different day temperatures (4).
4. Cultivars are classified according to treatment necessary to achieve a plant of desired height. Classifications of short, medium, and tall are based on the following criteria:
 - a. How quickly the plant forms roots.
 - b. Potential of plant for stem elongation.
 - c. Response of plant to growth regulators.
5. Most commonly used growth regulators in chrysanthemum production are B-Nine-SP (daminozide, SADH) and A-Rest (ancymidol).
6. B-Nine application is based on cultivar response:
 - a. Tall treatment varieties: 2 applica-
 - b. Medium treatment varieties: normally only 1 early application.
 - c. Short treatment varieties: 1 late application, if any.
7. Suggested concentration of B-Nine applications:
 - a. 2,500 ppm in winter
 - b. 3,750 ppm in spring and fall
 - c. 5,000 ppm in summer
8. To control height on extra-tall cultivars, a pre-plant dip or post-planting spray of B-Nine can be used in addition to recommended applications.
 - a. Dip plants in 1,250 ppm B-Nine just long enough to wet the stems and foliage. After planting let the foliage dry before watering. Used for rooted cuttings.
 - b. Spray 1,250 to 2,500 ppm 3 to 5 days after planting. Use for unrooted cuttings.
9. A-Rest sprays of 25 to 100 ppm give good height control.
 - a. Spray immediately after pinch to just wet the foliage.
 - b. Varieties requiring 2,500 ppm B-Nine require 25 ppm A-Rest.
 - c. Drench: 0.125 to 0.5 mg of A-Rest in 8 oz (240 ml) of water per 6 in. pot.
10. The response to B-Nine varies with cultivar and time of year. The cultivars shown in Figure 5 received one 2,500 ppm application of B-Nine.

Figure 4.






Figure 5.

VI. Problems

A. Diseases

1. Most diseases are eliminated by using culture indexed plants and growing in a pasteurized, well-drained soil.
2. Stem rot can be caused by *Pythium*, *Rhizoctonia*, and *Sclerotinia*. Control with fungicidal drenches.
3. *Botrytis* can be a problem especially on petals under conditions of high humidity.

B. Insects

1. Although there are potentially numerous insect problems, a preventive schedule of spraying and fumigation is often less costly than the losses that occur with infestation.
2. Common insect pests include aphids, spider mites, leafminers, thrips and caterpillars. The control chemicals are constantly changing due to introductions of new chemicals, labeling changes and the development of insect resistance. See Extension Bulletins E-1750 *The Glasshouse Ornamental Disease Control Handbook* and E-2014 *Insect and Mite Management in Commercial Greenhouses* or consult The Spartan Ornamental Network plant problem control program.

C. Physiological


1. Crown bud formations occurs when a bud is set under correct initiation conditions but the plant is not provided with correct developmental conditions. Vegetative shoots often surround the bud. Crown buds are caused by:
 - a. Light leaks or failure to apply black cloth consistently.
 - b. Insufficient light intensity during the long day period to keep plants vegetative.
 - c. Cuttings taken from shoots that were too long and thus initiated.
 - d. Too high temperature during short days.
2. Neckiness is a problem in which the flower stem below the bud is elongated and devoid of foliage. It can be prevented by an application of B-Nine within 1 to 3 days after disbudding. A later application may be ineffective.
3. Uneven flowering can be caused by several factors.
 - a. Light leaks in the black cloth.
 - b. Insufficient number of short days.
 - c. High temperatures.
 - d. Low temperatures (lower than 60°F) during flower initiation (first 3 to 4 weeks of short days).

- e. Low light during development.
- f. Ethylene pollution.

VII. Harvesting, Handling, and Marketing

- A. Maximize production light levels for good postharvest longevity. Low light levels during the final 3 to 4 weeks of production decrease longevity.
- B. Highest plant and flower quality occurs when temperatures are lowered to 56°F the last couple of weeks before sale.
- C. Termination of fertilizer (especially ammonium forms) at time of disbudding (approximately 3 to 4 weeks prior to marketing) extends longevity 10 to 14 days and acclimatized plants are more tolerant of shipping conditions.
- D. Maintenance of a good root system is critical for a long decorative life. Monitor watering closely to prevent overwatering.
- E. Leaf yellowing in some cultivars may be a direct result of over fertilization (especially with ammonium forms of nitrogen) rather than crowding.
- F. Flowers fail to open properly if plants are removed from the greenhouse too soon, i.e., when just showing color.
- G. A flower grows very rapidly during the last 10 days of development, often gaining 90% of its final dry weight. Inadequate light conditions during this period may result in black centers. Figure 6 shows an example of the amount of dry weight in the different plant organs over time for the cultivar 'Bright Golden Anne.' Note the large gain in flower dry weight during the last 10 days.
- H. In one experiment, flowering chrysanthemums shipped for 7 days at 70°F lasted 23 days under interior conditions while plants shipped at 75°F lasted for only 10 days. Plants shipped at 75°F were of poor quality after shipping.
- I. Maintaining potted chrysanthemums under interior conditions at 65°F increased longevity by 13 days compared to an interior temperature of 80°F.

Figure 6.



References

1. Cathey, H. M. 1955. A study of the effects of light and temperature upon the flowering of *Chrysanthemum morifolium* and *Tulipa gesneriana*. Ph.D. Thesis, Cornell University, Ithaca, New York.
2. Cathey, H. M. 1955. Temperature guide to chrysanthemum varieties. Bull. N.Y. St. Flower Grs. 119:1.
3. Cockshull, K. E. 1975. Premature budding in year-round chrysanthemums. Rep. Glasshouse Crops Res. Inst. 1974, 128-136.
4. Erwin, J. E. 1986. Environmental and physiological factors influencing the response of *Chrysanthemum morifolium* cvs. 'Bright Golden Anne' and 'Circus' to daminozide. M.S. Thesis, Michigan State University, E. Lansing, Michigan.
5. Hanzel, B., and W. H. Carlson. 1986. The pot mum pinch. How to do it. BPI News 17(4):1-2.
6. Karlsson, M. G., and R. D. Heins. 1986. Response surface analysis of flowering in chrysanthemum 'Bright Golden Anne.' J. Amer. Soc. Hort. Sci. 111:253-259.
7. Langhans, R. W. 1964. Light and photoperiod. In: Chrysanthemums. A manual of the culture, disease and insects and economics of chrysanthemum, p. 73-85. New York State Extension Service Chrysanthemum School.



MSU is an Affirmative Action/Equal Opportunity Institution. Cooperative Extension Service programs are open to all without regard to race, color, national origin, sex, or handicap.

Issued in furtherance of Cooperative Extension work in agriculture and home economics, acts of May 8, and June 30, 1914, in cooperation with the U.S. Department of Agriculture. W.J. Moline, Director, Cooperative Extension Service, Michigan State University, E. Lansing, MI 48824.

This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by the Cooperative Extension Service or bias against those not mentioned. This bulletin becomes public property upon publication and may be reprinted verbatim as a separate or within another publication with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company.