

## **MSU Extension Publication Archive**

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Controlling Field Bindweed and Hedge Bindweed

Michigan State University Extension Service

IPM Facts

Fred Salzman, Karen Renner, Jim Kells, Department of Crop and Soil Sciences

Issued December 1992

2 pages

The PDF file was provided courtesy of the Michigan State University Library

**Scroll down to view the publication.**

## Controlling Field Bindweed and Hedge Bindweed

Fred Salzman, Karen Renner, and Jim Kells  
Department of Crop and Soil Sciences  
Michigan State University

### *What is a Perennial Weed?*


A perennial weed is any weed capable of surviving for three or more years. Perennial weeds are characterized by vegetative reproduction. Vegetative reproduction in these species is due to (a) rhizomes - underground creeping stems commonly found in perennial grasses; (b) stolons - prostrate stems or runners on the soil surface with roots at the nodes; (c) creeping roots; (d) tubers - underground enlarged storage stems; or (e) bulbs - underground storage organs consisting of a stem axis covered with many overlapping leaf scales.

Perennial weeds may or may not reproduce by seed. They always, however, have the potential to reproduce by vegetative means.

### *Description of Field Bindweed and Hedge Bindweed*

Field bindweed and hedge bindweed are two closely related members of the morningglory family. They are not morningglories since they are perennial weeds while morningglories are annual weeds. Both bindweed species have extensive root systems. The roots of field bindweed may reach a soil depth of 20 to 30 feet while hedge bindweed roots are much more shallow. Stems are smooth, slender, and range in length from 2 to 10 feet. Leaves of field bindweed are 1 1/2 to 2 1/2 inches long and are oblong in shape with

arrowhead-like bases. Leaves of hedge bindweed are up to 4 inches long, more triangular in shape than field bindweed, with a sharp point at the tip. Flowers on both species are funnel-shaped and white to pink. Flowers of hedge bindweed are larger than those of field bindweed. Seeds are dark brownish to gray and dull with one rounded and two flattened sides. Plants reproduce by seeds, stems, and creeping rootstocks.



### *Method of Control*

Methods of perennial weed control fall into three categories: (a) cultural, such as crop rotation; (b) mechanical, tillage including various implements such as plows, disks, or cultivators; and (c) chemical, using herbicides. Control of perennial weeds may require a combination of all these methods. Consider the energy and environmental implications when choosing a method of control.

### *Mechanical Control*

Mechanical control may increase or decrease perennial weed infestations. Tillage may increase infestations by moving perennial weeds to new areas of the field or breaking dormancy of underground buds resulting in new shoot growth. Tillage during cool, wet conditions results in reduced control.

Tillage may decrease perennial weed infestations if done frequently enough to deplete underground root reserves. The field should be tilled every two or three weeks. Warm, dry soil conditions increase the effectiveness of tillage for perennial weed control by drying plant roots on the soil surface.

# Chemical Control of Field Bindweed and Hedge Bindweed

## Soybeans

<u>Herbicide</u> <sup>1</sup>	<u>Rate</u>	<u>Timing</u>	<u>Effectiveness</u>
Blazer + NIS <sup>2</sup>	1.5 pt/A + 1/4%	POST	Poor
Cobra + COC <sup>2</sup>	0.78 pt/A + 1 pt/A	POST	Poor

<sup>1</sup> These treatments cause top growth suppression only. Two applications of Blazer may be necessary. 2,4-DB may be added to Blazer and nonionic surfactant removed to increase activity. Read the label as 2,4-DB can injure soybeans.

<sup>2</sup> NIS = nonionic surfactant; COC = crop oil concentrate

## Corn

<u>Herbicide</u>	<u>Rate</u>	<u>Timing (Weed height)</u>	<u>Effectiveness</u>
2,4-D amine <sup>1</sup>	1 pt/A	8" runners	Fair
Banvel	1/2 pt/A	8" runners	Fair
Banvel + 2,4-D amine	1/4 pt/A + 1/2 pt/A	8" runners	Fair
2,4-D ester <sup>1</sup>	1 qt/A	Preharvest application (after brown silk stage of corn)	Good

<sup>1</sup> 2,4-D is effective only on hedge bindweed.

## Spot treatments and between crops

<u>Herbicide</u>	<u>Rate</u>	<u>Timing<sup>1</sup> (Weed stage)</u>	<u>Effectiveness</u>
Roundup	2%	Spot treatment (full bloom)	Good
Roundup	3 - 4 qt/A	Full bloom or beyond	Good
Banvel	1 - 2 qt/A <sup>2</sup>	Full bloom	Good
Roundup + Banvel	2 qt/A + 1 pt/A	Full bloom	Good

<sup>1</sup> Do not treat when bindweed is under drought stress.

<sup>2</sup> Banvel at 1 qt/A will provide suppression; 2 qt/A will provide control.

This bulletin was originally prepared with the support of the U.S. Department of Energy, Grant No. DE-FG0276CS60204. However, any opinions, findings, conclusions or recommendations expressed herein are those of the author(s) and do not necessarily reflect the views of DOE.

**To protect yourself and others and the environment, always read the label before applying any pesticide.**



MSU is an Affirmative-Action Equal Opportunity Institution. Cooperative Extension Service programs and materials are open to all without regard to race, color, national origin, sex, handicap, age or religion. ■ Issued in furtherance of Cooperative Extension work in agriculture and home economics, acts of May 8 and June 20, 1914, in cooperation with the U.S. Department of Agriculture. Gail L. Imig, director, Cooperative Extension Service, Michigan State University, E. Lansing, MI 48824. ■ This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by the Cooperative Extension Service or bias against those not mentioned. This bulletin becomes public property upon publication and may be reprinted verbatim as a separate or within another publication with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company. *Produced by Outreach Communications and printed on recycled paper using vegetable-based inks.*

This publication contains pesticide recommendations based on research and pesticide regulations. However, changes in pesticide regulations occur constantly. Some pesticides mentioned may no longer be available, and some uses may no longer be legal. If you have questions about the legality and/or registration status for using pesticides, contact your county Cooperative Extension Service office.

