

MSU Extension Publication Archive

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Controlling Jerusalem Artichoke

Michigan State University Extension Service

Michigan Energy Conservation Program for Agriculture and Forestry

Fred Salzman, Karen Renner, Jim Kells, Department of Crop and Soil Sciences

Issued May 1990

2 pages

The PDF file was provided courtesy of the Michigan State University Library

Scroll down to view the publication.



CONTROLLING JERUSALEM ARTICHOKE

Fred Salzman, Karen Renner, and Jim Kells
Department of Crop and Soil Sciences
Michigan State University

What is a Perennial Weed?

A perennial weed is any weed capable of surviving for three or more years. Perennial weeds are characterized by vegetative reproduction. Vegetative reproduction in these species is due to (a) rhizomes, underground creeping stems commonly found in perennial grasses, (b) stolons, prostrate stems or runners on the soil surface with roots at the nodes, (c) creeping roots, (d) tubers, underground enlarged storage stems, or (e) bulbs, underground storage organs consisting of a stem axis covered with many overlapping leaf scales.

Perennial weeds may or may not reproduce by seed. They always, however, have the potential to reproduce by vegetative means.

Description of Jerusalem Artichoke

Jerusalem artichoke grows from characteristic white tubers that form on the tips of the roots. The stems of the plant are rough-hairy, branch near the top, and can reach a height of 9 feet. Leaves are large with a rough upper surface, hairy lower surface, and can vary from egg-shaped to lance-shaped. Leaf edges are saw-toothed. Flowers are daisy-like with yellow centers and petals, and are 2 inches in diameter. Seeds are flattened, wedge-shaped, smooth, and often

mottled with black. In addition to tubers, Jerusalem artichoke can also reproduce from rhizomes, creeping rootstocks, and seeds.

Jerusalem artichoke can be found in moist fence rows, roadsides, woods, and can be very troublesome in cultivated fields.

Methods of Control

Methods of perennial weed control fall into

three categories. These categories are (a) cultural, such as crop rotation, (b) mechanical; tillage including various implements such as plows, disks, or cultivators, and (c) chemical, using herbicides. Control of perennial weeds may require a combination of all these methods. Consider the energy and environmental implications when choosing a method of control.

Mechanical Control

Mechanical control may increase or decrease perennial weed infestations. Tillage may increase infestations by moving perennial weeds to new areas of the field or breaking dormancy

of underground buds resulting in new shoot growth. Tillage during cool, wet conditions results in reduced control.

Tillage may decrease perennial weed infestations if done frequently enough to deplete underground root reserves. The field should be tilled every two or three weeks. Warm, dry soil conditions increase the effectiveness of tillage for perennial weed control by drying plant roots on the soil surface.



Chemical Control of Jerusalem Artichoke

Soybeans

<u>Herbicide</u> ¹	<u>Rate</u>	<u>Timing</u> (Weed height)	<u>Effectiveness</u>
Roundup	Rope-wick ²	POST (10-20")	Fair-Good
Classic ³ + NIS ⁴	¾ oz/A + ¼%	POST (8")	Good
Pursuit + 28% N + NIS ⁴	¼ pt/A + 1 qt/A + ¼%	POST (6-10")	Good

¹ These treatments control topgrowth only.

² Roundup can be applied with a ropewick more than once during the growing season if new Jerusalem artichoke plants reemerge or all plants are not above the soybean canopy at the time of the ropewick application.

³ Do not use Classic if soil pH is greater than 7.0.

⁴ NIS=nonionic surfactant

Corn

<u>Herbicide</u>	<u>Rate</u> ¹	<u>Timing</u> (Weed height)	<u>Effectiveness</u>
Banvel	½ pt/A and repeat	6" ²	Fair
Banvel + 2,4-D amine	¼ pt/A + ½ pt/A and repeat	6" ²	Fair
2,4-D amine	1 pt/A and repeat	6" ²	Fair
Atrazine	1½ lb/A and repeat	PPI+POST ³	Poor-Fair

¹ Two applications of any treatment are required for adequate results. These treatments may need to be repeated for 2 or more years. The second application will probably require a directed treatment.

² Treatments should be made when Jerusalem artichoke is 6 inches tall and repeated when the regrowth is 6 inches tall. The second application will probably require a directed treatment.

³ Add crop oil concentrate at 1 qt/A to the postemergence application.

Spot Treatments and Between Crops

<u>Herbicide</u>	<u>Rate</u>	<u>Timing</u> ¹ (Weed growth stage)	<u>Effectiveness</u>
Roundup	2%	Spot treatment (bud stage)	Fair-Good
Roundup	3 qt/A	Bud to bloom stage	Fair-Good
Banvel	2 qt/A	Bud to bloom stage	Fair-Good
2,4-D ester	2 qt/A	Bud to bloom stage	Fair-Good

¹ Retreatment is usually required for complete control.

This bulletin was prepared with the support of the U.S. Department of Energy, Grant No. DE-FG0276CS60204. However, any opinions, findings, conclusions or recommendations expressed herein are those of the author(s) and do not necessarily reflect the views of DOE.

MSU is an Affirmative Action/Equal Opportunity Institution. Cooperative Extension Service program are open to all without regard to race, color, national origin, sex or handicap.

This publication contains pesticide recommendations based on research and pesticide regulations. However, changes in pesticide regulations occur constantly. Some pesticides mentioned may no longer be available, and some uses may no longer be legal. If you have questions about the legality and/or registration status for using pesticides, contact your county Cooperative Extension Service office.

To protect yourself and others and the environment, always read the label before applying any pesticide.

Issued in furtherance of Cooperative Extension work in agriculture and home economics, acts of May 8, and June 30, 1914, in cooperation with the U.S. Department of Agriculture, J. Ray Gillespie, Interim Director, Cooperative Extension Service, Michigan State University, E. Lansing, MI 48824.

This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by the Cooperative Extension Service or bias against those not mentioned. This bulletin becomes public property upon publication and may be reprinted verbatim as a separate or within another publication with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company.