

MSU Extension Publication Archive

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Controlling Wirestem Muhly

Michigan State University Extension Service

Michigan Energy Conservation Program for Agriculture and Forestry

Fred Salzman, Karen Renner, Jim Kells, Department of Crop and Soil Sciences

Issued May 1990

2 pages

The PDF file was provided courtesy of the Michigan State University Library

Scroll down to view the publication.



CONTROLLING WIRESTEM MUHLY

Fred Salzman, Karen Renner, and Jim Kells
Department of Crop and Soil Sciences
Michigan State University

What is a Perennial Weed?

A perennial weed is any weed capable of surviving for three or more years. Perennial weeds are characterized by vegetative reproduction. Vegetative reproduction in these species is due to (a) rhizomes - underground creeping stems commonly found in perennial grasses; (b) stolons - prostrate stems or runners on the soil surface with roots at the nodes; (c) creeping roots; (d) tubers - underground enlarged storage stems; or (e) bulbs - underground storage organs consisting of a stem axis covered with many overlapping leaf scales.

Perennial weeds may or may not reproduce by seed. They always, however, have the potential to reproduce by vegetative means.

Description of Wirestem Muhly

The rhizomes of wirestem muhly are short, creeping, and very scaly. Shoots are erect or grow along the ground and become erect at the tips. Stems range in length from 2 to 3 feet. Stems are smooth below joints, very tough, leafy, branch freely, and often root at the lower nodes. Leaves are flat, rough, pale green, scattered along the stem, and dense near the tip, giving the plant a brushy appearance. The seedhead is a panicle that barely protrudes from the leaf sheath.

Wirestem muhly is found in fence rows and along roadsides. It sometimes persists in cultivated fields, but is most commonly found in no-till fields. Growth starts later in the spring than quackgrass. Wirestem muhly is more easily injured by frost than quackgrass.


Methods of Control

Methods of perennial weed control fall into three categories: (a) cultural, such as crop rotation; (b) mechanical, tillage including various implements such as plows, disks, or cultivators; and (c) chemical, using herbicides. Control of perennial weeds may require a combination of all these methods. Consider the energy and environmental implications when choosing a method of control.

Mechanical Control

Mechanical control may increase or decrease perennial weed infestations. Tillage may increase infestations by moving perennial weeds to new areas of the field or breaking dormancy of underground buds resulting in new shoot growth. Tillage during cool, wet conditions results in reduced control.

Tillage may decrease perennial weed infestations if done frequently enough to deplete underground root reserves. The field should be tilled every two or three weeks. Warm, dry soil conditions increase the effectiveness of tillage for perennial weed control by drying plant roots on the soil surface.



Chemical Control of Wirestem Muhly

Soybeans

<u>Herbicide</u>	<u>Rate</u>	<u>Timing</u> ¹	<u>Effectiveness</u>
Assure + COC ²	1 pt/A + 1%	POST	Excellent
Fusilade 2000 + COC ²	1½ pt/A + 1 qt/A	POST	Good
Poast + 28% N + Dash	1¼ pt/A + 1 gal/A + 1 qt/A	POST	Fair

¹ Apply when wirestem muhly is 4 to 6 inches tall. A second application of these treatments or cultivation 14 to 21 days later may be required for satisfactory control.

² COC = crop oil concentrate

Corn

<u>Herbicide</u>	<u>Rate</u>	<u>Timing</u> ¹	<u>Effectiveness</u>
Atrazine	1½ lb/A + 1 ½ lb/A	PPI + PRE	Fair
Atrazine	1½ lb/A + 1 ½ lb/A	PPI + POST	Fair
Atrazine	1½ lb/A + 1 ½ lb/A	PRE + POST	Fair
Atrazine	1½ lb/A + 1 ½ lb/A	POST + POST	Fair

¹ Atrazine effectiveness is greatest when applied POST. Add 1 qt/A of crop oil concentrate to all POST applications.

Spot treatments and between crops

<u>Herbicide</u>	<u>Rate</u>	<u>Timing</u> ¹	<u>Effectiveness</u>
Roundup	2 qt/A	Fall or spring	Good
Roundup	2%	Spot treatment (see label)	Good

¹ Apply when western muhly is 8 inches or less in height. Spring applications will delay planting. For fall applications, check plants for frost damage before application.

"This bulletin was prepared with the support of the U.S. Department of Energy, Grant No. DE-FG0276CS60204. However, any opinions, findings, conclusions or recommendations expressed herein are those of the author(s) and do not necessarily reflect the views of DOE"

MSU is an Affirmative Action/Equal Opportunity Institution. Cooperative Extension Service program are open to all without regard to race, color, national origin, sex or handicap.

This publication contains pesticide recommendations based on research and pesticide regulations. However, changes in pesticide regulations occur constantly. Some pesticides mentioned may no longer be available, and some uses may no longer be legal. If you have questions about the legality and/or registration status for using pesticides, contact your county Cooperative Extension Service office.

To protect yourself and others and the environment, always read the label before applying any pesticide.

Issued in furtherance of Cooperative Extension work in agriculture and home economics, acts of May 8, and June 30, 1914, in cooperation with the U.S. Department of Agriculture, J. Ray Gillespie, Interim Director, Cooperative Extension Service, Michigan State University, E. Lansing, MI 48824.

This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by the Cooperative Extension Service or bias against those not mentioned. This bulletin becomes public property upon publication and may be reprinted verbatim as a separate or within another publication with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company.