

## **MSU Extension Publication Archive**

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Control of Annual Grass Weeds in Soybeans

Michigan State University Extension Service

IPM Facts

Fred Salzman, Karen Renner, Department of Crop and Soil Sciences

Issued December 1993

4 pages

The PDF file was provided courtesy of the Michigan State University Library

**Scroll down to view the publication.**



# FACTS

Extension Bulletin E-2260

Revised December 1993

## Control of Annual Grass Weeds in Soybeans

**Fred Salzman and Karen Renner,  
Department of Crop and Soil Sciences  
Michigan State University**

### **Biology**

Annual grass weeds complete their life cycle in one year by germinating, flowering, producing seed, and dying. Reproduction is by seed only; there are no overwintering vegetative parts. Summer annuals complete their life cycle from spring to fall, while winter annuals complete their life cycle from fall to late spring. Most common annual grass weeds in Michigan are summer annuals.

### **Cultural Control**

Cultural control utilizes crop management practices that allow a desirable crop to out-compete weeds. Two common cultural control methods are the use of a smother crop and crop rotation. Smother crops compete with weeds for light, nutrients, and moisture. Common smother crops include forage sorghum, cereals (barley, oats, wheat, or rye) and legumes (alfalfa or clover), although any highly competitive crop that is well adapted to an area may be suitable.

Crop rotation prevents the buildup of weeds common to a specific crop. An ideal crop rotation includes crops that have vastly different growth habits and planting conditions compared to other crops in the rotation and to the predominant problem weeds. Smother crops can be included in the crop rotation.

### **Mechanical Control**

Tillage can be used for both ground preparation and weed control. Tillage equipment can be subdivided into two categories, primary and secondary. Primary tillage tools include moldboard plows, chisel plows, and heavy disks. Secondary tillage tools include various types of harrows, field cultivators, row crop cultivators, finishing disks, and rotary hoes.

Rotary hoes are used to remove shallow-rooted weeds prior to or shortly after crop emergence. Row crop cultivators are used to remove weeds between crop rows.

Tillage can influence weed species communities. For example, grasses and perennial weeds are more predominant in reduced tillage systems.

Shallow cultivation may be used to enhance herbicide applications. Cultivate as shallow as possible to prevent bringing weed seeds to the soil surface. Do not cultivate fields receiving preemergence herbicide applications for at least 2 weeks following application; however, if dry weather persists for 7 to 10 days following a preemergence application, rotary hoe the field. Delay cultivation after postemergence herbicide applications for at least 7 to 10 days to allow time for the herbicide to translocate throughout the sprayed weeds.

### **Chemical Control**


Chemical control of annual weeds can be achieved with herbicides applied preplant incorporated, preemergence, or postemergence. Consider the energy and environmental implications when choosing a control.

**Preplant incorporated** herbicides are mechanically incorporated into the soil prior to planting. Incorporation of some herbicides is necessary to prevent losses from volatilization or photodecomposition.

**Preemergence** herbicides are applied to the soil surface after the crop has been planted but before the crop seedlings emerge through the soil.

**Postemergence** herbicides are applied after crop emergence. They either burn off the above-ground portion of the weeds or they may be translocated throughout the weed and kill the growing points. Most postemergence herbicides are applied broadcast. However, certain herbicides can be applied with directed equipment such as drop nozzles when the crop or weeds are too large for broadcast applications.

# GRASS SEEDLING IDENTIFICATION GUIDE



This key is a way to identify grass weed seedlings. To use the key you need to understand how the key is organized and be able to recognize a few simple plant characteristics.

This key uses drawings to show which part of the plant to examine. Identifying characteristics are circled.

Examine grass seedlings with great care. One characteristic alone seldom identifies the weed.

Then begin on the left side of the key and proceed step by step to the right. Do not skip any steps.

# Grass Weed Response to Herbicides in Soybeans

Herbicide	Rate/A <sup>a</sup>	Barnyardgrass	Crabgrass	Giant foxtail	Green foxtail	Yellow foxtail	Fall panicum	Witchgrass	Wild proso millet
		Control <sup>b</sup>							
<b>Preplant incorporated:</b>									
Command	1 1/2 pt	G	E	E	E	G	G	G	F
Dual	2 pt	E	E	E	E	E	G	G	F
Lasso/Partner/Microtech	2 qt	E	E	E	E	E	G	G	F
Lexone/Sencor 75DF	1/2 lb	F	F	F	G	G	F	F	P
Preview 75DG	6 oz	F	F	F	G	G	F	F	P
Prowl 3.3 EC	2.4 pt	E	E	E	E	E	E	E	F
Pursuit	1/4 pt	F	F	G	G	G	P	P	F
Scepter	2/3 pt	F	P	G	G	G	P	P	P
Sonalan	2 1/2 pt	E	E	E	E	E	E	E	F
Treflan	1 1/2 pt	E	E	E	E	E	E	E	F
<b>Preemergence:</b>									
Dual	2 pt	E	E	E	E	E	G	G	F
Lasso/Partner/Microtech	2 qt	E	E	E	E	E	G	G	F
Lexone/Sencor 75DG	1/2 lb	F	F	F	G	G	F	F	P
Linex/Lorox 50DF	1 1/2 lb	F	F	F	F	F	F	F	P
Lorox Plus	14 oz	F	F	F	F	F	F	F	P
Preview	6 oz	F	F	F	G	G	F	F	P
Prowl 3.3 EC	2.4 pt	G	G	G	G	G	G	G	F
Pursuit	1/4 pt	F	F	F	F	F	P	P	P
Scepter	2/3 pt	F	P	G	G	G	P	P	P

<sup>a</sup> Refer to Extension Bulletin E-434, "Weed Control Guide for Field Crops," for more specific weed control recommendations.

<sup>b</sup> P = poor; F = fair; G = good; E = excellent

# Grass Weed Response to Herbicides in Soybeans

Herbicide	Rate/A <sup>a</sup>	Barnyardgrass	Crabgrass	Giant foxtail	Green foxtail	Yellow foxtail	Fall panicum	Witchgrass
		Control <sup>b</sup>						
<b>Postemergence<sup>c</sup></b>								
Assure II + COC <sup>d</sup> or NIS <sup>d</sup>	0.44 pt + 1% or 1/4%	G	G	E	E	E	E	E
Blazer + NIS <sup>d</sup>	1.5 pt + 1/8%	N	N	F	F	F	F	N
Fusilade 2000 + COC <sup>d</sup>	1 1/2 pt + 1 qt	E	G	E	E	E	E	E
Fusion + NIS <sup>d</sup> or COC <sup>d</sup>	1/2 pt + 1/4% or 1/2%	G	G	E	E	E	G	E
Option II + COC <sup>de</sup>	0.7 <sup>e</sup> pt + 1 qt	G	F	E	E	G	G	G
Poast + COC <sup>d</sup> or Dash	1 pt + 1 qt or 1 qt	E	G	E	E	E	E	E
Poast Plus + Dash	1 1/2 pt + 1 qt	E	G	E	E	E	E	E
Pursuit +28% N <sup>d</sup> + NIS <sup>d</sup>	1/4 pt + 1 qt + 1/4%	F	F	G	G	G	F	F
Reflex + NIS <sup>d</sup> or COC <sup>d</sup>	1 pt + 1/4% or 1%	P	P	F	F	F	P	P
Scepter +NIS <sup>d</sup> or COC <sup>d</sup>	1/3 pt + 1/4% or 1 qt	N	N	F	F	F	N	N
Select + COC <sup>d</sup>	1/2 pt + 1 qt	E	G	E	E	E	G	G

<sup>a</sup> Refer to Extension Bulletin E-434, "Weed Control Guide for Field Crops," for more specific weed control recommendations.

<sup>b</sup> N = none; P = poor; F = fair; G = good; E = excellent.

<sup>c</sup> Refer to labels for size of grasses controlled by each postemergence herbicide and specifics on application rates, spray pressure, gallonage, tank mixes, etc.

<sup>d</sup> NIS = nonionic surfactant; COC = crop oil concentrate; 28% N = 28% liquid ammonium nitrate

<sup>e</sup> Apply Option II at 0.5 pt/A for giant and green foxtail control.

This bulletin was originally prepared with the support of the U.S. Department of Energy, Grant No. DE-FG0276CS60204. However, any opinions, findings, conclusions or recommendations expressed herein are those of the author(s) and do not necessarily reflect the views of DOE.

**To protect yourself and others and the environment, always read the label before applying any pesticide.**

This publication contains pesticide recommendations based on research and pesticide regulations. However, changes in pesticide regulations occur constantly. Some pesticides mentioned may no longer be available, and some uses may no longer be legal. If you have questions about the legality and/or registration status for using pesticides, contact your county Extension office.



MSU is an Affirmative-Action Equal Opportunity Institution. Extension programs and materials are open to all without regard to race, color, national origin, sex, disability, age or religion. ■ Issued in furtherance of Extension work in agriculture and home economics, acts of May 8 and June 20, 1914, in cooperation with the U.S. Department of Agriculture. Gail L. Imig, director, Michigan State University Extension Service, E. Lansing, MI 48824. ■ This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by Extension or bias against those not mentioned. This bulletin becomes public property upon publication and may be reprinted verbatim as a separate or within another publication with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company. Produced by Outreach Communications and printed on recycled paper using vegetable-based inks.

