

MSU Extension Publication Archive

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Using Bt to Control Gypsy Moth

Michigan State University Extension Service

Leah S. Bauer, and Norman R. DuBois, USDA Forest Service, Deborah G. McCullough,
Departments of Forestry and Entomology, MSU

Issued August 1999

2 pages

The PDF file was provided courtesy of the Michigan State University Library

Scroll down to view the publication.

USING BT TO CONTROL GYPSY MOTH

By Leah S. Bauer, Deborah G. McCullough, and Norman R. Dubois¹

Reprinted August 1999

Extension Bulletin E-2421

Michigan State University Extension

Did you notice trees with partly or completely eaten leaves last summer? Did you see lots of big, hairy caterpillars feeding on the leaves? Even worse, were you plagued by the frass (feces) and the many caterpillars that fell from the trees? If so, the offender was probably the gypsy moth caterpillar. Other caterpillars feed on trees in Michigan, but the gypsy moth is the most common and annoying in many areas.

Why get rid of gypsy moths?

Gypsy moth caterpillars not only strip trees of their leaves but their frass and decaying bodies can cause a nuisance in picnic areas, recreation areas and backyards. In addition, people living where gypsy moth populations are high can experience skin or eye rashes or allergies due to caterpillar hairs.

To prevent severe defoliation from occurring this year, you may see airplanes or helicopters spraying insecticide in the spring. The insecticide used in Michigan is called Bt, also known as Thuricide, Foray, or Dipel. Bt is used around the world to control many kinds of caterpillars, including pests of agricultural crops. It has proven to be very safe and effective over the years. Bt has been used in the northeastern United States to control gypsy moth caterpillars since 1961.



What is Bt?

Bt is the nickname for *Bacillus thuringiensis* var. *kurstaki*, a bacterium found naturally on leaves and in the soil. In the early 1900s, scientists discovered that when caterpillars ate Bt, the insects stopped eating and died. Scientists took advantage of Bt's natural ability to kill caterpillars and used it to develop a new type of insecticide. Insecticides made with

microorganisms like Bt are called either microbial insecticides or biological insecticides.

How was Bt developed as a microbial insecticide?

Insecticides that use Bt as the active ingredient are available now because of a tremendous effort made during the last 80 years by government, university, and industry researchers throughout the world.

¹Leah S. Bauer
USDA Forest Service-NCFES
Pesticide Research Center
Michigan State University

Deborah G. McCullough
Departments of Entomology and Forestry
Michigan State University

Norman R. Dubois
USDA-Forest Service-NEFES
Hamden, Connecticut

How is Bt different from chemical insecticides?

Conventional chemical insecticides are manufactured poisons. We know that such a poison can kill most insects that consume it or come in contact with it. Since insects are biologically similar to animals, people will react to high dosages of these poisons in the same way. In addition, some chemical insecticides remain in the environment for a long time and can create problems for humans and other animals.

Microbial insecticides like Bt use a naturally occurring microorganism such as a bacterium, fungus, virus, or nematode as the active ingredient. Such a microorganism causes specific insects to get sick and eventually die. They often affect only one or a few species. Like manufactured poisons, microorganisms must be eaten or must invade the insect's skin or other natural opening. However, microbial insecticides kill only certain insects – they are harmless to people, other animals, and even most insects. In addition, microbial insecticides usually break down quickly and don't build up in the environment.

What does Bt do to gypsy moths?

Gypsy moth larvae pass through four different life stages: egg, caterpillar, cocoon and adult. Only one stage – the caterpillar – feeds on trees. Therefore, to be

effective, an insecticide must be able to kill caterpillars. The active ingredient of most microbial insecticides prepared from Bt is 1) a crystal protein made by Bt and 2) the spore or resting stage of Bt. When larvae eat leaves sprayed with Bt, the crystal dissolves and releases the protein that damages their stomachs. At this point, the caterpillars become sick and stop eating. During the next 2 to 3 days, bacteria enter the caterpillar's blood, eventually killing them.

Is Bt dangerous to people, pets, livestock, or honeybees?

One of the advantages of Bt is that only young caterpillars are affected. Bt is harmless to people, even at very high doses. Bt does not harm other mammals, birds or fish. Most beneficial insects, including honeybees and ladybugs, are not affected by Bt. Furthermore, Bt does not affect plants. It is considered so safe that it can be sprayed on food crops until the day they are harvested.

Does Bt contaminate the environment?

After Bt is sprayed, most of it is quickly destroyed by sunlight and other microorganisms. Larvae killed by Bt will drop to the soil and decay harmlessly. Bt does not multiply or accumulate in the environment.

Why is Bt sprayed so early in the season?

Young caterpillars are much more sensitive to Bt than older, larger caterpillars. The best time to spray is just after gypsy moth eggs have hatched and before larvae grow too big.

Who produces Bt?

Several private companies produce the Bt products that are used in Michigan. The large market for Bt has stimulated many new and existing companies to begin research on Bt and to develop improved Bt products.

Is it possible to buy Bt to spray caterpillars in my own garden?

Yes there are several commercially available biological insecticides that use Bt. Both wettable powders and liquid formulations are available from local garden shops. Be sure to follow directions on the label.

Are there other types of Bt for control of other garden or urban insect pests?

Yes, many types of Bt exist. The most common type of Bt affects caterpillars like the gypsy moth. Another important type is used to control mosquito and black fly larvae in places like ponds, streams, and gutters. A third type of Bt is used to control the larvae of the Colorado potato beetle and the cottonwood leaf beetle.



MSU is an Affirmative-Action Equal-Opportunity Institution. Michigan State University Extension programs and materials are open to all without regard to race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, marital status, or family status. • Issued in furtherance of Extension work in agriculture and home economics, acts of May 8 and June 20, 1914, in cooperation with the U.S. Department of Agriculture. Arlen Leholm, Extension Director, Michigan State University, E. Lansing, MI 48824. • This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by MSU Extension or bias against those not mentioned. This bulletin becomes public property upon publication and may be printed verbatim with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company.