

## **MSU Extension Publication Archive**

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Entomophaga maimaiga - A Natural Enemy of Gypsy Moth

Michigan State University Extension Service

Lyle Buss, M.S. Research Assistant, Dept. of Entomology, Dept. of Forestry; Dr.

Deborah McCullough, Assistant Professor, Dept. of Entomology, Dept. of Forestry; Dr.


David R. Smitley, Associate Professor, Dept. of Entomology

May 1996

2 pages

The PDF file was provided courtesy of the Michigan State University Library

**Scroll down to view the publication.**



# *Entomophaga maimaiga* - A Natural Enemy of Gypsy Moth

Michigan State University Extension

Lyle Buss  
M.S. Research Assistant  
Dept. of Entomology  
Dept. of Forestry  
Michigan State University

Dr. Deborah McCullough  
Assistant Professor  
Dept. of Entomology  
Dept. of Forestry  
Michigan State University

Dr. David R. Smitley  
Associate Professor  
Dept. of Entomology  
Michigan State University

The gypsy moth (*Lymantria dispar* L.) is an exotic pest of urban and forest trees. Gypsy moth caterpillars feed on the leaves of oaks, aspens, and many other hardwood and conifer trees. During gypsy moth outbreaks, trees may be completely stripped of leaves. Although gypsy moth caterpillars rarely kill trees by themselves, trees weakened by heavy defoliation may become more susceptible to drought, disease or other insect pests. In addition, the large hairy caterpillars annoy people living or recreating in outbreak areas.

Managing gypsy moth requires the integration of a variety of control tactics. Biological control, the use of natural enemies to control a pest, can be an important part of an integrated pest management program for gypsy moth. One biocontrol agent that has recently shown much promise is a fungal pathogen, *Entomophaga maimaiga*.

## Origin of *Entomophaga maimaiga*

*Entomophaga maimaiga* is a common disease in gypsy moth populations in its native country of Japan. The fungus was first released into the United States near Boston in 1910 as part of a program to introduce natural enemies of gypsy moth. Scientists could find no evidence that the fungus had become established and the project was abandoned a few years later. However, the fungus appeared unexpectedly in several northeastern states in 1989 and caused high mortality in many gypsy moth populations. Although scientists have several theories, the strange reappearance of the fungus is still a mystery.

*Entomophaga maimaiga* was first brought into Michigan in 1991 by scientists at Michigan State University and the USDA Forest Service. It was released in three sites in northern lower Michigan and monitored closely.

Additional introductions have since occurred and the fungus also is spreading naturally. *Entomophaga maimaiga* is now found in much of the northwestern Lower Peninsula and in some areas of the southern Lower Peninsula.

## Life history of *Entomophaga maimaiga*

*Entomophaga maimaiga* passes the winter as a tough, thick-walled "resting spore" in the soil and on tree bark. In May and June, resting spores germinate and produce sticky spores at the end of a stalk that grows just above the soil surface. Gypsy moth caterpillars come into contact with these spores in the spring as they search for suitable leaves to feed on. The fungus digests its way through the exoskeleton of the caterpillar and grows inside the body of the caterpillar. Infected caterpillars may die within one week.

When young caterpillars are affected early in the summer, the fungus will produce a second type of spore called conidia. These microscopic spores are spread by the wind and can infect other caterpillars. The cycle of conidia production and infection may occur four to nine times during the summer. When the fungus develops in large caterpillars, it produces the overwintering resting spores.

Weather plays an important role in determining how effective *Entomophaga maimaiga* will be. Like most fungi, its spores need moisture and high humidity to germinate. Frequent rainfall during May and June contributes to the start and spread of *Entomophaga maimaiga* through a gypsy moth population. Temperatures of 50 to 80 degrees F enhance fungal growth.




Figure 1. Gypsy moth larva killed by NPV hanging in an inverted "V" position.



Figure 2. Dead larva covered with conidia of *Entomophaga maimaiga*.



Figure 3. Larvae killed by *Entomophaga maimaiga* often remain attached to trees.

### Distinguishing *Entomophaga maimaiga* from NPV

Another disease is common in outbreak populations of gypsy moth. NPV (nuclearpolyhedrosis virus) is a virus disease that often causes gypsy moth outbreak populations to collapse. One important difference between the two diseases is that NPV is seldom prevalent until gypsy moth populations reach very high levels. In contrast, *Entomophaga maimaiga* may be found even when gypsy moth populations are low.

Caterpillars killed by NPV often remain attached to the stem or branches of trees. The bodies of the dead caterpillars are soft, filled with a brown liquid and disintegrate rapidly. Usually they hang limply in an upside-down "V" position (Fig. 1).

Caterpillars killed by *Entomophaga maimaiga* will also remain attached to tree stems or branches. However, the bodies tend to be stiff and straight, and the legs extend stiffly from the body. Some of the dead caterpillars may have tiny white conidia attached to the hairs on the body (Fig. 2). The cadavers may remain on the stem well into the winter (Fig. 3).

### The future of *Entomophaga maimaiga*

*Entomophaga maimaiga* may become an important biological control of gypsy moth in both low and high populations. Infections may be more common in years with rainy spring weather than in years with dry spring weather. Scientists have found that the fungus is established in a number of areas in Michigan. Laboratory and field studies have shown that *Entomophaga maimaiga* is host specific and poses little risk to other insect populations. It will not affect other animals or humans. Introductions and evaluation will continue. Although there is not likely to be any "silver bullet" for gypsy moth, *Entomophaga maimaiga* should improve our ability to manage this pest in Michigan.

MICHIGAN STATE  
UNIVERSITY  
EXTENSION

MSU is an Affirmative-Action Equal-Opportunity Institution. MSU Extension programs are open to all without regard to race, color, national origin, sex, disability, age or religion.

Issued in furtherance of Cooperative Extension work in agriculture and home economics, acts of May 8, and June 30, 1914, in cooperation with the U.S. Department of

Agriculture. Gail L. Imig, Director, Michigan State University Extension, E. Lansing, MI 48824.

This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by the MSU Extension or bias against those not mentioned. This bulletin becomes public property upon publication and may be reprinted verbatim as a separate or within another publication with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company.

New 5:96-3M-LJ-COL-Price \$.50. Single copy free to Michigan residents.

File 27.35 (Pests and Management)

Printed on recycled paper using vegetable-based inks.