


MSU Extension Publication Archive

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Weed Seedbank Dynamics
Michigan State University
Michigan State University Extension
Karen A. Renner, Department of Crop and Soil Sciences
Issued January 2000
4 pages

The PDF file was provided courtesy of the Michigan State University Library

Scroll down to view the publication.



Weed Seedbank Dynamics

Karen A. Renner, Professor
 Dept. of Crop and Soil Sciences
 Michigan State University

Agricultural soils contain thousands of weed seeds per square foot. The density of weed seeds in the weed seedbank is influenced by past farming practices and will vary from field to field (Table 1, Renner, 1999) and even between areas within fields. In intensively cropped fields in the north central corn belt, the weed seedbank ranged from 56 - 14,864 seeds per square foot (Forcella *et al.*, 1992).


Table 1

How many weed seeds are in the weed seedbank?	
Site	Number of seeds per square foot
W.K. Kellogg Biological Station — Long Term Ecological Research Site, Hickory Corners, Michigan	465 - 1,394
MSU Agronomy Farm, East Lansing, Michigan	1,394 - 2,787
U.S. Corn Belt	56 - 14,864

Composition of Weed Seedbanks


Seedbanks are made up of numerous weed species although only a few species will comprise 70 to 90 percent of the total seedbank. Common lambsquarters (*Chenopodium album*) is the dominant weed seed in many field soils in the north central region of the United States, including Michigan. Common lambsquarters dominated the weed seedbank in three of five cropping systems at the Long Term Ecological Research (LTER) site at the Kellogg Biological Station in Michigan (Figure 1).

Figure 1



This project supported by Project GREEN.

MICHIGAN STATE
 UNIVERSITY
 EXTENSION



Weed Seed Distribution in Soil

The location of seeds in the weed seedbank is influenced by the tillage system. More weed seeds will remain near the soil surface when tillage is reduced or no-till farming is practiced (Figure 2). These changes in the distribution of the weed seeds in the weed seedbank will influence weed emergence and the resulting weed population in farm fields.

Figure 2




Table 2

Typical Michigan weed seed production			
Weed	Number of seeds per plant	Weed density (per 33 feet of crop row)	Crop
Velvetleaf	400 - 1,500	90	corn
Giant foxtail	2,500	100	corn
Common lambsquarters	57,000	8	soybean

contains weed seeds on farm equipment or moving weed seeds to other fields when harvesting crops.

Sources of Weed Seed

Weed seeds can reach the soil and become part of the seedbank through several avenues. The main source of weed seed in the seedbank is from weeds that matured in the field and set seed. Annual weeds produce large numbers of seeds (Table 2, Renner, 1999). Weed seed can also enter


the seedbank by wind, water, animals, birds, and human activity. Some weed seeds (such as dandelion) are wind-dispersed. Weed seeds can reach a field site following flooding of drainage ditches or adjacent rivers. Wildlife and livestock can spread weed seed, either directly or by spreading of manure. Farming operations can add weed seed to the seedbank by moving soil that

Weed Seed Fate

A weed seed can have numerous fates once it is dispersed in a field (Figure 3, Renner, 1999). Some weed seeds will decay in the soil. Other seeds will not decay but will no longer have the ability to germinate (the seeds are not viable). Some weed seeds will germinate and die, while other weed seeds will germinate and emerge. Some weed seed will be predated by various predators including birds, rodents, crickets, carabid (ground) beetles, and ants. Seed predation occurs mainly on or near the soil surface. Many weed seeds will remain dormant in the soil and not germinate regardless of environmental conditions. However, dormancy is not permanent and seeds of many



Figure 3



seed longevity in soil since weed seeds usually remain viable longer if they are buried. Seed on or near the soil surface is exposed to predators and seed decay which reduces seed persistence.

Managing the Weed Seedbank

The best way to manage the weed seedbank is to not allow weeds to set seed in the field. Over a six-year period in Colorado, common lambsquarters and redroot pigweed seeds were reduced to 6 and 1 percent, respectively, of the original seedbank in a continuous corn rotation where herbicides were applied and the fields cultivated (Schweizer and Zimdahl, 1984). In a Nebraska study, the broadleaf and grass weed seed density in soil declined by 95 percent over a five-year period. However in the sixth year weeds were *not* controlled and the weed seedbank increased to within 90 percent of the original level at two of five locations (Burnside et al., 1986). These studies illustrate two important points in weed seedbank

weed species change from a state of dormancy to non-dormancy. This is called dormancy cycling (Figure 4, Renner, 1999). Seed dormancy is a survival mechanism and it is a major barrier to weed control in agroecosystems. Only a fraction of the weed seeds (less than 10 percent of most weed species) germinate each year. Therefore dormant seeds perpetuate the weed seedbank and weed populations in farm fields.

Weed Seed Persistence

Under agricultural conditions the average time that a weed seed will persist in soil and still be capable of germinating (remain viable) is less than five years. Some weed species will have more persistent seed than others. Velvetleaf (*Abutilon theophrasti*) and clovers (*Trifolium* sp.) have persistent seedbanks. Tillage also influences



Giant Foxtail



Redroot Pigweed





Figure 4



Adapted from Baskin & Baskin

management. First, there is a rapid decline in the weed seedbank when weeds are not allowed to set seed. Secondly, the few weed seeds remaining in the weed seedbank are capable of infesting the farm fields and returning the number of weed seeds in the seedbank to high levels. Therefore weeds must be managed every year to reduce the weed seedbank.

Other farming practices can influence the weed seedbank. Burying weed seed by tilling the soil increases the longevity of weed seeds in the seedbank. Leaving weed seeds on the soil surface exposes weed seeds to

predation which will reduce the number of weed seeds in the seedbank. Leaving weed seeds on or near the soil surface may increase the number of weed seeds that decay after being infected by fungi or other microorganisms. Livestock manure that is stored has fewer viable weed seeds compared to fresh manure. Cleaning tillage and harvest equipment can reduce the movement of weed seed from field to field.

Understanding weed seedbank dynamics is the first step in managing the weed seedbank. Reducing the number of weed

seeds in the weed seedbank will improve our management of weeds in agroecosystems.

References

Burnside, O.C., R. G. Wilson, G. A. Wicks, F. W. Roeth, and R. S. Moomaw. 1986. Weed seed decline and buildup under various corn management systems across Nebraska. *Agronomy Journal*. 78:451-454.

Forcella, F., R. G. Wilson, K. A. Renner, J. Dekker, R. G. Harvey, D. A. Alm, D. D. Buhler, and J. Cardina. 1992. Weed seedbanks of the U.S. corn belt: magnitude, variation, emergence, and application. *Weed Science*. 40:636-644.

Renner, K. A. 1999. Weed ecology and management. Pages 51-68 in *Michigan Field Crop Pest Ecology and Management*. M. Cavigelli, ed. Michigan State University Extension Bulletin E-2704.

Schweizer, E.E. and R. L. Zimdahl. 1984. Weed seed decline in irrigated soil after six years of continuous corn and herbicides. *Weed Science*. 32:76-83.

MSU is an affirmative-action equal-opportunity institution. Michigan State University Extension programs and materials are open to all without regard to race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, marital status, or family status. • Issued in furtherance of Extension work in agriculture and home economics, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Arlen Leholm, Extension director, Michigan State University, E. Lansing, MI 48824. • This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by MSU Extension or bias against those not mentioned. This bulletin becomes public property upon publication and may be printed verbatim with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company.

Produced by ANR Communications.
New - 1.00 - KMF - LP - 1.5M, Price 50¢, Single copy free to Michigan residents.