

MSU Extension Publication Archive

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Giant Hogweed
Michigan State University
Michigan State University Extension

Issued May 2005
4 pages

The PDF file was provided courtesy of the Michigan State University Library

Scroll down to view the publication.

Giant Hogweed

An attractive but dangerous federal noxious weed –
Have you seen this plant in Michigan?

Report giant hogweed in Michigan

Call MDA 1-800-292-3939
Select “3” from menu



Hogweed is hazardous

Giant hogweed is a majestic plant that can grow as tall as 20 feet. Although attractive, giant hogweed is a public health hazard because it can cause severe skin irritation in susceptible people. The plant exudes a clear, watery sap that causes photodermatitis, a severe skin reaction. Skin contact followed by exposure to sunlight may result in painful, burning blisters and red blotches that later develop into purplish or blackened scars. The reaction can happen within 24 to 48 hours after contact with sap, and scars may persist for several years. Contact with the eyes can lead to temporary or permanent blindness. Wash immediately with soap and water if skin exposure occurs. If

possible, keep the contacted area covered with clothing for several days to reduce light exposure.

Giant hogweed (*Heracleum mantegazzianum*) is a federal noxious weed, so it is unlawful to propagate, sell or transport this plant in the United States. The U.S. Department of Agriculture (USDA) has been surveying for this weed since 1998 and has identified several infestations in Michigan. The USDA cannot check all of Michigan, however, so we need your help in locating sites of giant hogweed. If you see this weed, please report its occurrence to the Michigan Department of Agriculture (MDA), the USDA, Michigan State University Extension or MSU Diagnostic Services.



It's a tall, majestic plant, but **DON'T TOUCH IT!**

Leg burn from giant hogweed.

Use common sense around giant hogweed

Don't touch or handle plants using your bare hands.

Don't allow children to play in giant hogweed. They may sometimes use the long, hollow stems for telescopes or peashooters.

Don't transplant or give away giant hogweed plants or seeds. These actions are illegal.

Do wash immediately with soap and water if giant hogweed sap gets on your skin.

Do report locations of giant hogweed plants by calling the Giant Hogweed Hotline at 1-800-292-3939.

How to recognize giant hogweed

Giant hogweed is a biennial or perennial herb capable of reaching 6 to 12 feet in height in Michigan. The best time to identify giant hogweed is during flowering. Plants sprout in early spring from seed or tuberous rootstocks. Plants can quickly form a solid canopy and displace native vegetation.

- **Stems** are 2 to 4 inches in diameter, hollow, stout and ridged with purple blotches and coarse, white hairs. Stems may approach 12 feet in height in Michigan.
- **Leaf petioles (leaf stems)** are hollow, purple blotched and sometimes nearly solid purple near the base. Coarse, white hairs are especially prominent circling the stem at the bases of the petioles.



Wearing gloves, a technician holds a giant hogweed stem.

- **Leaves** are very large -- up to 5 feet across. Lower leaves are compound with three large, deeply cut leaflets. Each leaflet has deep, irregular lobes and coarse, sharp teeth on the margins. Upper leaves are similar in shape, smaller, often not divided but simply deeply three-lobed. Hairs on the underside are stiff, stubby and approximately 0.25 mm long, and they may not be visible to the naked eye. The leaf underside looks smooth and scaly.



Left: Giant hogweed leaf – 3 feet long. Right: Flower.

- **Flowers** are white, clustered into a large, compound umbel with a flat bottom and gently rounded top. Umbels can be 2.5 feet wide. The plant flowers from June to August in Michigan.

Growth stages

1) Dead stems. After producing seeds in late summer, the plants die and leave stems standing into winter. At this point seeds have been dispersed to germinate the following spring or in future years.



2) Fruit. Mature fruit consist of two sections. Each section is oval in outline, 7 to 13 mm long, 6 to 10 mm wide, flattened and tan colored with usually four prominent dark lines (oil tubes).



3) Seedlings. Seeds germinate from early spring throughout the growing season. Seedlings form a vegetative rosette pattern of growth the first season.



4) Rosette of leaves. Leaf clusters sprout from overwintering roots each year for two to five years until the plant flowers.



5) Flowering stems (see picture at top right of page). Plants bolt and flower in midsummer after rosette plants accumulate enough energy reserves (this may take from two to five years).

Similar plants commonly mistaken for giant hogweed

Cow parsnip [*Heraclium maximum* (aka *H. lanatum*)] is a native plant very similar to giant hogweed. Cow parsnip is smaller, reaching a maximum height of 6 to 8 feet in Michigan. Unlike giant hogweed, the stem may be entirely green or have only a slight purplish cast. Stems are deeply ridged, often hairy below the nodes, often hairy

throughout and no greater than 2 inches in diameter. Hairs on cow parsnip are fine – soft and fuzzy – rather than coarse as on giant hogweed. Leaves are compound, up to 2.5 feet across, and contain three large, broad leaflets. Leaflets are deeply lobed with coarsely toothed margins. Stem leaves are smaller and their leaf stalks almost circle the

stem at the node. Although present on both leaf surfaces, the soft hairs are primarily confined to the lower leaf surface and have a velvety appearance. These lower leaf hairs are about 1 mm long and are often the best distinguishing characteristic from giant hogweed. Cow parsnip has white flower clusters similar to those of giant hogweed, but these

are flat-topped and smaller at 6 to 10 inches across. Cow parsnip generally flowers from early June through early July in Michigan, typically several weeks before giant hogweed. Mature fruits have two egg-shaped sections. Each section is 7 to 14 mm long and tan to pale tawny in color with four vertical purple lines (oil tubes).




Angelica (*Angelica atropurpurea*) can be easily distinguished from giant hogweed by several features. Stems are uniformly waxy green to purple, smooth, hollow and up to 2 inches in diameter. Mature leaves are double or triple compound with numerous leaflets and are capable of



reaching 2 feet in width. The globular, softball-sized clusters of greenish white flowers are up to 12 inches in



diameter. Angelica seldom reaches 8 feet in height and typically flowers from mid-May through mid-June in



Michigan. Mature fruit have two oblong-oval sections that are ribbed, 4 to 6.5 mm long and usually flattened.

Poison hemlock (*Conium maculatum*) is a multibranched, non-native biennial plant ranging from 4 to 9 feet tall. The waxy stem has purple blotches and the entire plant is smooth. The fernlike leaves are bright green and may appear glossy. Small, white flowers are arranged in numerous, small, flat-topped clusters on all the



branches. Leaves and flowers resemble those of wild carrot. Poison hemlock has a disagreeable "mousy" odor, and the entire plant



is poisonous if ingested. It generally flowers in late May through late June in Michigan. Mature fruit have two sections that are



and yellowish to gray-brown. Each fruit section is 2 to 3.5 mm in length with prominent, wavy, somewhat knotty lengthwise ribs.

Wild carrot (*Daucus carota*), also known as Queen Anne's lace, is a common biennial weed in Michigan that forms a basal rosette of leaves the first year followed by an erect, 3- to 4-foot flowering stalk the following year. Stems are erect, branched, hollow, rather thin and covered with short, coarse hairs. Leaves are very finely dissected, lacy to



fernlike and virtually identical to those of cultivated carrot. Lower leaves are considerably larger than the upper stem leaves. White to sometimes pinkish



flowers form a flat-topped cluster, often with a single, purple flower in the cluster center. Wild carrot flowers from July to September. Mature fruits have two egg-shaped



sections. Each section is flattened on one side, yellowish to grayish brown, 2 to 4 mm in length and ridged with barbed spines.

Wild parsnip (*Pastinaca sativa*) is a non-native species easily distinguished from the above species by its yellow flowers. Wild parsnip may cause photo-dermatitis similar to that caused by giant hogweed.



About giant hogweed – Where did it come from?

Giant hogweed, native to the Caucasus region of Eurasia, is a member of the carrot or parsley family (Apiaceae) that was introduced into Europe and North America in the early 1900s. Its massive, majestic size and imposing appearance made it desirable for arboretums and gardens. Because of its persistent and invasive habit, giant hogweed soon escaped cultivation and became established in rich, moist soils along roadside ditches and stream

banks, and in waste ground, tree lines and open wooded areas. In the United States, giant hogweed is known to occur in Connecticut, the District of Columbia, Maine, Maryland, Massachusetts, Michigan, New York, Ohio, Oregon, Pennsylvania, Washington and Wisconsin. It is also well established in southern Ontario. The plant's name comes from Hercules, of ancient mythological fame, and giant hogweed is aptly described as robust.

Prevention and control

If you suspect giant hogweed is on your property, please call MDA at 1-800-292-3939 and describe the plant so our staff can verify its identity. If the plant is giant hogweed, an MDA staff member will arrange to visit your property, assess the site and discuss management strategies with you. MDA, USDA and MSU staff members know which herbicide combinations are effective in controlling this noxious weed. Once a control program is initiated, MDA staff members will visit your property periodically to determine the success of the efforts and to check for any new seedlings that may have sprouted. Giant hogweed seeds may remain dormant in the soil for at least five years, so eradication requires a long-term commitment – something we are prepared to make.

Do not mow, cut or weed-whack to try to control the plant mechanically – its large perennial root system will soon send up new growth. Also, these tactics are risky

Michigan distribution of giant hogweed by county, July 2005

Branch	Gogebic	Kent	Saginaw
Berrien	Ingham	Manistee	
Calhoun	Jackson	Oakland	
Clinton	Kalamazoo	Ottawa	

A controlled patch of giant hogweed.



because they increase the opportunities for you to come in contact with the plant's sap.

Giant hogweed is spread naturally by seeds, which can be wind-blown and scattered several feet from the parental plant, or may be carried by water to invade new areas. However, people are usually responsible for spreading giant hogweed over long distances. Seeds or young plants from a friend's garden, planted in new locations, help spread this weed quickly over distances much greater than the plant would spread naturally. The dried fruit clusters are sometimes used in decorative arrangements, and when discarded outdoors, they can start a new infestation of giant hogweed.

We strongly encourage homeowners and landowners to report suspected giant hogweed to MDA or MSU Diagnostic Services. Proper identification is the first step toward eradicating this invasive, hazardous weed.

Michigan State University Extension

MSU Diagnostic Services

Michigan State University
101 Center for Integrated Plant Systems
East Lansing, MI 48824-1311
Phone: 517-355-4536; Fax: 517-432-0899
Web: www.pestid.msu.edu
www.ncpdn.org

Michigan Department of Agriculture

Pesticide, Plant Pest Management Division
P.O. Box 30017 Lansing, MI 48909
Phone: 517-373-1087
www.michigan.gov/mda

U.S. Department of Agriculture

Animal and Plant Health Inspection Service
Plant Protection and Quarantine
11200 Metro Airport Center Drive, Suite 140
Romulus, MI 48174
Phone: 734-942-9005

Report giant hogweed in Michigan

Call MDA 1-800-292-3939. Select "3" from menu.

MICHIGAN STATE
UNIVERSITY
EXTENSION



Diagnostic Services



Bulletin last revised May 2005 by Steven A. Gower and Robert J. Richardson, Michigan State University, in collaboration with the MSU IPM Program.

MSU is an affirmative-action equal-opportunity institution. MSU Extension programs and materials are open to all without regard to race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, marital status, or family status. Issued in furtherance of Extension work in agriculture and home economics, acts of May 8 and June 20, 1914, in cooperation with the U.S. Dept. of Agriculture. Thomas Coon, MSU Extension director, Michigan State University, E. Lansing, MI 48824.