

Asparagus Miner

Asparagus Miner

Ophiomyia simplex Loew
(Diptera: Agromyzidae)

Introduction

The asparagus miner was introduced into North America from Europe, and was first found along the northeast coast of the United States in 1869. This pest has two full generations per growing season in Michigan (Tuell & Hausbeck, 2008), and asparagus is its only known host (Chittenden, 1898). The asparagus miner has been recorded in every county in Michigan where there is commercial asparagus production, including Mason, Oceana, Van Buren, Allegan and Cass counties (see Figure 1). This factsheet provides general information on the biology of the asparagus miner and options for its management.

Identification

Adult asparagus miner flies are shiny black and about 3.5 mm to 5 mm long (see Figure 2). Their eggs are difficult to detect because they are small and translucent, later taking on the color of the asparagus stem. Females lay the eggs under the outer layer of stems. Larvae are creamy white and up to 5 mm long (Barnes, 1937). There are three instars (immature stages), the first is small and oval-shaped (see Figure 3A), the second is longer and tube-like (see Figure 3B) and the third is banana-shaped (see Figure 3C). Pupae are pale brown at first, later becoming dark brown, and are about 5 mm long (see Figure 3D).




Figure 1. Known distribution of the asparagus miner in Michigan.

Life History and Life Cycle

In Michigan, asparagus miners overwinter as pupae in mined asparagus stem debris (Lampert, Cress, & Haynes, 1984). Two complete generations take place in Michigan in each growing season (Lampert et al., 1984; Tuell & Hausbeck, 2008).

Adult flies normally emerge in late May to early June, and reach their first peak abundance around mid-June (see figure on page 3). Shortly after emerging, the adults mate and the females lay eggs near the base of asparagus stems, either right above or just below the soil surface.




Figure 2. Adult asparagus miners are shiny black and 3.5 mm to 5 mm in length.

Asparagus Miner




Figure 3. Life stages of the asparagus miner: A) First instar, B) Second instar, C) Third instar, D) Pupa.




Figure 4. A) Healthy asparagus stem, B) Stem damaged by asparagus miner activity, C) Miner pupating underneath the outer layer of an asparagus stem.




Figure 5. Number of parasitoids reared from 79 pupae that were collected from 5 commercial fields (n=630 stems) from July to September 2010.

Females are attracted to asparagus plants in fern, which means that young fields and volunteer asparagus are vulnerable targets for egg-laying early in the season. Adults feed at the base of unopened asparagus flowers and where asparagus beetle damage has occurred (Ferro & Gilbertson, 1982). Adults from the second generation are most abundant around mid-August, when the flies mate again, lay eggs, pupate and overwinter.

Damage and Importance

Adult flies do not cause damage, but the larvae tunnel in the asparagus stem and bore mines into the cortex around the base of the plant (see Figure 4). Mining usually does not affect nutrient transport, because only a small percentage of the plant's stem is damaged. However, mining is considered cosmetic injury and affects the harvestable yield and marketing of asparagus. In extreme situations, the larvae can girdle the stem.

Damage from asparagus miner has been associated with increased *Fusarium* crown and root rot and early decline of asparagus fields (Gilbertson, Manning, & Ferro, 1985; Tuell, 2003; Tuell & Hausbeck, 2008). *Fusarium* spores have been found abundantly on all stages of the asparagus miner (Tuell & Hausbeck, 2008). Pathogenic *Fusarium* fungi are likely able to enter the asparagus plant through the physical damage of miner oviposition holes and larval tunnels. Early decline from *Fusarium* can shorten the economic lifespan of asparagus fields by 5 to 8 years (Elmer, Johnson, & Mink, 1996).

Control Measures

Broad-spectrum insecticides are currently used to control the adult asparagus miner (Bishop, 2006). These products are not effective in reducing immature miner life stages that develop protected within the stems. Currently, control options are being tested for larval control.

Several biological control agents were described from the United Kingdom in the early 1900s, including larval parasitoids belonging to the families Pteromalidae, Eulophidae and Braconidae (Barnes, 1937).

Asparagus Miner




Illustration by Marlene Cameron.



Asparagus Miner

In a recent survey of commercial fields in Michigan, parasitoids belonging to these families have been found over the course of a growing season in Oceana County (see Figure 5).

Research at Michigan State University is currently being conducted to improve management of this pest through:

- Better monitoring tools.
- New integrated pest management tactics.
- Improving biological control.

Literature Cited

- Barnes, H. F. (1937). The asparagus miner (*Melanagromyza simplex* H. Loew) (Agromyzidae; Diptera). *Annals of Applied Biology* 24(3), 574–518.
- Bishop, B. A. (2006, December). *Asparagus miner research trial*. Presentation at the Great Lakes Fruit, Vegetable and Farm Market EXPO, Grand Rapids, MI. USA, pp 2. Retrieved from www.glexpo.com/abstracts/2006abstracts/Asparagus2006.pdf
- Chittenden, F. H. (1898). Insects that affect asparagus. In L. O. Howard (Ed.), *Some miscellaneous results of the work of the Division of Entomology* (Bulletin 10, pp. 54–62). Washington, DC: U.S. Department of Agriculture, Division of Entomology. Retrieved from http://naldr.nal.usda.gov/NALWeb/Agricola_Link.asp?Accession=CAT10415833
- Elmer, W. H., Johnson, D. A., & Mink, G. I. (1996). Epidemiology and management of the diseases causal to asparagus decline. *Plant Diseases*, 80(2), 117–125.
- Ferro, D. N., & Gilbertson, R. L. (1982). Bionomics and population dynamics of the asparagus miner, *Ophiomyia simplex* (Loew), in Western Massachusetts. *Environmental Entomology*, 11, 639–644.
- Gilbertson, R. L., Manning, W. J., & Ferro, D. N. (1985). Association of the asparagus miner with stem rot caused in asparagus by *Fusarium* species. *Phytopathology*, 75(11), 1188–1191.
- Lampert, E. P., Cress, D. C., & Haynes, D. L. (1984). Temporal and spatial changes in abundance of the asparagus miner, *Ophiomyia simplex* (Loew) (Diptera: Agromyzidae), in Michigan. *Environmental Entomology*, 13(3), 733–736.
- Tuell, J. K. (2003). *Fusarium and the asparagus miner (Ophiomyia simplex) in Michigan* (Master's thesis). East Lansing: Michigan State University, Department of Plant Pathology.
- Tuell, J. K., & Hausbeck, M. K. (2008). Characterization of the *Ophiomyia simplex* (Diptera: Agromyzidae) activity in commercial asparagus fields and its association with *Fusarium* crown and root rot. *Proceedings of the 11th International Symposium on Asparagus*, 776, 203–210.

Authors

Zsofia Szendrei, Department of Entomology, Michigan State University

William R. Morrison III, Department of Entomology, Michigan State University



This fact sheet was produced with support from Project GREEN, the C. S. Mott Foundation and MSU AgBioResearch. For more information, visit the Michigan State University Vegetable Entomology Szendrei Lab website online at vegetable.ent.msu.edu/.

MICHIGAN STATE UNIVERSITY | Extension MSU is an affirmative-action, equal-opportunity employer. Michigan State University Extension programs and materials are open to all without regard to race, color, national origin, gender, gender identity, religion, age, height, weight, disability, political beliefs, sexual orientation, marital status, family status or veteran status. Issued in furtherance of MSU Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Thomas G. Coon, Director, MSU Extension, East Lansing, MI 48824. This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by MSU Extension or bias against those not mentioned.