



Troublesome turf weeds in South Australia

Dr Terry Mabbutt travels to the other side of the world in his search for exciting new turf weeds.

First visit to a new country and climate on the other side of the world always provokes excitement at the prospect of hitherto unseen plants, even if they are troublesome turf weeds.

Like many plant scientists I started out as a botanist trained to appreciate plants for what they are in their natural environment, but graduated out of necessity into applied disciplines. These included managing plants growing in the wrong places such as amenity grass and fine professional turf of golfing green quality.

That apart some plant species have become so well adapted to turf that for all intents and purposes they are in their 'natural' environment. *Bellis perennis* (common daisy) in the United Kingdom immediately springs to mind. Moreover, you can appreciate such plants for their

ability to colonise turf which is not the most weed friendly of environments. With the UK behind me and still carrying some botanist's 'baggage' I arrived in the Adelaide Hills of South Australia intrigued about what weeds I might find growing in turf.

I was not to be disappointed. Turf in this relatively wet and mild southernmost area of South Australia up to 700 metres above sea level is forced to contend with many different weeds on par in range, number and frequency to the UK. This is not surprising since apart from three to four hot dry summer months (December to March) turf grass and weeds grow relatively unrestrained.

Distance and climate apart the same attributes which make for successful weeds in UK turf apply in South Australia, and in some cases to the very same species. Turf

weeds in South Australia include a high proportion of non-native species arriving as contaminants or escaping from gardens and subsequently adapting to turf. Some which originated in cool temperate climates like the UK appear to have discarded their normal perennial habit to become annuals, thereby allowing survival as seed through the long hot and dry South Australian summer.

Rosette and matt-forming weeds with creeping stems (stolons) or tap roots and fine-leaved plants (grasses and non-grasses) are clearly at the same advantage as in the UK. However, there are visibly but more Australian turf weeds with underground stems as bulbs and corms. They grow in winter, flower in spring and go dormant in summer. As such they are clearly well adapted to exploit good growing conditions in winter and spring



Cape dandelion is a classic rosette turf weed



Creeping oxalis is a serious weed of close cut



Flat weed (cat's ear) appears to be a bigger problem



Lamb's tongue (ribwort plantain) and trifoliate burr medic



One leaved Cape Tulip, pretty as a picture but a weed nevertheless



The leaves of onion grass blend in well but its pink

and subsequently survive adverse summer conditions. Moreover bulbs and corms are an ideal mode of reproduction and spread. Many like sparaxis/harlequin flower (*Sparaxis* sp) and one-leaved Cape tulip (*Moraea flaccida*) [all garden escapes] produce stunning spring flowers highly prized in UK horticultural and garden industries, but for South Australian greenkeepers and groundsman they are simply weeds.

The following account describes some of the most frequently seen weeds of turf in the Adelaide Hills area of South Australia some 20km south east of the capital Adelaide.

All were observed and photographed in amenity turf. Australia has a vibrant sports turf industry with a sophisticated arsenal of herbicides, its 'guns' apparently not yet spiked to the degree experienced in the UK courtesy EU (European Union) legislation. Weeds mentioned here can establish in amenity grass, wide area sports turf including fairways and poorly managed professional turf. However, like the UK only a minority of turf weeds are likely to be frequently seen in close-cut, well-managed professional turf.

Rosette forming weeds

Cape dandelion, Cape weed or Cape marigold (*Arctotheca calendula*) is a classic rosette-forming turf weed of the family Asteraceae (Compositae) with an extremely rapid growth rate. Deeply lobed basal leaves lay flat on the ground up to 25cm in length when fully grown, thus translating into massive rosette diameters of 50 cm. And all the more impressive because Cape dandelion unlike other related turfweeds (eg. English dandelion) is an annual herb in South Australia and clearly unable to survive the summer drought unaided.

Like many others is an alien species in this instance a native of the South African Cape with a similar Mediterranean-type climate.

Cape dandelion dies out in the summer heat and drought after shedding its high seed load. Seeds germinate in autumn and plants grow rapidly throughout the mild wet winter months. Key to survival and spread is its prolific spring and early summer flowering when poorly managed turf suddenly becomes a carpet of large bright brash daisies with light yellow ray (outer) florets and black/purple disc (inner) florets.

Seeds are covered with pale brown wool which clearly helps dispersal

by wind, animals, on clothing and most significantly by grass cutting machinery. Cape dandelion possesses a tap root but apparently not robust enough to survive the summer months.

Other 'big' rosette-forming turf weeds of the Asteraceae need no introduction to UK groundsman because Australia was 'cursed' with early introduction, by seed or through contaminated soil by *Taraxacum officinale* (British dandelion), *Hypochoeris radicata* (common cat's ear) and *H. glabra* (smooth cat's ear).

In the UK dandelion is more widespread and frequent than cat's ear but the opposite appears to be the case in South Australia where they do not appear to be as robust or free flowering as those in the UK, although the flowering period is much longer and may extend over the entire year. As in the UK *Taraxacum officinale* is a perennial with robust tap root and apparently

ful given new plants that can grow from root fragments.

A similar relationship exists for two plantain turf weeds, ribwort plantain (*Plantago lanceolata*) called lamb's tongue in South Australia and common plantain (*P. major*) called broad leaf plantain. In spread, frequency and size ribwort plantain outstrips broad leaf plantain, which is opposite to the situation in the UK.

Ribwort plantain covers turf with multitudes of rosette-forming plants which block sunlight and smother grass plants beneath, it's well developed tap-root facilitates year round survival in South Australia.

Ability to maintain the same perennial 'lifestyle' exhibited in its native cool temperate (UK) environment appears to be secret of its success as a turf weed in South Australia, putting the plant at relative advantage over similar rosette weeds including other plantains.

In the UK dandelion is more widespread and frequent than cat's ear but the opposite appears to be the case in South Australia where they do not appear to be as robust or free flowering as those in the UK.

able to withstand the summer heat and drought.

Cat's ear is called 'flat weed' in Australia due to its large flat basal rosettes of leaves, and appears more widespread, frequent and larger than in its native UK turf environment. Plants spread at speed smothering and killing turf grass underneath. Its broad basal leaves form large flat rosettes that merge to cover sizable areas to leave 'dead spots' when the plants die. *H. glabra* has maintained its annual habit.

The more common *H. radicata* is predominantly perennial as in the UK, but in drier areas of South Australia adopts an 'annual habit', expiring in summer after seed set and dispersal.

Cat's ears are therefore common even in the drier areas where the tap root is unable to survive the summer drought. *H. radicata* and *H. glabra* are normally difficult to distinguish and widespread hybridization between the species makes this virtually impossible.

Limited infestations are controlled by severing tap roots 2-4 cm below soil level using dedicated hand-tools. Same procedure can be used for dandelions using similar but deeper-penetrating tools, but as in UK is generally less success-

Stems above and below ground

Creeping Oxalis (*Oxalis corniculata*) is perennial weed spreading quickly in all types of turf and particularly persistent in low cut grass due to flat stems (stolons) that creep across the soil invariably escaping the mower's blades. Trifoliate leaves of three heart shaped leaflets together with the running stem habit makes creeping oxalis look just like a clover, but it comes from a completely different plant family called the Oxalidaceae (Wood Sorrel family). The plant is additionally called yellow wood sorrel because of its yellow flowers which though small are conspicuous in close-cut turf. Seed capsules shaped like miniature cucumbers throw ripe seeds several feet in all directions further aiding dispersal and spread.

O. corniculata is a locally abundant weed in the UK called 'Sleeping Beauty' found mainly near gardens from where it migrates into turf. In Australia it is completely cosmopolitan turf weed with many sub species making its exact alien origin unclear.

Other Oxalis species found in South Australian turf include *O. purpurea* (purple or large flowered wood sorrel) native to South Africa.



Plenty of *Poa* (winter grass) and a single 'shining'



Purple flowered wood sorrel persists and spreads by corms



Soursob is one of the most difficult weeds to control



Three cornered garlic grows literally everywhere

Its stunning carmine-coloured flowers open in the sun and hardly make for a classic turf weed but this garden escape is still a plant in the wrong place at the wrong time and as such a weed of turf. It lacks the stolons of creeping oxalis but has underground stems (corms) presenting their own problems for weed persistence. Limited infestation can be removed by hand-digging because the corms are in a shallow situation just beneath the surface of the ground.

Underground stems (bulbs and corms) are clearly to the advantage of any turf weed in South Australia with three to four months of extreme heat and drought. Two bulb/corm bearing species, both non-native garden-escapes and now invasive weeds, grow literally everywhere

including woods, waste-land, gullies in gardens and in turf. Three cornered garlic (*Allium triquetum*) after its three-sided flower stalks and native in the Mediterranean and Soursob (*Oxalis pes-caprae*) with striking yellow flowers and native to South Africa, can quickly invade and take over whole areas of amenity turf. Soursob, so called because of the sour acidic taste of its leaves and flowers, has a long underground stem bearing numerous bulbs making it one of the most difficult weeds to control. Soursob thrives in infertile soil and is suppressed by sound fertiliser programmes.

Few areas of turf grass outside the tropics escape white clover (*Trifolium repens*) and South Australia is no exception. This classic

matt-forming leguminous weed is just as widespread and problematic as it is in its native UK environment. White clover adopts the same perennial habit despite shallow rooting. Spread is by stolons (runners) which root at the nodes of the ground hugging stems where they come into contact with the soil. White clover's ability to adapt to most soil types together with a relatively long winter flowering period (May to September) clearly helps ensure its success as a turf weed in South Australia.

There are other closely related matt-forming leguminous weeds such as Burr Medic (*Medicago denticulate*), a larger and more robust looking version of UK turf's black medick (*Medicago lupulina*). Like white clover this southern

European native has small trifoliate leaves, the middle top leaflet on a longer stalk than the other two. Flowers are small yellow-orange and pea-like in solitary clusters on thin axillary stems.

Burr medic lacks the stolons of white clover but compensates with an efficient annual habit including prickly seed pods possessing hooked burrs which attach themselves to virtually anything they touch.

Burr medic is becoming more problematic with high infestations slowing down mowing and increasing the volume of clippings. Successful management requires effective control before plants come into flower during spring.

Grass plant look-alikes and the real thing

Plants with grass-like growth habits and leaves have a real advantage as turf weeds, not least because they are difficult to detect and therefore manage.

One such weed is onion grass or Guildford grass (*Romulea rosea*), another South African native and so called because its leaves look like those on onion or chive seedlings. This monocotyledonous species with bulbs (corms) is from the plant family Iridaceae, and therefore not

too distant from the Allicaceae (onion family). Onion grass is a small erect perennial herb bearing extremely fine linear leaves with thickened edges and a prominent midrib. You would not know it was there until spring when turf starts to sparkle with numerous tiny 'star-shaped' (6 petals) pink or rosy purple flowers.

South Australian turf facing hot dry summers and wet but essentially frost free winters offers the ideal 'meeting point' for grasses of both temperate and tropical origins. As such it has become overburdened with alien grass species from almost every part of the world, with rough grass weeds in turf generally divided 'winter' grass or 'summer' grass depending on when they grow and reproduce.

Main 'winter grass' requires no introduction. *Poa annua* (called 'winter grass in Australia) crops up everywhere in South Australia where it behaves as a winter annual as opposed to a summer annual in the UK. It flowers in winter and dies out in summer, although some perennial sub species are reported on golfing greens.

As in the UK controversy continues over its value as a constituent of golf greens. Winter grass germinates in the autumn and grows quickly

and flowers profusely throughout winter, finally seeding in spring when heads can cause an uneven playing surface on the fine turf such as golf greens. Perennial types of winter grass can become established in some fine turf situations. Winter grass continues to thwart those greenkeepers striving for 100% bent-grass playing surfaces. It remains widely present on greens (average golf green throughout Australia has been variously estimated to comprise 40-80% *Poa annua*) and is common on fairways.

'Summer grass' infestation is mainly by *Paspalum dilatatum* a tufted perennial tropical grass originating in South America. It is nominally frost sensitive but since South Australian winter temperatures rarely fall below freezing it can survive winter in a dormant state.

Moreover it is relatively drought tolerant and a combination of watering and high nitrogen during summer sees growth and infestation take off. Seed is sticky and spread easily on shoes clothing and grass cutting machinery. Strict hygiene with machinery and footwear scrupulously cleaned after working in *Paspalum*-infested areas of turf is essential to minimise seed dispersal and weed spread.



This article comes to you courtesy of the BIGGA Learning and Development Fund.
Thankyou to all our key sponsors

Continue to learn

at Harrogate Week 2010
Sunday 17 to Thursday 21 January 2010

The Flagship Education Event for the Sports and Fine Turf Industry

www.harrogateweek.org.uk

Visit our Harrogate Week page on...

facebook

Venue
Harrogate International Centre
North Yorkshire
England



