

# TORONTO C-15 BENT SOD & STOLONS

"PURE TO STRAIN"

Mowed at putting green height with  
"Greens King"

For Tees and Collars — FYLKING SOD  
WINDSOR SOD

For Fairways — BLUEGRASS BLEND SOD

## PENNCROSS BENT SOD



H & E Sod Nursery, Inc.  
4301 W. Flossmoor Rd.  
Tinley Park, Illinois  
312 798-2210

## GOLFERS PLAY

... while we install your  
irrigation system

one fairway at a time!

Free literature and list of installations on request.

**K**irchdorfer  
Irrigation, Inc

P. O. Box 4124 • Louisville, Ky. 40204  
Phone 585-4305 • Area Code 502

**C. E. STEWART**  
and Associates  
Civil Engineer

Irrigation, Drainage, Water Supply, Pumping Plants  
Design — Reports — Consultation — Supervision  
18357 Homewood Ave. Homewood, Illinois

### EVERSPRAY CO.

• UNDERGROUND LAWN SPRINKLING SYSTEMS  
**IRRIGATION EQUIPMENT  
AND SUPPLIES**

GEORGE WELLEK

9250 GOLF ROAD  
DES PLAINES, ILLINOIS  
296-5555

### COOPERATIVE EXTENSION SERVICE

University of Illinois

#### WILL BIRCH TREES GO THE WAY OF THE ELM?

Birch trees are some of the most beautiful plants available for gardeners in Chicagoland.

Sadly, these magnificent plants are threatened by an insect pest which can kill the tree outright. This pest is the bronze birch borer. In some localities nearly all white or paper birches even large old trees, have been destroyed by this pest.

Infestation is first noticed when tips of upper branches turn brown. Large branches progressively die back, until the entire tree succumbs. Infested branches often appear swollen and brown, with ridges around smaller ones.

An examination of the inner bark of infested branches reveals numerous burrows, each four to five feet long, crossing and recrossing, completely cutting off circulation of sap. The burrows are tightly packed with sawdust and each contains a slender white grub about 3/4 of an inch long.

The larvae pupate in spring and emerge in June as greenish bronze beetles which cut their way out of the branches, leaving small, slightly oval holes.

Females deposit eggs in cracks in the bark. Hatching takes place in about ten days and the tiny borers work their way into the tree to repeat the cycle, further injuring the tree.

Control of this pest is difficult. Hatching larvae are the only stage exposed to chemical control. The recommended material is Cygon (dimethoate). It should be applied to trunk and branches the first of June so that the larvae encounter it as they try to burrow into the tree.

Trees should be fertilized and watered to maintain active growth. Some overwintering larvae can be destroyed by pruning out and burning dead branches in early spring.

The most promising control appears to be a small, parasitic wasp which is capable of boring through the bark with its ovipositor to lay an egg on the borer larva. Research is now in progress to find ways of rearing and establishing this beneficial predator.

When calling Mr. Jones  
I find it most appalling  
To have his secretary ask,  
"May I tell him who is calling?"  
One of these days  
When I find it all too taxing,  
I'll come right back to her and say,  
"May I ask you, WHO IS ASKING?"

A dollar won't do as much for people as it used to, but then, people won't do as much for a dollar as they used to.

Bea von Boeselager