

We dug into the tall fescue control problem. So now you don't have to.

Introducing LESCO TFC™ Herbicide. The new, easy way to remove unwanted clumps of tall fescue without digging or spottreating with a non-selective herbicide. LESCO TFC is the first product to provide se lective spot control of tall fescue in Kentucky bluegrass, bentgrass, fine fescue, bahiagrass and bermudagrass, while allowing desirable grass to fill in. This water dispersible granule offers convenience and accurac Order today! Nationwide • (800) 321-532 In Ohio • (800) 686-7413 Always read and follow label instructions before using any chemical product. LESCO, Inc. 20005 Lake Road, Rocky River, Ohio 44116 (216) 333-9250

Sulfur Use on Greens Dr. Norm Hummel Cornell University

Soil reaction, or pH, is an important consideration in managing fine turfgrass areas. The pH is a measure of the hydrogen ion concentration in the soil solution. While most turfgrass species tolerate a wide range of soil pHs, optimum growing conditions exist in a pH range of 6.0 to 7.0. At excessively low pHs, the solubility of aluminum and manganese increase to a point where these elements can reach toxic levels in the soil. Excessive alkalinity, or very high pH, can result in deficiencies of several micronutrients.

Throughout much of New York State, calcareous sands are used to topdress goif course greens. Managing the pH on greens topdressed with these sands has been a major concern of superintendents. Applications of elemental sulfur are frequently used on an annual or biannual basis to reduce the pH of greens to a optimum range. Sulfur is an effective acidifying amendment thanks to the action of *Thiobacillus* bacteria that live in the soil. The bacteria obtain their energy from sulfur by oxidizing it into the sulfate form. A by-product of this oxidation process are hydrogen ions that reduce the soil pH.

Recently, isolated reports of damage from sulfur applications have surfaced. The symptoms have varied from what looks like an incurable infestation of dollarspot to a general discoloration. Damage has resulted from granular, flowable, and wettable powder forms of sulfur. The dollarspot-type damage caused by granular sulfur may not occur until several months after application.

While sulfur still remains a recommended acidifying material perhaps some caution should be exercised when using it. The following are suggested guidelines for using sulfur on bentgrass greens:

1. Do not use sulfur unless soil pH is above 7.5. Believe it or not, you can grow beautiful bentgrass greens on slightly alkaline soils. The use of micronutrients may be necessary as the pH rises much above 7.0. As an alternative to sulfur, acidifying nitrogen fertilizers such as ammonium sulfate may be used.

2. Apply no more than 2 pounds of sulfur per 1,000 square feet per application.

3. Only apply sulfur when temperatures are expected to be cool, preferably spring and fall.

4. Use a sulfur product the way it was intended to be used. For example, a wettable powder should be sprayed, not broadcast and watered in. **Read the label.**

5. A wettable powder or flowable sulfur material can be applied more uniformly than granular sulfur.

6. If damage occurs, do not aerify. Aerification will increase microbial activity, including the oxidation of sulfur by *Thiobacillus* bacteria. Aerification may actually increase damage.

7. Once damage occurs, there is nothing known to counteract the symptoms. If you feel compelled to do something a light application (2 lb./1,000 sq. ft.) of hydrated lime may or may not alleviate the problem. Fortunately, damage is not permanent and recover should be rapid.

Credit: Our Collaborator 8/89