(On the Waterfront cont'd.)

of PVC fittings removed from installations and tested, it appears that the ... observations may not hold true for fittings. At least the number of cycles to failure may be considerably less, due to stress concentrations at points of direction change in the fittings."

NEXT MONTH: BURST FAILURE AND LONG TERM PRESSURE FAILURE

This month's article is pretty vague. Next month's will be more understandable. I'll bet most of you didn't think I could write something this exciting. Hope this doesn't take as long to type as "War and Peace".

Management of Fungicide Resistance


by Patricia L. Sanders Plant Pathology Department Penn State University

Fungicides can be divided into two groups according to where they act to protect plants. CONTACT or PROTECTANT FUNGICIDES are those that stay on plant surfaces and provide a barrier against the fungi that cause disease. ERADICANT or SYSTEMIC FUNGICIDES are absorbed by plants, and thus can work to protect plants from within, in the same way that antibiotics act to eradicate "germs" inside human bodies. Most systemic fungicides also have protectant properties in that they can provide barriers to fungi on plant surfaces. Systemics have the advantage of long residual action, protection of plant crowns and roots, movement within plants to protect newly-formed tissues, eradication of fungi already inside plants, and protection from washoff and weathering.

The chief disadvantage of systemic fungicides has been the problem of resistance to these fungicides in many important turf pathogens. Resistance in fungi to systemic fungicides occurs because these fungicides generally poison fungi at only a single location in their growth and development cycles. It is, therefore, relatively likely that some individuals will be present in populations of disease-causing fungi that are able to circumvent or short-circuit the poisoned site. These individuals will be able to grow and increase in the presence of the fungicide. With repeated, continuous application of the same systemic fungicide, the naturally-resistant individuals in a fungal population will multiply until the population is composed primarily of fungicideresistant individuals, and disease control fails. This has happened in countries all over the world where systemic fungicides have been used. In the U.S.A., most of the disease control failures from resistance to systemic fungicides have occurred on turfgrass. There are published reports of resistance control failures of Tersan 1991 on dollar spot, Subdue on Pythium blight, and Chipso 26019 on dollar spot and pink snow mold.

Identification and development of new fungicides is costly and time-consuming. Therefore, we must learn to use systemics in ways that will prolong their useful lives. In order to prevent or delay fungicide resistance in populations of disease-causing The Ex Stands For Excellence

Why the EX STANDS FOR EXCELLENCE in professional turf care.



Par Ex is the only professional fertilizer line containing IBDU, undisputedly the finest slow release Nitrogen source in the world.

IBDU gives you the highest level of WIN, or Water Insoluble Nitrogen.

IBDU delivers all of its Nitrogen in a single growing season.

The Nitrogren release rate of IBDU is generally affected by only two factors: soil moisture and granule size.

Slower release makes Par Ex products perfect for your Fall Fertilization program.

Fewer applications are required, labor costs are cut.

The salt index of IBDU is the lowest of all commonly used fertilizer materials.

IBDU fights nitrate pollution because very little Nitrogen is lost through leaching or volatilization.

Choose from over 60 different product formulations.

Territory Manager Thomas E. Skinner Keokuk, IA 319/524-8912



(cont'd. page 22)

(Fungicide Resistance cont'd.)

fungi, it has been suggested that systemic fungicides should be alternated or used in mixtures. However, there are very few published research studies on which recommendations for preventing or delaying fungicide resistance can be based. Alterations will be effective in cases where the resistant individuals in the population are not as competitive as sensitive individuals. Thus, the population will fluctuate; the resistant component increasing when the resistance-prone fungicide is applied, and the more vigorous sensitive component increasing when the fungicide selection pressure is not present and the alternate fungicide is being used. Unfortunately, many times the resistant individuals in fungal populations are just as competitive and vigorous as the sensitive ones. In such cases, an alternating program will result in a steady increase in proportion of resistant members, until finally the population is predominantly resistant — a condition we are trying to avoid. In populations of equally-fit resistant and sensitive components, mixtures have been found to be effective in keeping resistant proportions stable in experimental populations.

Assuming that fungicide mixtures are able to keep resistance levels stable in fungus populations, they must be effective in controlling disease. Obviously, we cannot use full rates of fungicides in mixtures, because to do that would increase financial and environmental costs. We need to be sure that reduced rates of fungicides in mixtures will give satisfactory field control of diseases. Field and greenhouse studies have shown that reduced-rate mixtures can give disease control equal to, and sometimes greater than, the additive control of the individual mixture partners alone at the reduced rate. Although much more research is needed, it appears that reduced-rate mixtures can give acceptable field disease control, as well as delaying problems with resistance.

There are several important things to consider when selecting fungicides for use in alternations or reduced-rate mixtures. First, only fungicides with different ways of controlling the target fungus can be used in alternations or mixtures to delay or prevent control failures resulting from fungicide resistance in fungal populations.

The three systemic fungicides registered for Pythium blight control (Banol, Aliette, and Subdue) have different modes of action, and therefore, can be used in alternations or two-component, half-rate mixtures for resistance management and disease control. Three-component, third-rate mixtures of Banol/Aliette/Subdue may also be effective for these purposes, but research to test this is not completed.

The broad-spectrum systemic fungicides that control other turf diseases fall into three groups according to their mode of action: the benzimidazoles (Tersan 1991, Fungo 50, CL 3336), the dicarboxymides (Chipco 26019, Vorlan), and the sterol inhibitors (Banner, Bayleton, Rubigan). Any fungus that is resistant to one of the benzimidazole fungicides will be resistant to them all. The same is true within the dicarboxymide and sterol-inhibitor groups of fungicides. Therefore, for resistance management, broad-spectrum systemic fungicides must be mixed or alternated BETWEEN but not WITHIN groups. Systemic fungicides may also be mixed or alternated with any contact fungicide that will give the disease control desired.

(cont'd. page 23)

PRECISION BLENDED TOP DRESSING

Custom blended to your specifications by our modern equipment.

Buy when you need — Eliminate costly storage

We sell an air-dried, uniform and free flowing top dressing.

ASK THE MAN WHO HAS USED IT.

HENRY FRENZER

Area Code 708 658-5303 620 Webster St. Algonquin, IL 60102

Specializing in Golf Course Renovations

- Tees Greens Traps Fairways
- · Hydroseeding · Plant Installation ·

don fiore

landscape contractors & architects golf course contractors

28846 Nagel Court Lake Bluff, Illinois 60044 708-234-0020 Fax 708-234-0922

Certified Member: American Society of Landscape Architects Midwest Association of Golf Course Superintendents In addition to mode of action differences, the length of disease control provided by mixture components must be matched to avoid resistance selection. If a short-residual fungicide is included in a mixture for delaying resistance, an interspray of the short-residual chemical probably will be necessary.

If they are available, it is probably much better to use systemic fungicides in mixtures for resistance management. The reason is that the turfgrass plant itself can "unmix" mixtures of contact and systemic fungicides. If you apply a contact/systemic mixture, the mixture will be present on plant surfaces, but the systemic fungicide will be present alone inside the plant. As an example, in the case of a Subdue/Fore mixture. Subdue alone will be acting against **Pythium** that already has invaded the plant. For this reason mixtures of systemics are safer for resistance delay than contact/systemic mixtures.

The management of fungicide resistance in populations of disease-causing fungi is an area where much more research is needed. Additive, synergistic, or antagonistic effects may be possible with particular fungicide mixtures. It is, therefore, important that alternations and mixtures of various fungicides be tested, both for disease control and for resistance delay, in as many use settings and turfgrass/pathogen systems as possible.

Although there is much more we need to know about how we can best use systemic fungicides to avoid disease control failures from fungicide resistance in fungal populations, one thing is clear. We cannot safely use any systemic fungicide repeatedly and exclusively for disease control. Sensible and prudent use of systemic fungicides dictates diversity in chemicals used. Turf managers should be very skeptical of recommendations suggesting that any systemic fungicide can be used alone and continually without risk of resistance problems.

Editor's Note: Brian Bossert sent me this article as a followup on work Peter Hahn was doing at Ridgemoor C.C.

"The Good News Is ..."

by Vicki Lynn Sims, C.P.A.

THE GOOD NEWS IS ...

There are many income items which we receive tax-free. These qualify, because of their natures, under the Internal Revenue Code as "income exclusions".

JOB RELATED EXCLUSIONS

These payments benefit the employee, truly serve as compensation, provide a tax deduction for the employer, and are not taxed to the employee. In the case of the non-profit employer (as are many golf clubs), there is no need for the tas deductibility, but any exclusion available to the employee is still important.

MEDICAL COVERAGE

The most common area of income exclusion, this coverage represents TWO nontaxable benefits. We do not pay tax on the value of th premium paid on our behalf, and we do not pay tax on the value of the care provided.

DEPENDENT CARE ASSISTANCE

This is one of the best, but least used, employee benefits available. The employee may exclude amounts paid by the employer for furnishing dependent care on behalf of the employee. The employer directly pays the day care, nursing care, etc., the employer deducts the payment as an employee benefit and the employee is not taxed on the payments. The payments are excludable, up to the amount of earned income (for a married employee, the earned income of the lower earning spouse), subject to a \$5000 a year limit (\$2500 for married filing separately). Special rules apply to the situation in which one spouse is a student or is incapacitated.

MEALS & LODGING

The value of meals or lodging and utilities furnished to an employee and the employee's family for the convenience of the employer generally is not taxable. The exclusion does not apply if the employee can take cash instead. The exclusion related to housing applies only if the employee must accept the lodging on the employer's premises as a condition of employment. GROUP LEGAL SERVICES

Amounts contributed by an employer to a group legal services plan may be excluded from income by employees, within limits. This exclusion was scheduled to expire after 1988, but has been extended until September 1990 by this Fall's tax act, the Omnibus Budget Reconciliation Act of 1988 (OBRA). EDUCATIONAL ASSISTANCE

Employees may exclude the first \$5,250 in educational benefits provided under an educational assistance program of the employer. This exclusion was also set to expire after 1988, but has been extended by OBRA until September 1990.

FRINGE BENEFITS

- No additional cost services (course play privileges, flights for airline personnel) provided by an employer to an employee for free, or at a reduced price, are excludable from the employee's income provided:
- (1) the services are for sale to customers in the ordinary course of business, and
- (2) the employer incurs no substantial additional cost in providing such services.
- Qualified employee discounts may be excluded if the property or services are ordinarily offered for sale to customers.
 There are no exclusions for discounts on personal investment or real property.
- Working condition fringes include travel advances, the use of a company car, employer provided vehicles and employer provided parking.
- Travel advances are excluded if the employee uses the advance for specific activities that are deductible, proves that payment was made, and returns any excess to the employer.
- Use of a company car for business purposes is excludable. Some "merely incidental" personal use may qualify as excludable (de minimis fringe), but use of the vehicle to commute, according to the tax code, is to be included in income. There are many employers who do not follow this reporting requirement. That is between them and the IRS, though it is possible that the employee could be assessed tax should a dispute arise.
- An employer-provided vehicle used in an employer's business is not included in your income. However, if you also use that vehicle for commuting or other personal purposes, or use it in another trade or business, the value of such use is includable in your income. Please see USE OF A COMPANY CAR, above, for further comment.
- Parking provided by the employer may be excluded, whether the employers pays directly or pays a parking allowance to the employee. (cont'd. page 24)