

Still Simply The Best Greens Fertilizers You Can Use Today

You can count on Par Ex® Greens Grade Fertilizers with IBDU® slow-release nitrogen to deliver what you're looking for in a greens fertilizer.

SAFETY IBDU® slow-release nitrogen is unmatched when it comes to application safety. With other products you may have to worry about footprinting or even burn, but not with Par Ex® Greens Fertilizers.

HIGH WIN LEVELS The key to a safe greens fertilizer, Par Ex® products continue to carry the highest WIN (water insoluble nitrogen) percentages of any line of greens products offered today. In other words, with Par Ex®, you get more of what you're paying for - more *working* WIN which delivers long-lasting results with greater safety.

CONSISTENCY IBDU® slow-release nitrogen provides the most consistent pattern of slow-release nitrogen available. Effective even in sand, Par Ex® eliminates the peaks and valleys that are typical of many other greens fertilizers.

PRODUCT VARIETY With the flexibility to offer a wide variety of formulations, you can select a Par Ex® grade that offers the exact level of WIN, phosphorus, potash and micronutrients that *your* turf needs.

For more information contact:
Vigoro Industries, Inc.
Specialty Products Division
800/767-2855

par ex
THE EX STANDS FOR EXCELLENCE

Par Ex® and IBDU® are registered trademarks of Vigoro Industries, Inc.

MIDWEST TURF EXPO

JANUARY 17 - 19, 1995
INDIANA CONVENTION CENTER
INDIANAPOLIS, IN

- ◆ GCSAA SPRAY TECHNICIANS SEMINAR
- ◆ IN-DEPTH WORKSHOPS ON MASTER PLANNING, DISEASE AND WEED CONTROL, PESTICIDE STORAGE AND RINSATE PADS
- ◆ TRADE SHOW FEATURING EQUIPMENT AND PRODUCTS FOR ALL AREAS OF TURF AND LANDSCAPE MANAGEMENT

Join your friends from Purdue University to celebrate the 50th anniversary of the Midwest Regional Turf Foundation

CONTACT JO HORN AT (317) 494-8039

Specializing in Golf Course Renovations

- Tees • Greens • Traps • Fairways
- Hydroseeding • Plant Installation •


company inc
don fiore

landscape contractors & architects
golf course contractors

28846 Nagel Court
Lake Bluff, Illinois 60044
708-234-0020
Fax 708-234-0922

Certified Member: American Society of Landscape Architects
Midwest Association of Golf Course Superintendents


The Long And The Short Of It!



If your present supplier isn't providing you with a foolproof pest and weed control formula, call on the SHAW'S TEAM! Bill Crotty and Brian McGuffin are backed by 45 years of experience in producing turf plant foods and turf protectants, as well as, over 142 years of combined staff experience in management, marketing and sales. Join the Shaw's Team for the finest turf NPK and controls products available today!

Bill Crotty (708)279-4342

Brian McGuffin (708)957-7954



KNOX FERTILIZER CO. / KNOX, IN 46534 / (219)772-6275

LESCO HAS IT!

Everything you need for professional turf management.


- Fertilizers •Control Products •Seed
- Irrigation •Equipment •Replacement Parts
- Golf Course Accessories •Safety Products
- Service Brochures • Low Voltage Lighting

(800) 321-5325 Your Growth Partner

LESCO, Inc., 20005 Lake Rd., Rocky River,
Ohio 44116

"THE PRODUCTION MACHINE"

- ✓ High Capacity
- ✓ High Floatation
- ✓ Economical
- ✓ Cutting Units Operated Individually
(use 1, 2, or all 3 units)
- ✓ 40 H.P. Diesel
- ✓ 126" Cut
- ✓ 2 or 4 Wheel Drive
- ✓ Up to 5 years lease available.



CALL:

**BILL MORGAN
TOM RALPH
PAYLINE WEST
225 N. RANDALL RD.
ST. CHARLES, IL 60174**

**708/584-8700
FAX 708/584-4453**



Groundwater, Turf Management, and Public Perception

by Michael L. Agnew, Iowa State University

Ground water is defined as any water which occurs beneath the surface of the earth in a saturated geological formation of rock or soil. It accounts for the drinking water of half the total United States population and 95 percent of the rural populations. At one time, groundwater was generally thought to be protected from contamination by impervious layers of subsoil, clay, rock, and the soils' own degradation process. However, in 1979 the pesticide Aldicarb was found in wells on Long Island and in Wisconsin. This along with detection of nitrate in groundwater, forced groundwater contamination to become the top environmental issue.

The primary sources of groundwater contamination can be classified as either point of source or nonpoint source contamination.

Point source contamination can be traced back to a specific source. In 1988, the U.S. Environmental Protection Agency reported that deficient septic tanks, leaking underground storage tanks, and agricultural activities (i.e. fertilizer application) were the most frequently cited sources of groundwater contamination.

A nonpoint source of contamination is one that cannot be traced back to a specific source. In water that did not meet state use designations by the EPA, nonpoint sources of pollution were cited as the cause of water quality degradation in 76 percent of lake acres, 65 percent of stream miles, and 45 percent of estuarine water. Examples of nonpoint sources

of contamination include agricultural fertilizer and pesticide runoff, agricultural fertilizer and pesticide movement through the soil, and sediment from construction sites.

Factors Influencing Contamination

Understanding the soil type, solubility of chemicals, water table depths, topography, and vegetation can assist in the site evaluation for groundwater protection.

Soils that have higher infiltration and percolation rates are more susceptible to groundwater contamination. Sandy soils, modified sand golf greens, and modified sand athletic fields are examples of areas having high percolation rates. With the exception of native sandy soils, these areas are constructed in a 12 to 24-inch soil profile with water diverted from the modified soils to soils with lower percolation rates. However, native sandy soils can be found in most states. These areas are highly susceptible to groundwater contamination.

The solubility of pesticides can directly influence groundwater contamination. The EPA has identified several turfgrass pesticides as having potential for leaching into the groundwater. They are Carbaryl, Chlorothalonil, 2, 4-D, DCPA, Dicamba, Fenamiphos, and Trifluralin. Only a few of these products actually remain soluble in water. The Farm Chemical Handbook provides information on pesticide solubility.

Fertilizer sources also vary in their rate of solubility. Nitrogen is more likely to move into the groundwater when present in the soil in a soluble form. Soluble forms of nitrogen include synthetic nitrogen sources (ammonium nitrate, ammonium sulfate, calcium nitrate) and urea. Slow-release nitrogen

(continued page 15)



Since 1930

A Tradition

*More than 60 years of
complete, dependable and
economical tree service.*

Nels J. Johnson Tree Experts, Inc.

Nels J. Johnson, Jr.
President

Karl G. Johnson
Vice President

Fully Insured

Main Office: 912 Pitner Avenue, Evanston, IL 60202

Telephone: (708) 475-1877

Fax: (708) 475-0037

Stop Snow Mold

With a Chipco® 26019/Daconil® Combination.



RHONE POULENC AG COMPANY
P.O. Box 12014, 2 T. W. Alexander Drive
Research Triangle Park, NC 27709
Call 800/334-9745 for product information.

CHIPCO is a registered trademark of Rhone-Poulenc.

Daconil is a registered trademark of ISK Biotech.

(Groundwater continued)

sources have a lower water solubility than the soluble forms of nitrogen. Within the slow-release nitrogen group, some slow release nitrogen sources are more soluble than others. For example, ureaform and milorganite are less soluble than short chain methylene urea.

Nitrogen Sources

Slow Release	Water Soluble
Natural Organics	Synthetic Inorganics
• Milorganite	• Ammonium Nitrate
• Sustane	• Ammonium Sulfate
• Restore	• Calcium Nitrate
Synthetic Organics	Synthetic Organics
• Ureaform	• Urea
• Methylene Urea	
• Sulphur Coated Urea	

The depth of the water table directly affects the susceptibility of the groundwater to contamination. Shallow water tables are more likely to be contaminated than deep aquifers. In Iowa, much of the drinking water is from shallow water sources.

The topography of the site also influences the movement of fertilizers and pesticides. Heavily sloped areas are more like to lose water, nutrients, and pesticides through runoff. All other conditions being the same, it stands to reason that the greater degree of slope, the greater the water loss due to the increased velocity of water flow. The length of the slope also influences the movement of fertilizers and pesticides. The

greater the extension of the sloped area, the greater the concentration of the flooding water.

The presence of vegetation on the soil surface will greatly affect the loss of fertilizers and pesticides through both runoff and leaching. The kind of grass, the thickness of the stand, and the vigor of its growth greatly affect runoff and are of great importance in the control of pesticide and fertilizer movement. A thick, healthy stand of cultivated turfgrass is much less susceptible to runoff than are pastures. Pastures are more compacted and are not as thickly vegetated as lawns. In research conducted at Penn State, nutrient loss through runoff was greater on seeded sites than on sodded turfgrass sites. The loss of water by percolation is also less on vegetated lands than bare soil. The roots of a turfgrass plant will be in the upper 8 to 12 inches of the soil profile. These roots are excellent extractors of soil water.

In summary, sandy turfgrass sites treated with soluble chemicals are more prone to leaching loss, whereas heavy clay turfgrass sites on sloped areas are more prone to runoff loss.

Management Practices That Protect Groundwater

The manager of a turfgrass site has ultimate control on protecting the groundwater. This is especially true for sandy turfgrass sites. Thus the nitrogen source, nitrogen application rates, timing of nitrogen application, and irrigation practices can directly influence groundwater contamination on sandy sites.

(continued page 18)

WE CAN DO...IBDU

BAGGED, BULK or BULK BAG FERTILIZER, WE CAN DO IT!

NOW YOU CAN TOO!

- An outstanding source of slow-release nitrogen
- Not affected by temperature
- Not affected by microbial activity
- Activated by hydrolysis
- Available in 50 pound bags
- Available in bulk truck with Lely spreaders
- Available in 1000 pound bulk bags
- No flushes of growth
- Constant color during the growing season
- Resistant to nitrogen leaching
- Low WSN and high WIN content

FOR INFORMATION, CALL:

Arthur Clesen, Inc.

543 Diens Drive • Wheeling, IL 60090 • (708) 537-2177 • FAX (708) 537-2199



TORO®

Thatcher/Seeder



MODEL 83 AERO THATCH

- Power Take-Off Driven
- Thatches or Aerates
- Forward or Reverse Rotation
- 48" Swath

MODEL 93 SEEDER

- 100 lb. Seed Capacity
- Ground Driven
- Seed Tubes for Accuracy
- Variable Seed Control


Chicago Turf & Irrigation Inc.

708-773-5555
FAX 708-773-4273

1170 W. Ardmore • Itasca, Illinois 60143



Aerators Spray Sculpture Floating Fountains



**We make the
solution to water
quality problems
crystal clear!**

708-773-5555
FAX 708-773-4273

Chicago Turf & Irrigation Inc.

1170 W. Ardmore • Itasca, Illinois 60143



Upcoming Events — Mark Your Calendar

November 8-10 — Penn State Golf Turf Conference, State College, PA

November 10 — Midwest Clinic & Annual Meeting at Medinah C.C.

November 28-30 — NCTE at Pheasant Run Resort

December 7 — CPR Seminar at Oak Brook Hills Resort, Oak Brook, IL

January 17-19 — Midwest Turf Expo, Indiana Convention Center, Indianapolis, IN

February 12-17 — Indiana-Illinois Turfgrass Short Course

February 20-27 — GCSAA Conference & Show, San Francisco

Don Maske is moving to Kansas City, MO where he will become the District Manager for Agr-Evo. Don will be starting December 1st. We all wish him well and will miss him in our area.

Ed Esgar, currently interning at Flossmoor C.C., is seeking an assistant's position. Call 708/957-5262.

Len Berg is making a BIG MOVE, he is heading to Summit, New Jersey to take over at the Canoe Brook C.C., a 36 hole private club. The club was built by Donald Ross in 1901. Len starts on December 1st. Good luck, Len and thanks for all your work with the MAGCS.

Ken Lotzer at Twin Lakes G.C. is looking for an assistant. Please call 708/934-3006. (Ken & Ed, you need to get together).

Mike Bavier and Gordon Witteveen got together on Long Island for 4½ days in late September and played 7 different golf courses.

The experimental test green at Cantigny Golf Club is finished and all are welcomed to come and visit. Give Tony a call so he can explain what all it is and will be able to do.

For Sale: Foley Reel Grinder, good condition. Call Art at Pinecrest Golf Club, 708/669-3130.

Rick Wilson, Glenview Park District Golf Course will be returfing and leveling a few tees this fall. He also likes topdressing with push spreaders using dry sand for his greens.

Art Benson Sr., One of the Deans!

by Peter Leuzinger

Art Benson Sr., passed away on October 17, 1994 from congestive heart failure. He had a heart of gold. One of Art's trademarks was his generosity, kindness and willingness to each. He taught more than greenkeeping. For example, he always considered the current president, his best friend. He told young people getting started that, "if you want to save money, spend less than you make." He understood the planning and patience needed to grow specimen trees on a golf course. He spaced them far enough apart from each other and well away from the fairway. He loved greenkeeping and the game of golf. He felt fortunate to be making a living doing something he loved. He allowed his enthusiasm to be shared with others, especially younger people.

Art Sr., worked at 3 golf courses in his life time; all in the Fox Valley. They were Fox Valley C.C. in the 30's, Aurora C.C. in the 40's and 50's and St. Charles C.C. from 1956 to 1979, when he retired completely. He was an Honorary Member at St. Charles C.C. and one of the most respected and highly regarded individuals ever to walk the grounds.

Art was a tall, friendly Swede. He always knew who to kid or what to do if a person needed a helping hand. He was a great guy and one of our best friends.

Bill Aiston, Lake Shore C.C. will be spending his time removing the bent from the banks of his greens and resodding with bluegrass.

Dan Dinelli at North Shore C.C. has a new toy; a weather station that does everything except apply the chemicals when needed. It has all of the "bells & whistles" from soil moisture sensing of a green, dew, wind, humidity, air temperature at 6" and other sensing features. Also call Dan and ask what he is doing with hydrogen peroxide and black layer.

A contribution is being donated to the American Cancer Society in the name of Terry Petric, a deceased MAGCS member. These funds are from the June meeting at Aurora C.C.

The Dave Meyer Family did very well this past year at the local and state level, in showing their livestock. At the DuPage County Fair, Austin, 17, won Champion Steer and Champion Market Barrow Hog. Sister Kara, 13, won Champion Market Lamb, which was purchased at auction by Max Armstrong for \$485.00. Then Austin won the Grand Champion Market Drouc Hog at the State Fair and it was auctioned off and Spike O'Dell won it for \$3,100.00. Well done, kids. (See page 30 for a picture of Austin and his champion hog).

This years College Golf Champion is being shared by two excellent turf schools — Penn State & Purdue University. There were eight teams entered this year and their order of standing is: Danville 3rd, University of Illinois 4th, Wisconsin 5th, Southern Illinois 6th, Michigan State 7th and last, Kishwaukee.

(Groundwater continued)

As stated previously, slow-release nitrogen sources have a lower solubility than inorganic nitrogen sources. Slow-release nitrogen sources are recommended for use on sandy soils. Research has shown that nitrate leaching is less when applied as a natural organic form (Milorganite) or a synthetic organic form (ureaform).

If soluble nitrogen sources are preferred, rates should be adjusted to prevent movement through the soil profile into the groundwater. For example, nitrogen applications with urea on high sand content golf greens should be at a rate of .1 to .25 lb. N/1000 sq. ft. per application. Anything greater may leach below the root zone. Once this occurs, the nitrogen is no longer available for plant use. However, if slow-release nitrogen sources with a high water insoluble nitrogen ratio are used, N rates can be as high as 2 lb. N/1000 sq. ft. per application on Kentucky bluegrass.

Certain types of weather will favor nitrogen leaching. For instance, cool rainy weather favors the movement of nitrogen beyond the root zone into the groundwater. Increased leaching potential occurs because cool temperatures decrease denitrification, volatilization, microbial activity and plant nutrient uptake. Thus, application of high rates of nitrogen on sandy sites during the late fall, winter or early spring can lead to nitrate movement into the groundwater.

Irrigation practices that result in water movement below the root system will increase potential nitrogen and pesticide leaching. Irrigation on a daily basis during cool months will increase leaching losses. On the other hand, infrequent deep irrigation to well below the root system will more than likely move nutrients with the water. Irrigation should only be provided to replace what water has been removed by plant uptake and evaporation.

Source: NYSTA Spring 1990 Bulletin 138

Golf Course Drainage

A Hands-On Report by Geoffrey Corlett
of Turf Drain Inc.

It is now common knowledge that good drainage is an important ingredient for the success of a golf course. Lesser known are the causes for poor drainage, and further, how to rectify the situation. Outlined in the following paragraphs are four typical drainage problems on a golf course with options that may be applied to permanently improve drainage.

High water table, low lying area

For a golf course built with a river as the predominant feature, there is a strong possibility that at least one site on the course will exhibit the following characteristics: Commonly found in a natural low point of the water shed, a basin is formed with slopes on at least three sides where surface and ground water collects. The soils exhibit the following characteristics of anaerobic conditions: A foul odour and very dark in colour. Standing water remains long after a rain and the turf is spongy throughout the season. The scale of this problem area can range from isolated pocket of less than one acre to a situation where most of the golf course is built within the basin.

Extensive drainage is required to remove the volumes of water that collect in the basin and maintain the water table at a depth conducive to growing turf. Spacing and depth of the tile is based on the grade available for fall in the drainage line and the soil characteristics. Native soil backfill is sufficient as the objective is to lower the water table, not to cut off flowing ground water. Once the depth of the soil is controlled, water holding capacity of the soil is greatly increased, allowing for the infiltration of surface water soon after a rain.

Sidehill seepage

Sidehill seepage exists anytime there are two distinct elevations separated by a well defined slope. This scenario is particularly evident where a sidehill has been excavated to accommodate a fairway. Distinctive features are soft soil along the lower run of the slope, water boiling from the ground on the lower elevation and in more severe conditions, slumping soil along the base of the slope. Sidehill seepage is the result of a head created by the ground water in the upper elevation causing an unusually high water content in the soil directly along the base of the slope.

The objective here is to intercept the ground water as it flows down the slope before reaching the lower elevation. A series of drainage lines should be installed, with granular backfill, allowing for the site characteristics for grade and outlet.

Subsoil layering

Subsoil layering is a problem not easily identified. Although water problems do occur naturally as a result of subsoil layering, they are most pronounced when the native subsoil has been disturbed through excavating and forming the architecture of a golf course. The natural ground water flow is altered by a new and non-uniform arrangement of materials in the subsoil, such as a pocket with high permeability isolated by a impervious layer of clay. Small variances in the texture of the subsoil are difficult to detect without removing a three

(continued page 20)

WE MOVED ...

CLAUSS BROTHERS, INC.

OUR NEW ADDRESS:

**360 SCHAUMBURG ROAD
STREAMWOOD, IL 60107**

PHONE: (708) 830-7405

FAX: (708) 830-9652

EFFECTIVE OCTOBER 1ST

Work from the soil up with Nature Safe®.

- Homogeneous
- Non-Burning
- Micro-Nutrient Rich

Creating a soil
structure
to grow turf.

**Nature
Safe**
Natural & Organic Fertilizers

(800)252-4727

(606)572-2549



Also available in
10-3-3 and 7-1-14

42ND ANNUAL

MIDWEST TURF CLINIC

SPONSORED BY:

MIDWEST ASSOCIATION OF GOLF COURSE SUPERINTENDENTS

THURSDAY, NOVEMBER 10, 1994
MEDINAH COUNTRY CLUB

PROGRAM SCHEDULE

9:00 - 10:00	Registration - coffee and rolls
10:00 - 10:10	Opening remarks
10:10 - 10:30	Mr. Don Ferreri Superintendent, Seven Bridges Golf Club "The Wonderful World of Wetlands"
10:30 - 11:00	Mr. Dave Ward Superintendent, Olympia Fields Country Club "Establishing Native Prairies"
11:00 - 12:00	Mr. Connor Shaw Owner, Possibility Place Nursery "How To Succeed With Native Woody Plants on the Golf Course"
12:00 - 1:15	Lunch
1:15 - 2:30	Mr. Joseph O'Brien Chief Operations Officer, GCSAA "The Changing Times for the Industry, Association and Profession"
2:30 - 3:00	Mr. Ed Hightower NCAA Final Four Basketball Official "Making the Right Call"
3:15 - 5:00	M.A.G.C.S. Annual Meeting
5:30	Dinner

A Commitment to Quality Turf Care Products For You - The Turf Care Professional



Your Tee Time Distributor Offers You:

- Greens and Fairway Fertilizers with Nutralene
- Quality Sulfur Coated Urea Fertilizer
- Combination Products
- Granular Insecticides
- Granular Fungicides

For additional product
information, contact
Craig Filley 1 (800) 225-2639

**The
Andersons**

Lawn Fertilizer Group
P.O. Box 119
Maumee, Ohio 43537
1 (800) 225-2639

(Golf Course Drainage continued)

or four foot core sample. Poor drainage under these conditions will appear in several forms. A spring boiling from an otherwise well draining fairway with moderate grade and spongy pockets randomly located in an area which appears to be well graded. (See fig. 1)

Once identified, drainage lines should be installed continuous with the clay outcrop to a suitable outlet. If the wet area is localized, three to fifteen meters in diameter, one or two lines into the problem area are sufficient to remove the trapped ground water.

Surface grading

It is a well known fact that high quality surface grading is critical to the success of every golf course. Unfortunately, in practice, surface grading rarely, is well executed in both design and construction. The result area areas that suffer extensive erosion while on the same project there will exist an area with pockets of standing water scattered throughout.




Fig. 1. Ground water movement restricted by impermeable layer of clay. Result is an underground ponding area.

Depressions in the surface grade are the most visible and inconvenient water problem for the membership. The condition is mostly pronounced on fairways with topsoil having more than 30% clay content, as the surface water is very slow to infiltrate after a rain or an irrigation application. For fairways with minimal grade, it is critical that depressions are eliminated during construction.


Due to numbers and random locations of depressions, the best approach is the installation of a main line initiating at an open outlet or tied into an existing line. From the main, several short lines are installed passing through each depression.

Erosion as a result of excess water flow through a restricted area is evidence of misformed or improperly designed surface grading. Soil type will determine the extent of soil movement. Sandy and gravelly soils are most subjected to erosions.

Total acreage that is supplying surface water to the area must be determined in order to calculate volumes of flow during peak periods of the year. To demonstrate this point, a hypothetical situation is outlined:

(continued page 22)

PRESSURE WASHERS



250
to
2,000
PSI

NEW AND USED

FREE DEMO

- SALES
- SERVICE
- PARTS
- CHEMICALS

CALL TOLL FREE 1-800-666-3900

*The Cleaning Equipment Specialists Since 1976**



TIM LAYDEN
General Manager

708-893-0777 • FAX (708) 893-1045

1-800-666-3900
Sales Service Parts Rentals Chemicals
25 S. Park Street • Roselle, Illinois 60172