flexible enough ...

...to gather the best quality and technology available. ...to respond to future market opportunities. ...to deliver genuine value to the turf manager.

United Horticultural Supply has earned a leadership role in discovering and delivering the best quality and technology available on the market.

Competitors tied to limited resources or technologies cannot make that statement.

That's why you can trust the performance and quality of UHS Signature Brand Fertilizers.™

Contact your local UHS representative for more information.

(800) 411-5115 www.uhsonline.com

UHS Signature Brand Fertilizers are a product of United Horticultural Supply



"you have our name on it"



BEST TURF MOWERS FOR A QUALITY CUT.

The Bush Hog TD Series Tri-Deck Finishing Mowers are available in 15 or 17-foot cutting widths and are ideal for sod growers, golf courses, land owners, municipalities and recreational sites.

These mowers are designed for precise cutting on even or uneven terrain . . . they feature a 30° up and 10° down wing flex and are rated for 40 and 45 minimum tractor horsepower. Gearbox rpm is 540, and deck construction is rugged, 7-gauge steel. Blade tip speeds are over 15,500 fpm, to give you finer, cleaner cutting on each pass. Wing decks flex side to side and front to rear, for closer mowing.

Come in today and see why Bush Hog Tri-Deck Finishing Mowers are clearly your best choice for quality.



BUSH HOG® DIVISION OF ALLIED PRODUCTS CORPORATION Building the Legend.

Roseman Tractor Equipment Co. Inc 2620 Crawford Avenue Evanston, IL 60201 847-864-1842 Fax 847-864-1938

Using Composts to Improve Turf Ecology

F. Dan Dinelli

t is said that if any of the billions of organisms inhabiting the soil had hands, the fate of the world would be in them. An important soil function is the harboring of a diverse community of organisms that includes bacteria, fungi, protozoa, nematodes, mites, springtails, millipedes, sowbugs, earthworms and many others. This community drives the decomposition of organic residues; recycles important nutrients like carbon, nitrogen and phosphorus; and contributes to the formation of new soil and soil structure. With these activities, soil organisms contribute to other important soil functions, such as supporting the growth of plants and absorbing, neutralizing and transforming compounds that might otherwise become pollutants in the environment. Basically, soil organisms play a critical role in shaping and maintaining terrestrial communities and ecosystems.

As the superintendent at North Shore C.C. in Glenview, IL, I became interested in applying compost as a soil amendment after reading research suggesting its many agricultural benefits. Dr. Michael Boehm, Ohio State University, and Dr. Eric Nelson, Cornell University, have done helpful work, specifically about the effects of compost on turfgrass. Generally, researchers and practitioners recognize that incorporating high-quality compost does several things:

1. Adds food for nearly every kind of organism needed by a healthy soil.



Topdressing fairways with compost.



Testing different composts on bentgrass mowed at 1/8".

2. Adds a diversity of organisms to the soil.

3. Encourages plant growthpromoting substances in soils. Compost can also have an effect on soil structure, nutrient cycling, disease suppression, nematodes and other biological activity.

In fact, the use of composts on turf is not new. A book given to me by my grandfather, Frank Dinelli (retired greenkeeper at Northmoor C.C.), titled *Turf For Golf Courses*, by Charles V. Piper and Russell A. Oakley, printed in 1917, has a chapter devoted to "Manures, Composts and other Humus Materials." Yet because compost is not widely used on golf courses, I wanted to participate in further research prior to investing in the process at North Shore C.C.

Phase I: Experimentation

In 1996, we got just that opportunity by participating in a two-year study of various composts and organic materials under the direction of Dr. Michael Cole of the University of Illinois and (continued on page 14)

Only With A Diamond

Diamond FriKut **5000**

Diamond Turf

quipment Inc.

ONew - Built new to exacting standards.

●Affordable – Our factory-direct agency assures you the best price possible. Call today for current pricing.

Dependable – DTE has been manufacturing greens equipment since 1982. Only top quality OEM parts are used, engines such as Honda assure dependability.

OReliable – Every product is backed by a two year bumper-to-bumper warranty. Diamond's exclusive Five-Star Service Program is unmatched in the industry. It's designed to keep you running and get you back up fast should repairs or parts be needed.



Aerate 18 greens in 3 hours without disrupting play or an average fairway in about 2 hours For greens and tees, AerWay's new tow-behind Greens Express penetrates 6" without cores, and won't disrupt play.

(815) 469-8500 (708) 301-8500

The roller leaves greens instantly playable. Aerway lets you aerate whenever you want, for better turf conditions, and happier golfers.

For fairways and roughs, patented Aer-Way Shatter Tines relieve compaction 7" and deeper. Recovery is fast: a mowing pass the opposite direction and you're ready for play. No complex machinery. No cores. No PTO. Take a look at Aerway today.

For the name of your nearest dealer contact:

Tee Shot Marketing Inc. 800-950-4288



Using Composts to Improve . . .

(continued from page 12)

GreenCycle, Inc. (operator of several composting facilities) of Northfield, IL. The study was a replicated 10' x 10' plot design on our #5 fairway comprised of creeping bentgrass and Poa annua maintained at 1/2" mowing height. During the field evaluation, all observations were noted. However, our main objective

was to observe any diseasesymptom differential between the various plots.

Our first application was in the fall of 1996 to observe snow mold (*Gerlachia nivalis, Typhula spp.*) suppression. None of the materials demonstrated any noticeable snow mold suppression. However, plots treated with compost had a notably earlier green-up and recovery rate versus the control plots. We then repeat-

ed applications in late spring of 1997. Observations through the remaining growing season showed strong dollar spot (*Sclerotinia homoeocarpa*) suppression—up to 80% reduction; improved turf color and density; and increased earthworm castings. Thus, while our initial objective of snow mold suppression was not observed, our experiment to test organic products as they improve overall turf ecology proved quite successful.

Phase II: Implementation

Based on favorable results after two seasons of field evaluation of compost topdressing, we implemented the strategy on all our fairways. During our normal coring of fairways, the process involves the following steps:

- Coring with hollow tines;
- Breaking up the soil cores with a vertical mower;
- Topdressing with compost;

- Mixing the soil with compost as it is matted into the surface with a section of chain-link fence;
- Blowing the remaining tufts of turf and thatch into rough via a three-point hitch blower;
- Picking up debris in the rough with an out-front rotary mower fitted with a bagging attachment; and



'Homemade' compost tea.

• Irrigating the area well.

We have been coring fairways like this for several years. Adding the extra step of compost topdressing has not significantly impacted the workload. The cleanup is about the same and we can still get our targeted nine holes (15 acres) done in one day.

(Note: Part of our IPM cultural program is poling, by dragging a chain over the fairways each morning to remove leaf moisture and guttation. This process also manages earthworm casting buildup).

Phase III: Results

The results so far are much the same as in the test plots: improved turf density and color, rapid healing of cored turf, dollar spot *(Sclerotinia homoeocarpa)* suppression, increased earthworm castings and thatch reduction have been observed. We continue to monitor the impacts of compost use on turf and maintain computerized spreadsheets to evaluate our results. Over time and continued applications, we hope to document improved soil structure and suppression of other diseases.

Selecting Quality Compost Is Key

Selecting quality compost is very important; you have to do your homework. Compost products are not yet standardized, so the chal-

lenge is in obtaining consistent, high-quality compost. The procedure we use to assure the compost we obtain is optimal for our turf involves a series of tests. We analyze chemical, physical and biological activity.

- Chemical Analysis In the chemical analysis we look for:

- Carbon:nitrogen ratio <20:1, best at 15:1.
- pH at 6.5-8.5.
- None to trace amounts of ammonium, sulfide and nitrite.
- Low concentrations of soluble salts, especially sodium.
- Moreover, we strive towards elemental balance and recommended ratios favoring the high side of potassium and calcium. Also, biosolids need to meet U.S. EPA's Part 503 technical rule for biosolids. All biosolids tested for coliform and other diseases. Biosolids composted properly have been heated sufficiently to kill viruses, coliform and other diseases. Metals in biosolids are often high and should be considered.

- *Physical Analysis* Physically we look for:

• Fine texture < or = 1/8". (continued on page 16)



Leading the industry in turf and ornamental Management

1

1

1

Trent Bradford Tim Keating John Meyer

630-514-8749 630-514-8997 630-514-8748 **Eric Foerster** John Lebedevs Bruce Schweiger

630-514-8750 630-514-8745 630-514-8763

2N255 County Farm Road, West Chicago, IL 60185

Office: Fax:

630-231-8441 630-231-8453 800-383-3230 800-457-7322

or

or

Where Success is based on a commitment to people...

PETROLEUM TECHNOLOGIES EQUIPMENT. INC.

ONE TOWER LANE . SUITE 1700 OAKBROOK TERRACE, ILLINOIS 60181 PHONE 630-573-2910 . FAX 630-579-9087

"Don't Be Caught Off Guard By The 1998 Underground Storage Tank Regulations Act"

SERVICES OFFERED

• Underground Tank Removal, Installation & Upgrades

Above Ground Tank Installation

- Petroleum Equipment Distributors
 - Distribution & Installation of Compressed Natural Gas Equipment
 - Pump Sales & Service
 - General Contracting
 - Key and Card Fuel Management Systems

"FULL SERVICE PETROLEUM EQUIPMENT INSTALLATION SPECIALISTS"

Using Composts to Improve ... (continued from page 14)

- Light, crumbly structure, parent material nonvisible.
- Moisture at 30-40%.
- Dark brown to black in color (caution with dark black compost, for the compost might have gotten too hot and burned).

- Microbiological Analysis

A great diversity of bacteria, fungi, protozoa and beneficial nematodes occur in good compost. Healthy compost has been hypothesized to have between 10,000 to 20,000 species of bacteria per gram. The DNA analysis required to establish the set of species in highly diverse compost awaits molecular microbiologists. The study of microorganisms is very complex and dynamic. Many of the billions of organisms that exist have not yet been described or their functions understood. It is a frontier ready to be explored.

Several laboratories and universities are exploring methods to assess soil and compost microbial activity. Soil Foodweb Inc. has done testing for us, including bacteria, fungi, protozoa and nematode counts. Some feel (continued on page 18)

| | | STANDARDS FOR COMPOSTS (SOILS) | AGRI-ENERGY COMPOST | BREW WASTE | EARTHWORM CASTS | GREENCYCLE | BIOSOLIDS | WOOD BIOCOMPOSTE | B & G COMPOST |
|-----------------------------|------------|--------------------------------------|------------------------|---------------|--------------------|------------|-----------|---------------------|------------------|
| Sample Number | | (5 - 1 - 2 / | 73 | 63 | 72 | 217 | 310 | 554-1 | 555-1 |
| Date | | | 3/18/98 | 3/31/98 | 3/18/98 | 5/22/96 | 8/4/97 | 12/17/98 | 12/17/98 |
| Aerobic Plate | | | 6.00E+08 | 1.10E-09 | 4.50E+08 | 2.50E+09 | 1.20E+10 | 2.10E+09 | 6.00E+08 |
| Anaerobic | | 10:1 Ana:Aer | 1.80E+07 | 9.40E+08 | 1.10E+08 | 2.50E+08 | 5.30E+08 | 8.00E+07 | 2.30E+05 |
| Yeasts/Molds | | 1E+3 to 1E+4 | 1.10E+05 | 1.60E+05 | 1.40E+06 | 9.70E+05 | 1.60E+04 | 1.70E+05 | 1.20E+05 |
| Actinomycetes | | 1e+6 to 1E+8 | 1.30E+07 | 1.10E+03 | 1.10E+07 | 2.30E+07 | 2.20E+06 | 3.10E+07 | 8.30E+05 |
| Pseudomonads | | 1E+3 to 1E+6 | 2.90E+05 | <1.00E+02 | 2.00E+05 | 2.50E+05 | 2.10E+06 | 1.40E+05 | 3.50E+05 |
| N-Fixing Bacteria | | 1E+3 to 1E+6 | 2.00E+04 | <1.00E +03 | 7.70E+05 | 3.00E+06 | 1.50E+05 | 6.70E+05 | 2.00E+03 |
| Soil Moisture | | 30 to 40 % | 57 | | 85 | 78 | 91 | 76 | 13 |
| Total Species Richness Di | versity | | | | | 10.4 | | 9.4 | 3 |
| Total CEC (meg/100g) | | >60meq/100g | 36.35 | 6.66 | 38.56 | 49.79 | 36.92 | 39.54 | 58.24 |
| pH | | 6.5 to 8 | 7.6 | 5.4 | 6.7 | 7 | 7.4 | 6.7 | 7.2 |
| Organic Matter (%) | | 20 to 35% | 13.4 | 73.2 | 37.5 | 28.5 | 27.1 | 32.74 | 9.25 |
| Estimated N Release | | | | >130 | >130 | >130 | >130 | 121 | |
| PHOSPHOROUS | | | | | | | | | |
| Easily Extractable P | P as P2O5 | | 4403 | 1289 | 249 | 2760 | 10659 | 2171 | 11661 |
| Bray II | ppm of P | | 961 | 281 | 54 | 603 | 2327 | 474 | 2546 |
| | P as P205 | | 1085 | 85 | 376 | 1025 | 1045 | 916 | 870 |
| | ppm of P | | 237 | 19 | 82 | 224 | 228 | 200 | 190 |
| | | | | | | | | | |
| SOLUBLE SULFUR | | 15 to 40 | 184 | 28 | 188 | 147 | 1937 | 615 | 753 |
| | ppm | 15 10 40 | 104 | 20 | 100 | 147 | 1357 | 015 | , |
| EXCHANGE CATIONS | | | | | | | | | |
| calcium | lb/A | | 5748 | 602 | 10798 | 10066 | 6929 | 10476 | 7196 |
| | ppm | 1300 to 3000 | 2874 | 301 | 5399 | 5033 | 3465 | 5238 | 3598 |
| magnesium | lb/A | | 2088 | 526 | 1574 | 2552 | 3338 | 1512 | 3776 |
| | ppm | 140 to 300 | 1044 | 263 | 787 | 1276 | 1669 | 756 | 1888 |
| potassium | lb/A | | 7940 | 90 | 662 | 8658 | 1710 | 2050 | 11818 |
| | ppm | >140 | 3970 | 45 | 331 | 4329 | 855 | 1025 | 5909 |
| sodium | lb/A | | 790 | 96 | 280 | 324 | 932 | 360 | 3182 |
| | ppm | <67 | 395 | 48 | 140 | 162 | 466 | 180 | 1591 |
| BASE SATURATION % | | | | | | | | | |
| calcium | % | 60 to 85% | 39.53 | 22.6 | 70.01 | 50.54 | 46.91 | 66.24 | 30.89 |
| magnesium | % | 8 to 12% | 23.93 | 32.91 | 17.01 | 21.36 | 37.66 | 15.93 | 27.0 |
| potassium | | 5 to 8% | 28 | 1.73 | 2.2 | 22.29 | 5.94 | 6.65 | 26.02 |
| sodium | | 3% | 4.72 | 3.13 | 1.58 | 1.41 | 5.49 | 1.98 | 11.8 |
| other bases | And States | | 3.8 | 6.6 | 4.7 | 4.4 | 4 | 4.7 | 4.3 |
| hydrogen | % | | 0 | 33 | 4.5 | 0 | 0 | 4.5 | 1 |
| 5.5-7 | | | | | | | | | |
| EXTRACTABLE MINORS boron | | 0.4 to 1.5 ppm | 1.77 | 0.61 | 1.11 | 5.46 | 2.88 | 2.44 | 3.9 |
| iron | | >50 ppm | 157 | 19 | 288 | 320 | 351 | 363 | 28 |
| manganese | | 8 to 35 ppm | 85 | 4 | 8 | 60 | 74 | 74 | 120 |
| copper | | 0.6 to 2 ppm | 3.48 | 0.77 | 2.61 | 3.68 | 14 | 5.46 | 21.1 |
| zinc | | 1.3 to 3.5 ppm | 36.8 | 6.8 | 4 | 25.7 | 198.7 | 73.27 | 95.5 |
| aluminum | | 1.5 to 5.5 ppm | 141 | 23 | 28 | 64 | 752 | 711 | 9 |
| | | | 141 | 23 | 20 | 04 | 152 | | |



excellent at reducing the impact of stress and stimulating root growth through enhanced water and nutrient uptake. In addition, **BioGain**(Fe) contains a 10% fully chelated iron for use with turf that displays an iron deficiency.

For added efficiency, use **Canteen**[™] yucca extract based spreader and soil penetrant to deliver **BioGain** more effectively into the crown and root zone. *BioGain and Canteen*—the natural way to



Everyone needs a great tree to enjoy a good book. Surumn Tree Care Expense

Autumn Tree Care Experts, Inc.

Porking With Nato

A full service arboricultural firm dedicated to tree preservation since 1979.

847.729.1963

2091 Johns Court • Glenview, IL 60025 www.autumntree.com

Using Composts to Improve . . . (continued from page 16)

nematodes could be used as a biological indicator. Four basic types of nematodes occur: bacterial-feeders, fungal-feeders, root-feeders and predatory nematodes, which feed on other nematodes. When organisms increase, an increase in those nematodes that feed on them occurs, thus providing an indication of relevance.

BBC Laboratories has also done testing for us (see chart). They analyze concentrations of six groups of organisms, which they feel are "key players" in soil ecology. The following six functional groups tested are:

- Heterotrophic (aerobic) bacteria—Finished compost should have 100 million to 10 billion colony forming units/gram dryweight (CFU/gdw). Compost with less than 100 million CFU/gdw will not perform as well as soil innoculants and may not be effective in suppressing plant diseases.
- Yeasts and molds (fungi)— Finished compost should have between 1,000 and 10,000 CFU/gdw. These organisms are important for breaking down organic compounds, soil nutrient cycling, stabilizing soil aggregates and controlling plant disease.
- Nitrogen-fixing bacteria—The number of free-living nitrogenfixing bacteria in compost varies a lot depending on the available nitrogen concentration but may be in the range of 1,000 to 1 million CFU/gdw. The populations of these free-living nitrogen-fixing bacteria will proliferate as the available nitrogen in the compost decreases. As a consequence, there is typically an inverse relationship between biologically available

nitrogen in the compost and the concentration of free-living nitrogen-fixing bacteria.

- Actinomycetes—Finished compost should have at least 1 million to 100 million CFU/gdw. Compost made with woody materials may have more. These organisms are important for many functions, including the breakdown and nutrient cycling of complex chemical substances such as chitin and cellulose, improving soil crumb structure and assisting in the reduction of plant pathogen pressures. They are particularly efficient in alkaline soils.
- Anaerobic bacteria—The ratio of aerobes to anaerobes in the compost should be at least 10:1 or greater. An overgrowth of anaerobes indicates the compost was not turned with sufficient frequency. It is important that anaerobic by-products in the

compost be degraded prior to use with plants of germinating seeds.

• Pseudomonads—Finished compost concentration should be between 1,000 and 1 million CFU/gdw. Depending on starting materials, this number could be lower, but is rarely higher. Pseudomonads are important in nutrient cycling, assisting plants with phosphorus availability, and some have been linked to the biological control of plant pathogens.

In addition, compost needs to be free of contaminants, such as weed seeds, plant parts, pathogens, stones, plastic, glass, wood, nails, etc. Compost also needs to be "mature," testing >50% on the maturity index, by BBC Labs. In-house maturity tests can be performed by planting grass seed in a pot, utilizing the (continued on page 20)





The Measurement of Perfect Turf.

For over 20 years, IBDU slow release nitrogen has been the university benchmark on all types of turf. And it's available through Par Ex blended fertilizer. So you get a premium fertilizer while your turf gets the highest available WIN. Par Ex is available in four size grades and multiple formulations. Call your nearest Par Ex

distributor for information. Or 1-800-233-0628.









Using Composts to Improve . . .

(continued from page 18)

intended compost as the growing medium, to observe seedling health and establishment. Another method is to fill a plastic bag with intended moist compost and allowing it to sit sealed in the sun for a few days. Upon opening the bag, the compost should have an earthy smell, not an offensive smell from ammonia or sulfur.

Following these procedures will help insure favorable results. Adverse effects can result when utilizing poor-quality compost. Starting slow and testing small areas first is always helpful. Developing a working relationship with local composters will help in understanding their product.

Additional Uses for Compost with Turf

In addition to our fairway compost topdressing program, we also use compost in our "soil and seed" mix for divot repair. Compost is used as topdressing while overseeding turf. In 1998, we constructed a 7,000-square-foot experimental putting green, featuring 20 different root zone mixes. Each mix used USGAapproved sand in a USGA root zone profile with various organic and inorganic amendments. The 90/10 sand/compost plots outperformed the others considerably in seedling establishment and development. We continue monitoring other effects as the putting green matures. Compost tea is made and applied as a protective biofilm on the phylloplane and to deliver plant growth promoting substances.

The Bottom Line

To apply compost topdressing to fairways we purchased a TY-Crop MH-400 for \$20,000. This material hauler/topdresser is used for other tasks as well, such as rapid refill of materials while topdressing greens and tees and applying sand in bunkers. The compost we currently use is a 50/50 mix of yard trimming compost and biosolids. Our cost for yard compost trimming is \$14.00/cubic yard. For us now, biosolids are freely available (EPA permits are needed). The rate used is approximately 17 vards (7 tons)/acre = 1/8"laver. Total material cost is \$119/acre. We offset some costs by reducing our other fertility inputs and decreasing fungicide treatments as part of our IPM program.

All Composts Are Not Created Equal

Understanding the chemistry, biology and science of compost is complicated. Parent material used, how it's managed during composting, and storage can all have a huge effect on the finished product and results. Yet our efforts to understand compost, particularly its microbial benefits, have paid off. Results using composts have been positive and the turf ecology is improving under our growing conditions.

To find out more about the composting practices at North Shore C.C., contact superintendent Dan Dinelli, CGCS, at 847-724-4963, or via e-mail at ddinelli@aol.com.

A NEW ADDITION TO OUR BULK FERTILIZER PROGRAM... LEBANON FERTILIZER IN 1000 POUND BULK BAGS!

- FEATURING HOMOGENOUS PRODUCTS
- REDUCES DISPOSAL AND LABOR COSTS
- USE THE BULK BAG LIFTER WITH ANY FRONT-END LOADER
- APPLY FERTILIZER ACCORDING TO YOUR SCHEDULE
- FEATURES AN EASY TO CLOSE SHUT-OFF BAG

543 Diens Dr. Wheeling, IL 60090 847-537-2177 847-537-2210 FAX 8050 W. 186th St. Tinley Park, IL 60477 708-444-2177 708-444-2199 FAX ARTHUR CLESEN INC.