

# MSMA

## A loaded gun to your head?

By Dr. Phil Busey

That's harsh language, but it metaphorically describes the situation of those of us who use MSMA. Monosodium methanearsonate (MSMA) is the most effective broad-spectrum postemergence grass herbicide for use in bermudagrass golf and sports turf. It's also a heavy element and Class A human carcinogen.

Each typical application of MSMA adds to the environment 1.0 kg/ha (nearly one pound per acre) of elemental arsenic. Two to four applications may be required for a single series of treatments to control grass weeds such as goosegrass.

Arsenic has only slight volatility, that is, tendency to evaporate; therefore additional applications will, for the most part, either accumulate at increasingly higher concentrations in soil, or move in water. The Florida Department of Environmental Protection's Leslie Smith said that, south of Orlando, one should be concerned about both a groundwater and soil problem. Her remarks (including the loaded-gun expression) were made at the United States Golf Association Green Section Regional Conference in Palm Beach Gardens, Nov. 14.

One attendee asked Leslie, "What's its half-life?" She correctly pointed out that as an element, arsenic never breaks down. To emphasize that in my own way, I say, "its half-life is infinity."

So you can study to no end the complex models for the transformation of arsenic compounds, some of which are more toxic than others, but the bottom line is, the majority of the arsenic stays around. (Unless the owner pays someone to excavate the golf course or sports field and move the arsenic to someone else's property.) What little leaches into the groundwater may be a serious health problem in some cases, but will not diminish greatly what is accumulated in the soil.

If a single application of MSMA were mixed thoroughly in the top 15 cm (about 6 inches) of soil, based on a bulk density of 1.5, there would be a concentration of 474 ppb elemental arsenic in the soil throughout the root zone. This exceeds the existing 400 ppb soil screening level of the US

Environmental Protection Administration. In one application. If even a small percentage of the arsenic reached the drinking water, it would also exceed the EPA limit of 10 ppb arsenic in drinking water, depending on the level of dilution.

Houston, we have a problem.

Or do we? Ironically, MSMA has a very low acute toxicity, based on a high LD50. It takes a relatively large lethal dose to kill 50 percent of laboratory animals. The acute oral LD50 is 2833 mg/kg for rats. For an 80-kilogram rat, about the weight of an adult human male, that would be equal to about 9 shot glasses, a pretty distasteful and unrealistic way to die. The chronic effects of small doses are not easily predicted, but arsenic is not good to be in contact with. If I were an 80-kilogram rat, I'd be more con-

cerned about chronic exposure to MSMA, than the unlikely shot glass. And that's why I always wear protective clothing, whatever I am spraying.

### When was MSMA first used in turf in Florida?

MSMA was first used publicly in 1963, and by 1967 Dr. Evert O. Burt of the University of Florida, Fort Lauderdale, reported that it was equal or slightly more effective than disodium methanearsonate (DSMA) for grassy weed control in bermudagrass.

Although that was a long time ago, both DSMA and MSMA were already late arrivals. The arsenicals as a group were the first chemicals widely tested for chemical weed control. Long before the 1942 discovery of the phenoxyacetic acid herbicides (e.g., 2,4-D), the US Army

Corps of Engineers was using sodium arsenite for control of water hyacinth in Louisiana, in 1902.

Some of the early arsenical products used in Florida turfgrass include the Florida East Coast Fertilizer Company's S.A.M.A. 70, a monosodium arsenical, possibly MSMA, and Dal-E Rad 70, a DSMA powder by Vineland Chemical Company. These products usually required four applications to give a serious chance to eliminate goosegrass, especially the more mature goosegrass. Very mature goosegrass could not be controlled.

Aware of the hazards of arsenic, scientists attempted for years to find replacements. In describing metribuzin as such a prospect in 1979, the University of Georgia's Dr. B. J. Johnson said as an afterthought, "If EPA takes MSMA off the market, we may be left with Sencor as our base material."

Rather than seeing MSMA disappear, however, Dr. Johnson made it better by developing (with others) the synergistic combination of tank mixtures with the triazine herbicide metribuzin.

Sencor was Mobay Chemical Company's formulation of metribuzin, and Dupont



had attempted to develop another formulation called Lexon. Another chemical, methazole, formulated as Probe, was being looked at along with metribuzin, but by 1993 its herbicide registration was voluntarily canceled by Sandoz Agro, Inc.

It was the MSMA + metribuzin tank mixture that proved to be very effective at selective control of goosegrass, even mature goosegrass, in bermudagrass turf. The number of sequential applications of MSMA could be reduced from four to two, and with a little bit of metribuzin as Sencor, there would be better goosegrass control than with MSMA alone.

I was present at a 1976 meeting in Arkansas when Dr. Johnson described the promise of metribuzin for goosegrass control in fairways. By 1978, Dr. Max Brown described in the South Florida Green (Volume 5 No. 3) that Sencor could be used for grass weed control. But tank mixtures with MSMA were not mentioned.

### History in the Making

The big breakthrough for Florida golf course superintendents and sports turf managers came around 1979, when Dr. B. J. Johnson described metribuzin as the "best product researched and now on the market" for grass weed control in bermudagrass, and he described a 1/8-pound-per-acre active ingredient metribuzin tank-mixed with 2 pounds active ingredient MSMA. At that time he had done some three years of research on MSMA + metribuzin tank mixtures. The interview was conducted by Dave Bailey, at that time superintendent of Atlantis Country

Club, and staff writer for *The South Florida Green* which was edited by Dan Jones.

There was also a flurry of abstracts (not full scientific reports), also in 1979, by Dr. Johnson, as well as by the University of Arkansas's Dr. John King, and the University of Florida's J. A. Tucker and Dr. Wayne L. Currey. There followed a full scientific article on the subject by Dr. Johnson, in 1980. I first became aware of the MSMA+metribuzin tank mix in the summer of 1980, when my bermudagrass breeding plots were overrun with crowfootgrass. But I opted not to include the metribuzin because it was still too new, and I didn't want to take a chance of messing up my experiment.

### The Lost Discovery?

One of the interesting mysteries about the MSMA + metribuzin tank mixture is that the first scientific report goes back to 1974. This synergistic mixture was reported in *Agronomy Journal*, a widely disseminated journal, by the University of Hawaii's Dr. Chuck L. Murdoch and David Ikeda. Dr. Johnson was aware of that paper in 1975, because he cited Murdoch and Ikeda's work when he published a study involving MSMA and metribuzin. But the most novel aspect of the Hawaii paper was the tank mixture, which was not a part of Johnson's studies until later.

In conclusion, MSMA appears to have been used in Florida since the mid-1960s for post-emergence goosegrass and other grassy weed control in bermudagrass turf. DSMA had formerly been used for the same purpose, but did become established.

MSMA was not very effective against mature goosegrass until 1979 or 1980, when the MSMA + metribuzin tank mixture swept the industry based largely on the research of Dr. B. J. Johnson. The same mixture was reported, and appears to have been ignored, from work in Hawaii in the early 1970s.

These dates are approximate, but based on written documentation. If anyone has a better memory of the history of MSMA use, I would appreciate hearing from you.

### Peter Harrison responds on Organic Arsenicals (MSMA question)

I have no issue with the question mark over using, and the soil accumulation data regarding the organic arsenicals. However, the arsenic while it remains in the soil is an unlikely problem for workers and users, and I also support your comments regarding sprayer user and protection, where chronic problems may occur, although I am not aware of reported problems. Movement from soil is a medium-term problem and of increasing interest.

Arsenic in soil and water, esp. mobile forms is an ongoing issue in a number of areas including widespread problems in at least one country (Bangladesh, where well water can be serious health problem if drunk), old precious-metal mining areas, older tanneries (where arsenic compounds were used at times) even soil in old animal yards and quite a lot of work is being done and some has been published about converting this to nonmobile forms in the soil/water. Some soil microbial solutions are being

## DRAGONFIRE-CPP™

# Organic Nematicide

DECREASE IN NEMATODES

Treatment	Change in Nematodes
Dragonfire-CPP	-63%
NemaCur	-32.5%
Control	+9%
Sesame Oil	+8.3%

(NemaCur® is a registered trademark of the Bayer Corporation)

- No REI
- EPA Exempt
- Sprayable
- Effective
- 100% Natural
- Registered Nematicide(FL)

University Tested

U.S. Patents Pending

**Turf Maintenance Products**

## NaEx™

Reduces Soil Sodium Levels

## NitroLoc™

Reduces Nitrate Leaching

## JETWET™

Wetting Agent/Soil Penetrant

**POULENGER USA, Inc.**  
Science in Sync with Nature

1-866-709-8102

863-709-8102 Fax: 863-644-4038

www.poulengerusa.com

touted, among various options.

It will not “go away” as you point out, but it may be made a minor issue for those areas where transfer to groundwater, etc. is a problem. Solutions to the issue may arise from left field in areas of bioremediation and phytoremediation sciences... but there is no doubt that these products do an excellent job with weeds... including many sedges, notwithstanding newer products that have become available. Often their use can be moderated by practicing some rotational use among products, a sound ecological practice anyway, although immediate costs are sure to rise.

Similar issues and concepts over the arsenicals are receiving thoughts in Australia, so Florida is not alone. I am looking forward to the balance of the articles.

*Peter G Harrison*

*Principal*

*Above Capricorn Technologies*

*Agricultural and Environmental Consultants*

*PO Box 736, Nightcliff 0814 Australia*

**MSMA vs. other sources of Arsenic:  
Phil Busey replies to Peter Harrison**

Peter correctly points out the global aspects of arsenic pollution arising from many sources. In Florida and other warm U.S. states, we have leftover CDVs (Cattle Dipping Vats) used by government mandate from 1917 to 1944 to dunk cattle in an arsenic solution (probably arsenic trioxide) to successfully eradicate Cattle Fever Ticks that were causing 50 percent mortality. This nice brew was aug-

mented later with DDT, toxaphene, and chlordane, and the arsenic was later dropped. Cattle could only be moved short distances; therefore Florida has a legacy of 3200 CDVs with a high concentration of arsenic in the soil.

Since then, many “safer” acaricides and insecticides and anthelmintics have been developed for veterinary use in the tropics, subtropics, and warm areas where animal husbandry people have so many difficulties with worms, bots, ticks, etc. Unfortunately, the narrow-spectrum “safer” pesticides are often prone to break down due to evolution of resistance in the target pest, whereas the legacy products of arsenic are dependable poisons.

Today most of the concern and press coverage on arsenic in the U.S. is about CCA (chromated copper arsenate) used to treat lumber for outdoor use, such as playgrounds. I had trouble understanding this, considering the seemingly small areas affected, and the fact that I didn’t eat playground equipment when I was a kid. However, as I point out below, CCA-treated wood accounts for about 60 percent of the import of arsenic to Florida. But the story is never simple.

In one instance in Broward County, after a playground was remediated by replacing soil and play equipment, the sampling extended into adjacent areas of bermudagrass turf maintained with MSMA, and, no surprise, there was arsenic there also.

Natural background levels of arsenic vary tremendously around the world. While the geometric mean of 441 near-pristine Florida soils was reported by the University of Florida’s Dr. Lena Q. Ma and co-

workers as 0.42 mg/kg (420 parts per billion), marl soils such as in Everglades National Park average around 5 mg/kg (around 500 ppb) which exceeds the Florida Department of Environmental Protection (FDEP) industrial soil cleanup goal of 370 ppb, and far exceeds the residential soil cleanup goal of 80 ppb.

No one is talking about excavating the Everglades to remediate a natural background arsenic level exceeding the environmental guidelines. Much of the arsenic there is tied up pretty well by calcite, organic matter, and with oxides of iron and aluminum, depending on pH, and the labyrinth of transformations that arsenic can undergo. The point is to use different background levels appropriate to different soils. And it is extremely difficult to predict what will cause arsenic to show up in drinking water.

Worldwide the major problem with arsenic involves entirely natural origins in well water used by people. The longest-term unintended experiments involving human consumption of arsenic in water have involved skin cancers in Taiwan, and internal cancers in Taiwan, Chile, and Argentina. Many of the natural sources exceeded 500 micrograms per liter (parts per billion), which is associated with approximately a 1-in-10 lifetime chance of internal cancer. The U.S. Environmental Protection Agency threshold is now 10 ppb.

The arsenic crisis in West Bengal and Bangladesh was due to the well-intentioned efforts to provide a safe drinking water supply, free from the problem of gastrointestinal microbes. For example, UNICEF and the Bangladesh Department of Public

# South Florida Grassing, Inc

Over 35 Years in the Grassing Industry

**TifSport**

**SeaIsle**  
Certified Seashore Paspalum

Sod, Sprigs and Rolls available

- We provide reworking and planting services for fairways, tees and greens
- Our grass is grown in Hobe Sound on gassed, sand-based soil
- Hydro-mulching, grassing and mulching also available

**Tifway 419**

**TifEagle**

**Tifdwarf**



**Certified Producer**

Southern Seed Certification Association

**(561) 546-4191**

[www.southfloridagrassing.com](http://www.southfloridagrassing.com)

**800-483-4279**

Health Engineering, and later private partners, installed some 3 million tube-wells, mostly since the 1980s, and unknowingly at the time the majority of wells were contaminated with horrendous levels of arsenic which was released from natural arsenic-bearing aquifer sediments.

Since the "safe" drinking water was not tested, the problem was discovered only in 1983 after people, eventually thousands, were diagnosed with arsenic poisoning symptoms, such as gross skin lesions in children. The confirmation of the problem as arsenic contamination of well water was confirmed in 1993, but by 1997, UNICEF was still patting itself on the back for surpassing its 2000 goal of "safe" drinking water.

Bangladesh is now grappling with the largest mass poisoning of a population in history, and if the estimated 200,000 victims of arsenicosis in West Bengal is any indication, the number affected in Bangladesh is far greater, based on 20 million people estimated to have been exposed. The British Geological Survey reported that among 9037 wells tested, 22 percent have arsenic concentrations above 100 micrograms per Liter (ppb).

Arsenic bioaccumulation by lowland plants and aquatic organisms contributes to elevated arsenic in some lowland soils, and bioaccumulation may also be a remedy. The University of Florida's Lena Q. Ma and coworkers showed in *Nature* magazine in 2002 that the brake fern *Pteris vittata* can accumulate up to 126-fold enrichment of arsenic, and the highest concentration was 22,630 ppm arsenic in the

plant. The fern naturally grows better in alkaline environments where arsenic is more available, and grows better in arsenic-contaminated soil than in uncontaminated soil.

As the FDEP's Leslie Smith pointed out on Nov. 14, turf fertilizer cannot be ignored as a possible source of elevated levels of arsenic in golf courses. But the most complete report that would shed light on arsenic in Florida is an extensive draft report, "Quantities of arsenic within the state of Florida, by University of Miami's Dr. Helena Solo-Gabriele and others such as UF's Dr. Timothy Townsend. The bottom line is that about 2500 metric tons of arsenic moved into Florida in the year 2000, 60 percent associated with CCA-treated wood, 20 percent from herbicides, 15 percent from geologic sources such as phosphate mining, and 4 percent from coal.

Although Florida has about 50,000 tons of "accessible" natural arsenic reservoirs, including geological reserves, roughly 50 percent is associated with CCA-treated wood, and between 7 and 20 percent is associated with arsenical pesticides. MSMA (and DSMA) were described as a "difficult dilemma since these chemicals are applied in liquid form directly on crops and golf courses. Contamination from these arsenical herbicides is immediate, quick to disperse, and thus difficult to control. Given these observations, efforts in Florida should focus on reducing the use of arsenical herbicides for controlling weed growth on crops and golf courses," and properly dealing with CCA-treated wood and wood waste.

*Reprinted with permission from the e-newsletter, "Turfgrass Management," Vol. 4, Number 22 and Vol. 5 Number 1. For information about the author, see inside cover.*

*Editor's Note: While not disputing the obvious import of arsenic into Florida soils through chemical use, I do question some of the figures in the report by Dr. Helena Solo-Gabriele referenced above. In the report they used the generally accepted figure of 150 acres per golf course times 1400 golf courses in Florida to estimate the amount of arsenic applied annually.*

*When you break down the actual acreage per golf course that might logically receive MSMA treatments combined with the fact that many of the 1400 courses don't treat wall to wall or even use MSMA other than some spot treatments, their figures need to be adjusted downward significantly. However, that factor does not relieve superintendents of the responsibility of reducing the use of a product whose final impact to the environment is under scrutiny.*

## ADA Guidelines Target Golf and Recreational Facilities

If your club is planning to expand or renovate its course or other facilities, you should be aware that the federal government has issued new ADA guidelines that specifically deal with golf courses and

# Neo-Tec<sup>®</sup>

## Effective, Environmentally Sound Nematode Control

It's true! **Neo-Tec** isn't like other nematicides. Neo-Tec is naturally derived and EPA registered under the safer pesticide rules. Neo-Tec is harmless to beneficial nematodes and is completely biodegradable.

- **Promotes healthier roots by controlling harmful nematodes**
- **Easy container disposal, minimal PPE and re-entry time**
- **Reduces damage to plant root in the following ways:**
  - ▶ **Disrupts the ability of plant parasitic nematodes to locate plant roots, thereby restricting penetration, feeding and reproduction**
  - ▶ **Reduced nematode feeding activity reduces the opportunities for invasion by pathogenic fungi**



**Parkway Research**

A Division of

**BRANDT**  
CONSOLIDATED  
Professional Agriculture

**800.442.9821**

info@parkwayresearch.com

Fax 281.590.3353

other recreational facilities.

The Americans With Disabilities Act (ADA) is a comprehensive federal law that prohibits discrimination on the basis of disability. Among its provisions is a requirement that places of public accommodation and commercial facilities be readily accessible to and usable by individuals with disabilities.

The ADA's public accommodations requirements went into effect in 1992. They generally require facilities that are open to or used by the public to be accessible and usable by individuals with disabilities by removing architectural and communications barriers and providing certain aids to assist individuals with disabilities.

Due to the unique features of recreational and outdoor facilities, in July 1993, an advisory committee was convened as the first step in developing new guidelines. After nearly 10 years in development, a final draft rule applying ADA to golf courses and recreation facilities was released Sept. 3. The Department of Justice is now incorporating the new guidelines into the existing ADA framework. When this process is complete, these new rules will carry the force of law.

As with the original provisions of the ADA, private clubs remain exempt. However, to the extent your club is open to nonmembers or the general public, you may be required to abide by the ADA. The regulations summarized below represent the final language awaiting adoption by the Department of Justice.

### Golf Courses Overview

- The proposed accessibility guidelines would apply to newly constructed or altered golf courses, driving ranges, practice putting greens, and practice teeing grounds.
- Special emphasis has been placed on the use of golf cars to make the game accessible to the majority of persons with disabilities. As a result, the proposed guidelines provide for access, via a golf car passage (a continuous passage on which a motorized golf car can operate), to various elements of the facility.
- Generally, accessible routes must be located within the boundary of the golf course, must be 48 inches wide (minimum) and connect to the bag drop areas, accessible teeing grounds, and putting greens. Additionally, where handrails are required, the accessible route must be at least 60 inches wide.
- All of the amenities (such as snack bars, toilet rooms, and weather shelters) on a golf course must be accessible and connected by a golf car passage.

### Driving Ranges and Practice Tees

- Where practice tees or driving ranges are provided, at least 5 percent of the practice tees, but not less than one tee, must have a minimum clear area of 10 feet by 10 feet with a practice slope which does not exceed 1:48 in all directions.
- The area must be constructed so that a golf car can enter in a forward motion and maneuver.

### Golf Car Passages

- Golf car passages must be at least 48 inches wide. This dimension is based on the standard width of most golf cars.
- Where curbs or other manmade barriers exist, openings of at least 60 inches wide, at intervals of 75 yards, must be provided for access to fair ways by golf cars.

### Putting Greens

- Putting greens must be designed and constructed to allow a golf car to enter, maneuver within, and exit the putting green.

### Weather Shelters

- Weather shelters must be designed and constructed to allow a golf car to enter and exit in a forward direction.
- A clear floor or ground space of 60 inches by 96 inches (minimum) is required to allow a golf car to be driven directly into a shelter and exit in a forward motion.

### Boating Gangways

- The proposed guidelines note the difficulties for gangways (bridges that link land or fixed structures with floating piers) due to changing water levels that may affect the slope of such structures. As a result, transition plates at the top and bottom of gangways are permitted to provide for level landings. Transition plates should have slopes that are less than or equal to 1:12. Several techni-

## Thinking about remodeling your course?

One call can save you Time,  
Money and Aggravation.



*The Links at Madison Green  
Royal Palm Beach, FL*



John Sanford Jr.,  
*Member of*  
American Society of  
Golf Course Architects

*The Links at  
Madison Green*

*Voted "Florida's Best  
New Public Golf  
Course" by Florida  
Golf Magazine*



**SANFORD  
GOLF DESIGN**

866.233.4SGD

**DESIGN • REMODELING • CONSTRUCTION MANAGEMENT**

[www.sanford-golf.com](http://www.sanford-golf.com)

cal exceptions are provided to this requirement.

### Slips

- Where boat slips are provided, the guidelines require at least 3 percent of the slips, but not less than one slip total, comply with accessibility requirements. The new regulations provide a chart listing the number of slips required by total marina size.

Persons with disabilities should have access to different types of boat slips; however, the slips may be clustered in the same area.

### Swimming Pools, Pool Entry, and Exit

- Swimming pools must have at least two means of entry and exit. A sloped entry or lift must be the primary means of access for swimmers requiring access accommodations. The secondary access can duplicate the primary means of access and may include transfer walls, transfer systems, or stairs.
- Swimming pools with less than 300 linear feet of swimming pool wall may (as an exception) have only one means of access, but that means of access must be either a lift or sloped entry.

### Shooting Ranges

- Shooting facilities: Where fixed firing positions are provided at a site, at least 5 percent, but not less than one of each type of firing position must be accessible.
- Fixed firing positions: Fixed firing positions must contain a 60-inch-diameter space and have a slope no steeper than 1:48.

*Editor's Note: This is a condensation of the some of the major points affecting golf courses and recreational facilities. The Americans with Disabilities Act - What you need to know" will be the subject of the GCSAA "Current Issues in Golf" program at the Atlanta Conference and Show on Thursday, Feb. 13 from 2:30-4:30 p.m.*

### On the Web

The complete long version of the Federal ADA document can be viewed at the GCSAA website by registered GCSAA members only. Go to [www.gcsaa.org](http://www.gcsaa.org). Select Government Relations and follow the links.

## Current News & Issues...

### Curfew Granted 24C Label

According to a recent e-mail from Dr. Brian Unruh of the UF/IFAS West Florida REC in Milton, Dow Agro Science's two-year wait has ended as the United States EPA has granted a 24C label for the soil fumigant Curfew. After being limited to applying Curfew to only 5,000 acres per year for the past two years under an experimental use permit, Dow will now be able to take orders for treatment of more fairway acreage.

Curfew, a soil fumigant, has been used in agriculture for years. Two years ago, Dow conducted test applications on several Florida golf courses in cooperation with the Florida DEP. The state signed off on the use of Curfew for nematode control on golf course fairways, but the U.S. EPA was not able to come to a decision within the mandated 90-day review period and the decision-making process has dragged on for two years. Dow enlisted the aid of superintendents to write their legislators about the importance of this product as a potential alternative to Nema-cur, which is being phased out after the federal agency seemed to ignore the state of Florida's acceptance of the product's use on golf courses.

Those courses that were able to book fairway treatments the last two years reported excellent results and turf response especially during the tough drought-induced growing conditions.

### Arsenical Herbicides Under Review

Herbicides containing arsenic compounds are the subject of a statewide task force in Florida. The action is the result of the arsenic levels found in soil and water samples taken from golf courses in South Florida. The issue was moved to the front burner when land sales transactions were put in limbo as the arsenic levels found in samples during routine environmental audits were found to exceed health-concern levels.

Arsenic is a naturally occurring element and can be found in most any soil and water sample in Florida, which complicates the regulatory process. In fact natural background levels can exceed the regulatory level being proposed by Florida Department of Environmental Protection.

Right now, applying arsenical herbicides is an obvious, easy target source of arsenic being added to the environment. But, arsenic is also found in fertilizers (it combines readily with phosphorus), bio-solids, mulch, native soil and fill dirt, native limestone marl and rock formations, and waste water. Determining the source of the arsenic in a sample is nearly impossible. When the sample is processed, the elemental arsenic remains with all the attached molecules having been removed in the process.

Stakeholders will meet in Tallahassee in late January to discuss the use of arsenical herbicides. An informal survey of courses concerning the current use patterns of the commonly used grassy weed herbicide, Monosodium methanearsonate (MSMA) revealed the modern trend of spot treatment versus wall-to-wall spraying common 20-30 years ago. With the advent of best management practices, and integrated pest management principles, and new chemistry - the overall use of MSMA has declined considerably. Tropical signalgrass is cited as the weed still requiring use of MSMA for good control. Other products including pre-emergent herbicides are available as alternatives to using MSMA to control the goosegrass, crabgrass and the various sedges.

Protecting the environment should be our number-one concern, but as in any regulatory action, science should play the major part in determining the facts of risk and exposure and environmental harm. Arsenic is known to the general public as a poison and recently was the focus of a controversy concerning the treated-lumber industry. While no medical evidence showed a real health concern, the public perception of arsenic and the political nature of the issue forced the industry to change to a different preservative. The same situation is very possible for turf applications of arsenical herbicides.

Superintendents should take a serious look at their weed-control programs and determine what role MSMA plays in the conditioning of the course. We may be forced to reduce or eliminate the use of MSMA entirely. A cost analysis should be done to include pre-emergent weed control products and other more expensive, but effective post-emergent products.

See the related article on MSMA by Dr. Phil Busey in this section. Dr. Busey gives a detailed account of the history of MSMA use in two excerpts from his e-newsletter, "Turfgrass Management." The complexity of the issue is evident from the commentary in the article.

### Water Restrictions - The New Way Of Life

It is probably safe to say the drought in Florida is over. Polk County posted a new total annual rainfall record set in 1948; the new record for 2002 is 72 inches. Regardless of lake levels returning to near normal and more frequent rains helping with turf irrigation, the growth and development of Florida rushes on and that prospect has water-management districts for the most part, keeping watering restrictions in place.

Out of necessity there has been increased cooperation between the golf industry and the water management districts, but the five water-management bodies still approach their sovereignty with their own particular viewpoints and must be dealt with independently.

At the last check South Florida, Suwannee and Northwest Florida districts had no watering restrictions. Southwest Florida still has two-day-per-week restrictions in general, but has acknowledged golf courses' need for flexibility in turf management and has relaxed scheduling constraints, but cautions users and permit holders to stay within permitted amounts and follow best management and conservation practices.

The St. Johns River district is still on twice-per-week watering with the usual prescribed exceptions for hand-watering hotspots, overseeding, frost and wilt protection and new turf establishment.

The bottom line is we need to continue our involvement with the various districts so they know we are trying to do the best we can in managing water resources for golf which is a major contributor to the state's economy.

*Compiled by Joel Jackson, CGCS*