stresses, at different times of the year are quite likely to react in different ways. One of these different reactions will undoubtedly be with hormone regulation, and this is consistent with the variability in plant response to hormone applications in research results and field trials across the country. There currently is no evidence to suggest that applications of plant hormones will yield favorable or consistent results with respect to improved plant health. Furthermore, adding hormones to plants beyond normal levels may produce an inhibitory or undesirable effect. Without research information to identify and quantify treatment regimes, it may be wise to avoid tampering with plant hormonal activity (7). Anecdotal evidence and testimonials have been the substitute for independent research results repeated at multiple locations.


Another type of growth stimulant available on the market contains humate or humic acid. These are naturally occurring organic compounds that are the end products of biological decomposition. Accordingly, they are extremely resistant to further decomposition. Products containing humates claim to increase cation exchange capacity, increase microbial activity, and chelate micronutrients. Kussow reviewed manufacturer recommendations for amending a sandpeat root zone mix with humate and found it to be a very expensive means of increasing the CEC by 13% (9). His re-

Evaluating Independent Research

- Who (principal investigator) did the research?
- Where was the work done (lab or field, sand or soil)?
- Look for replication, good comparative treatments, and statistically significant differences.
- Have the results been duplicated at another site by another independent researcher?
- Have the results been published in a refereed journal?
- Slick brochures can be confusing! Don't be fooled by sales techniques.

view further concluded that iron, copper, manganese, and zinc are rarely deficient in turfgrass soils, thus enhancing micronutrient availability may only provide negligible benefits. Another study clearly demonstrated that since humates are the end result of decomposition and thus resistant to further breakdown, they do not stimulate increased microbial activity (25). Yet another study reviewed

the effects of six non-traditional growth-promoting products on the establishment of creeping bentgrass in high-sand-content rootzones. Only one of the products produced significant differences from the control, and the product contained humate. Upon chemical nutrient analysis of the product, however, it was discovered to contain 6% N, 5% P, 2% K, 4% S, and 4% Fe. Using this product at





the recommended application rate was equivalent to applying an additional 0.75 pound N, 1.3 pounds of P, and 0.34 pound of K per 1,000 square feet per month (7). It may well be that this response could have been duplicated with conventional fertilizer, and it would seem to request an independent nutrient analysis of any growth stimulating products you intend to try.

Finally, there have been studies that indicate humates and humic acids can reduce the efficacy of pesticides by reducing their absorption by plants and pathogens (9). It is also reported that the fulvic acid component of humates can actually increase the solubility of pesticides and possibly increase mobility (2S). Most of the studies that claim any benefit from adding humates were in either nutrient culture or sand culture systems, not in field situations. The variation in humic substances from different sources and lack of research that supports their use on turfgrasses currently do not jus-

tify their use in turf management.

Carbohydrate fertilizers, another biostimulant, have not been proven to improve turfgrass stress tolerance or have any lasting impact on soil microbial populations. Again, research on turfgrass and carbohydrate application is lacking, but observations across the country indicate no observable benefits. Any stimulation of microbial activity is likely to be very short-lived.

Microbial Inoculants

Various microbial inoculants have been formulated for use on turfgrass, with claims of accelerated organic matter decomposition, improved nutrient use efficiency and availability, soil conditioning, disease control, mycorrhizal associations, and others. The success of these inoculants has been limited for a number of reasons. At this point, you should be aware that the microbial community is a very diverse and complex set of organisms The degree of natural com-

petition, antagonism and predation limits the successful establishment of introduced species. Persistence of applied organisms is further hindered by the continual temporal and spatial fluctuation of microorganism populations (6). Formulation and delivery of the organisms present even more problems for microbial inoculation (15). If the organisms can be kept alive until application, many are sensitive to UV light and must be applied frequently (in some cases nightly) to establish sufficient populations. Although there have been efforts to apply microorganisms through irrigation systems, the results remain largely inconsistent (2). Finally, some companies will not even list what organisms they have formulated, because they are proprietary. Without knowing what is being applied, it is impossible to gauge the potential benefits. These organisms could be detrimental to your turf by actually competing with the beneficial organisms already present in your soil (7).

Producers of Quality Turfgrasses for Golf Courses and Athletic Fields



Pike Creek Turf inc.
Route 2, Box 376-A, Adel, Ga. 31620
1-800-232-7453

We grow & install high quality sod and sprigs:

- ♦ Tifway
- ♦ TifSport
- ♦ TifEagle
- **♦** Tifdwarf
- ♦ Centipede Sod/Seed
- ♦ Meyer Zoysia

Producers of Fumigated Georgia Certified Bermudagrasses

Rowplanting Services Available

Composts

With little doubt, the most promising method of managing and enhancing the activity of soil microbes is with composted organic matter in wastes and other materials. Ironically, this is also one of the oldest agricultural practices. Composts have been shown to add an active microbial component to soils and to stimulate those microbes already present in the soil (14). Well decomposed organic matter provides excellent habitat and energy sources for soil microbes, and will provide more permanent benefit than inoculation with microorganisms. Composts will effectively enhance soil aggregation, provide nutrients, reduce compaction, and improve soil porosity. Sandy soils amended with compost will exhibit greater nutrientand water-holding capacity (10). While limited evidence exists, there is some data to suggest amending sand-based rootzones with compost can offer improved establishment and disease control over commonly used peat amendments (5, 14).

The use of composts in turfgrass management presents a viable means of recycling municipal and industrial wastes while improving turfrass quality. Composts can vary considerably, however, depending on their source. Commonly used composts include brewery sludge,

On-Site Testing Protocol

- Test products at several locations representing different conditions on the golf course.
- Use controls (no product) to establish comparisons.
- At least two years of field data are necessary to obtain an accurate assessment.
- Rate the plots monthly to track differences (color, disease, stress tolerance, rooting, etc.).
- Conduct an independent nutrient analysis of new products. You may be seeing a fertilizer response!
- Be honest! Is it the product or favorable weather, better cultivation, an improved growing environment, or other changes in management strategies?

yard wastes, poultry litter, animal manure, municipal wastes, and food wastes. It is recommended to have composts tested for organic matter content, ash content (especially if used as a topdressing), moisture content, pH, nutrients metals, and soluble salts (10). On-site composting operations should follow guidelines to ensure that the material has been properly and sufficiently

composted (14, 20, 28). The diseasesuppressive characteristics of composts will be discussed in the next section.

Biological Pest Control

In recent years, considerable focus has been placed on the biological suppression or control of various turfgrass pests, including diseases, insects, and weeds. Reducing the pesticide load on

If You're Looking for Improved Pest Control . . . Du Cor has an Easier Solution!

New Du Cor LEMWET can help improve your pesticide spray application! LEMWET is a lemon extract attractant and surfactant for use with most pesticides. LEMWET attracts nearby pests to your spray. In addition, it's non-ionic surfactants improve spray penetration and bring your pesticide into close contact with pests!

For over 35 years, Du Cor has been coming up with products to fit our customers needs. From liquid micronutrients, humates and fertilizers, to specialty dyes, surfactants, odor counteractants and more, we take pride in finding special solutions that help our clients.

So if your looking for something special to attract a nuisance, or maybe just a different solution to help you grow, give Du Cor a call. We may already have just the solution you need!



Du Cor International Corporation

P.O. Box 593298, Orlando, Florida 32859 407-859-4390 or Toll Free 1-800-382-6735



the environment is the primary impetus behind such study. While research has proven effective pest control with various biological entities in the laboratory, few have proven consistently effective in field studies.

Biological control operates on five basic interactions with the turfgrass-soil community: competition, antagonism, predation, parasitism, and pathogenicity (1). The two ways of exploiting these interactions include microbial inoculants and organic amendments. While dozens of organisms with potential as inoculants for disease control have been studied (17, 18, 24) few have demonstrated any efficacy in the field, and only one product (Biotrek 22G, (Trichoderma harzianum) has been registered for disease control on turf (11, 15). Biological control of insects has been somewhat successful in recent years with such organisms as entomogenous nematodes, soil bacteria and fungi although registered products are still limited (21, 26).

Serious shortcomings exist in the understanding of the pest control mechanisms themselves, relationships with other organisms in the community, and formulation and delivery technology. Furthermore, foliar disease control with inoculants is limited due to UV sensitivity of the organisms and wide fluctuations of environmental parameters in the turfgrass canopy.

The difficulty in delivering organisms to the roots has preempted much success in controlling root diseases. Because successful pest control typically depends on the establishment of high population levels, frequent (and arguably unsustainable) applications become necessary. Injecting organisms through irrigation systems has yet to be proven as an effective method of uniform and consistent microorganism application

Keep in mind that 1) population interactions within the soil are dynamic and interrelated, 2) introduced organisms are slow to colonize habitat and generally fail to persist, and 3) it is unclear whether the introduction of microbes in the environment will produce a lasting change and if the introduction will be beneficial in the long run (1, 15).

Organic soil amendments and additives, particularly compost, have perhaps a greater potential for effective biological control of diseases than do inoculants. Well-composted material (2-3 years) often exhibits disease-suppressive characteristics (14).

Studies at Cornell University have demonstrated significant and lasting disease suppression of Pythium root rot, dollar spot, and snow mold when composts were used as amendments or topdressing (14).

Continued research in this area to reveal the microbiological mysteries should help develop more reliable and predictable composts for disease suppression and soil conditioning. As alluded to earlier, proper composting techniques and laboratory testing coupled



Soft Swipe

Created especially for cleaning non-metal spikes
For all courses where non-metal spikes are used!



The aluminum base of the SoftSwipe Cleaner uses 4 prongs to penetrate the ground. Place it wherever non-metal spikes need to be cleaned.

PLAY IT SAFE. SWIPE YOUR SPIKES!






The Exclusive Distributor for the SoftSwipe Turf/Tee Spike Cleaner

Swipe away debris in seconds. Move the SoftSwipe with the tee markers for player convenience.

Will not damage any non-metal spike and extends spike life expectancy



Only \$29.95

Includes aluminum base and replaceable SoftSwipe cleaner disk

Available from any LESCO Stores-on-Wheels, LESCO Professional Golf Representative, or call 1-800-321-5325

Order today!

LESCO is a registered trademark of Lesco, Inc.
Stores-on-Wheels is a registered service mark of Lesco, Inc.
SoftSwipe Cleaner Disk (patent pending)



with on-site testing will reveal what to expect from composts.

New Products

Never before has the turfgrass industry had as many commercially available products for use. Financial responsibility and sound management dictate that product purchasing decisions are of extreme importance. So how does one choose between the good, the bad, and the ugly?

The first place to start is with the product label. There are products that have been registered with the EPA and can legally justify the claims of the product. These are products that contain active ingredients (29). There are unregistered products marketed for various uses, some of which are supported by inde-

pendent research. Then there are products marketed for various uses without supportive research. These products use testimonials and fancy marketing to make a sale, and often can be classified as snake oils.

Let's be sure we understand the independent, scientific research that supports product use. Be sure you know who conducted the research, where, under what conditions, and the relevancy to turfgrass systems. Also, look for replication in the study, good comparative treatments, and least significant differences. Check closely to see that the results have been duplicated at another site by another independent researcher, and that results have been published in a refereed journal. Make no mistake, slick brochures and displays can be confusing!

One product advertisement I recently reviewed claimed the product would cause no grow-in layer, extend the useful life of greens, reduce grow-in time, eliminate the possibility of nitrite (yes, they said nitrite, not nitrate!) and phosphate leaching, and reduce labor, among other things. This company may need legal counsel as much as scientific counsel. Finally, call the researchers and ask technical representatives what the active ingredients are and what are their modes of action (29). University extension personnel and USGA agronomists can also provide valuable information.

If a product you are interested in passes this initial screening, it is strongly recommended to conduct on site testing at your golf course. Many of these products are not cheap, and good manage-

Literature Cited

- 1. Berndt, W. L. 1996. Population interactions: the key to unlocking natural pest control. *The Turf Letter*. 1(3):1-8.
- 2. Bresnahan, J. 1998. Bioject: results of 1997 research trials. Presented at New
- England Regional Turfgrass Conference and Show Providence, Rl. March 4,1998.
- 3. Chiariello, N., J. C. Hickman, and H. A. Mooney. 1982. Endomycorrhizal role for interspecific transfer of phosphorus in a community of annual plants. *Science* 3:941-943
- 4. Dernoeden, P. H. 1987. Management of take-all patch of creeping bentgrass with nitrogen, sulfur, and phenylmercury acetate. *Plant Disease*. 71 (3):226-229.
- 5. Hall, R 1996. Composts vs. peats: composts win. *Landscape Management*.35(8):14.
- 6 Holl, F.B. 1997. The mysterious world of the turfgrass root zone. *Turfgrass Trends*. 6(11):1-8.
- 7. Isaac, S. 1998. Bio-stimulants and all that! *International Turfgrass Bulletin*. 199:9-10.
- 8. Karnok, K. J. 1989. The use of plant hormones as non-nutritional turfgrass growth enhancers. *Golf Course Management*. 57(7):28-38.
- 9. Kussow, W. R 1994. Humate and humic acid. *The Grass Roots*. 22(2):18.
- 10. Landschoot, P. 1998. Using composts to improve turf performance. *Turfgrass Management in the Pacific Northwest*. 1(1):32-34.

- 11. Lo, C. T., E. S. Nelson, and G. E. Harman. 1996. Biological control of turfgrass diseases with a rhizosphere competent strain of Trichoderma harzianum. *Plant Disease*. 80(7): 736741.
- 12. Loomis, R S., and D. J. Connor. 1992. *Crop ecology*. Cambridge University Press, New York, NY.
- 13. Moore, J. F. 1997. Let's give credit where credit is due. *USGA Green Section Record*. 35(2):20.
- 14. Nelson, E. s. 1996. Enhancing turfgrass disease control with organic amendments. *Turfgrass Trends*. 5(6):1-15.
- 15. Nelson, E. B. 1997. Biological control of turfgrass diseases. *Golf Course Management*. 65(7):60-69.
- 16. Nelson, E. B. 1998. Turfgrass pesticides and biological disease control: are they compatible? *CUTT* 8(4):7-9.
- 17. Nelson, E. B., and C. M. Craft. 1991. Introduction and establishment of strains of Enterobacter cloacae in golf course turf for the biological control of dollar spot. *Plant Disease*. 75:510-514.
- 18. Powell, J. F, M. G. Nair, and J. M. Vargas, Jr. 1994. Utilization of a bacterial metabolite for the management of dollar spot disease of creeping bentgrass (Agrostis palustris). *Phytopathology*. 84(10):1077.
- 19. Roberts, E. C. 1998. Turf research: nature, needs and net results. *Turfgrass Trends*. 7(1):i-9.
- 20. Senseman, R. Y. 1995. Composting: turn your eyesore into black gold. *USGA Green Section Record*. 33(3):8.

- 21. Shetlar, D. J. 1996. Field testing of biological pesticides. Turfgrass Trends. 5(8):1-9.
- 22. Stockwell, C. T., E. B. Nelson, and G. M. Craft. 1994. Biological control of Pythium graminicola and other soil borne pathogens of turfgrass with actinomycetes from composts. *Phytopathology*. 84-1113.
- 23. Thompson, D. C., B. B. Clarke, J. R. Heckman, and I. A. Murphy. 1994. Suppression of summer patch in turf with acidifying nitrogen fertility programs. *Phytopathology*. 84(10):1085.
- 24. Thompson, D. C., B. B. Clarke, and D. Y. Kobayashi. 1996. Evaluation of bacterial antagonists for reduction of summer patch symptoms in Kentucky bluegrass. *Plant Disease*. 80(8):856-862.
- 25. Varshovi, A. 1996. Humates and their turfgrass applications. *Golf Course Management*. 64(8):53-56.
- 26. Vilani, M. G.1996. Biorational control agents for Japanese beetle management. *CUTT 7*(1):1-7.
- 27. Wegner, T. 1997. Bentgrass response to non-traditional soil additives. *The Grass Roots*. 25(4):58-61.
- 28. Zontek, S.1974. Composting alchemy in action. *USGA Green Section Record*. 12(3):9-11.
- 29. Zontek, S. 1997. Buyer . . . beware. USGA Green Section Record. 35(1):17.

ment involves an economic analysis. Test the material at several locations on the golf course representative of different conditions, replicate (meaning include repeated treatments at each site), and use untreated controls and other treatments in side-by-side comparisons. All too often, new products are tried all over the golf course without a control; thus, it is impossible to determine what effect, if any, the new product has. Perceived benefits could be a result of favorable weather or other management techniques (7, 13). Take consistent, monthly ratings of the plots for color, disease, and rooting depth and mass, and note stress tolerance differences. Good tests require at least two years of field data. Because a product will cause no harm is not reason to use it, and such a decision is representative of poor management.

Conclusion

Turfgrass management is a continually evolving science, and as our understanding of the microbial community in turfgrass systems improves, new products will routinely hit the market. Some of these products will be useful, and many others will not. Independent reseach will be essential to the development of effective products. Perhaps companies marketing biological products would be wiser to fund some research than to purchase full page ads in popular trade magazines (if they have faith in their products)!


If completely organic management is ever realized, it will certainly be through a gradual phase-out of synthetic products. Along with the advent of biological products, golf course superintendents must also keep themselves apprised of advances in synthetic chemistry. Many new products have been developed from synthesized organic compounds that are effective at very low levels of active ingredients, have low water solubility, short half lives, and a strong binding potential with soil and organic matter. The new synthetic chemistries are better for the

environment than many of the older chemistries.

The importance of a strong microbial community cannot be questioned. The effectiveness of various products available to stimulate microbial activity can be questioned. Become familiar with soil microbiology and processes, check for duplicated independent research to support product claims, and test the material yourself to be sure it is effective and makes good economic sense. But whatever you do, don't forget the basic tenets of successful turfgrass agronomy: adequate sunlight, drainage, air circulation, proper fertility, good water management, traffic control, and cultivation.

CREDIT: USGA GREEN SECTION RECORD, **JULY-AUGUST 1998**

MATT NELSON is an agronomist in the Northeast Region of the USGA Green Section. He "bugs" superintendents to take a close look at product purchasing.



Safety Storage, Inc. prefabricated, relocatable buildings provide a low-cost solution to safe storage, containment, mixing, and dispensing of golf course chemicals and hazardous materials.

Safeguard personnel, avoid the liability arising from soil and groundwater contamination, meet fire safety needs, and achieve full compliance with federal, state, and local regulations.

Building sizes range from 5' to 32' in length with capacities up to 320 sq. ft. Standard features include continuously-welded, heavy-gauge steel construction, secondary containment sump, removeable fiberglass floor grating, and chemical resistant coating inside and out. Select from a full range of options

For complete details, call us toll free at: 1-800-344-6539 • www.safetystorage.com

...professional hazardous materials storage solutions



Petro-Chem Environmental Systems, Inc. 15310 Amberly Drive, Suite 250 Tampa, Florida 33647 800-330-6949 Fax: 813-972-0955



Microbes: The Real Millennium Bugs

icrobes! Those little microscopic links in the chain of life. We all know that they're out there in the soil, water, air and in our bodies. What we don't know exactly is how they really work, what affects their populations the most, and how we can harness them to maximize turfgrass management.

Just like many of you, I have anecdotal experience that would suggest that this product or that product did indeed produce some sort of positive results. The challenge comes in finding reams of peer-reviewed scientific studies that give us a better idea of what's going on and how predictable a product or its results might be. We may soon find ourselves on a path that separates the snake oil from salvation.

Synthetic chemicals and fertilizers are under constant attack these days. They have been quick, effective tools to solve turf problems for about 50 years. Now the politically correct and environmentally conscious activists are waging a holy war to eliminate as many pesticides and fertilizers whether or not they are truly harmful. We can fight back, but it is a constant battle and the end users are at the mercy of too many things they can't control like the media, and politics and the manufacturers' bottom lines.

So, what if we could find more natural, benign ways of producing strong, healthy turf in a very active, healthy, growing medium? Maybe we can reduce our dependence on synthetic products, which are likely to start disappearing anyhow. Maybe we can find less expensive ways to enhance the natural systems at work and eventually produce stronger, healthier playing surfaces. The following articles provide information, insight, caution and hope for microbial solutions, and we are poised to start funding more research on these millennium bugs.

Former Sales Rep is Sold on the Products, Not the Sales Pitch

Since I tried to sell this technology to most of you for two years, I felt obligated to respond to this Hands On topic, "Microbes – The Real Millennium Bugs." I'm not going to waste my time or yours continuing to sell you on this concept. By now you either believe that using microbial products is good science or you think it is a deception.

Now as a superintendent once again, I use the Green Releaf products that I sold, but not exactly in the way I sold them. I tried to sell programs that incorporated small amounts of all the products that Green Releaf sold on a 7- to 14-

day schedule. Green Releaf trained us to sell the products that way, because if you bought into the program you would not have room to use other people's products. The program I use incorporates several different brands of products.

I basically spray my greens every week with a mixture of nutrients. I determine which nutrients to use and the amount of nutrient needed by tissue sampling. In that mixture I incorporate two gallons of Green Releaf 5-10-5 plasma per acre. I use the plasma because of the carbohydrates, amino acids, and humates that are in the product. The nutrients in the plasma are just a bonus.

The theory is that if you increase the amount of food source for the microbes in the soil, you will increase their population. The more microbes that are in the

soil the more efficient the plants will become.

I'm pretty sure that all the so-called biological products are based on this theory. In my mixture, I add one gallon of fulvic acid per acre to help hold the nutrients in the soil. My own personal theory is that by using plasma and fulvic acid, I can get by with using the lower-priced liquid nutrients and get the same results as if I used the high-priced ones.

When I was selling microbes for Green Releaf, most microbiologist were skeptical of the validity of the products. Every researcher I spoke with said that, in a laboratory setting ,the microbes did all the wonderful things that Green Releaf claimed. However, only a few scientist felt the microbes could live long enough to get down into the root zone.

Even the most discriminating skeptics felt that if you could get the microbes into the root zone they would be notably beneficial. Because of this, I use the living microbes in two ways. I use them when I aerify, and I inject them with a hydroject.

When I aerify, I spray 5 gallons of Bio A and Bio B+ per acre after the cores are removed and before I topdress. I immediately flush the "bugs" into the soil with 10 to 15 minutes of irrigation. I then add 300 lb. per acre of Green Releaf 15-4-7 organic granular (to feed the microbes), then drag it into the holes with the topdressing.

What I see by doing this process is rapid recovery from the aerification, and a flush of root growth even in the summer months.

During the winter I inject microbes into the greens monthly. Last May going into the rainy season, the roots on my greens were 6 to 8 inches deep and dense (I'm not just blowing smoke, several people can attest to this). During the heaviest part of the rainy season the roots shrunk to 2 to 3 inches, but new root growth was always occurring.

So here is the question: Can I attribute the success that I have had to microbial products? I don't know! It could be the microbes and carbohydrates. Or it could be that the weather was good last year, or my water management was

choice for GREENS

AVAILABLE FOR <u>SAFE DAMAGE FREE</u> APPLICATION ON YOUR GREENS, TEE TOPS AND SLOPES!

APPLICATION IS PROVIDED BY

Prosource OneTM (Formerly TERRA)



Not affected by weather * No colters or knives used
 Multiple products can be injected * No disruption of play

Application bookings can be made through your local **ProSource One**™ Salesman or directly to the turf application specialist @ 1-813 240 8607 or 1 813 762 1177 Ext. 260.



better this year, or my nutrient management is better than when I used granular fertilizers.

I honestly don't think there is any way to tell for sure if the microbes and carbohydrates have made a difference without conducting a lot of expensive research. However, I am not willing at this time to take the microbial products out of the program. Not until someone proves to me that using microbial products is not good science.

Do I think there should be money laid aside to research microbial products? Absolutely! In my opinion, with all the radical environmental groups trying to get us shut down, anything we can do that is environmentally responsible is worth researching..

MIKE HAMILTON, CGCS Foxfire C. C., Naples

Editor's Note: At its fall board meeting the FGCSA agreed to submit a proposal from Dr. Monica Elliott on microbes to the GCSAA for research funding. The purpose of the project will be to determine if Bacillus microbes applied to bermudagrass putting greens do survive and establish on either the turf or in the soil. The project will also determine the effect of frequent, multiple applications on the Bacillus populations. The research will be conducted on a portion of the FGCSA's Otto Schmeisser Research Green at the University of Florida's Research and Education Center in Ft. Lauderdale

Microbes Play Varied Roles in All Facets of Human Existence

If champagne, a loaf of freshly baked bread and a fragrant cheese will figure prominently in your Year 2000 celebration, then you can thank your local microbe. Human beings are dependent on microbes for many foods such as yogurt, sauerkraut, sour cream, beer and other fermented drinks.

There is also a type of bacterium that lives in the human intestine that makes vitamin K, a factor needed to clot blood.

In addition we take advantage of bacteria for their ability to protect us from microbial invaders. Human beings have their own special bacteria that have adapted to life in a salty, ocean of sweat on our skin. These bacteria keep out pathogenic fungi and bacteria. We also have a host of bacteria living in our throats, noses, intestines and other areas of the body. These friendly bacteria are tremendously effective at keeping the disease-causing bacteria away from their homes, our bodies!

A diversity of bacteria that associate with plant roots serve the same purpose of holding pathogenic microbes at bay.

Biological control agents, such as bacteria and fungi, work through a variety of mechanisms. These mechanisms include the production of antibiotics and other inhibitors; simple competition for food, water and space; stimulation of the plant's own system of defenses; and an environment around the plant roots that encourages growth of beneficial microbes.

Beneficial bacteria and fungi can form a barrier against pathogen invasion of plant roots. Because turfgrass's roots are surrounded by a host of friendly bacteria, it is much more difficult for *Gaeumannomyces* (take all disease organism) to reach its intended target.

In laboratory testing, conclusive tests have shown that *Gaeumannomyces* is sensitive to substances produced by several different species of bacteria. When a diverse group of microbes exists in the soil in the plant root zone, many reciprocal benefits are derived from that relationship. When bacteria and fungicolonize the rhizosphere, they are fed a steady diet of plant sugars. The plants in turn benefit from colonization of non-pathogenic microbes.

What happens if friendly plant bacteria are not available to protect plants? Soil that has been treated with harsh chemicals may deplete soil of the beneficial bacteria and fungi. When fungi and bacteria are no longer available as a shield pathogenic microbes may then be able to invade plants.

Within the past 50 years, the public has been taught that the only good microbe is a dead microbe. Nothing could be further from the truth. Without beneficial microbes, our plants would die and our own existence would be in jeopardy. And yet we

continue to douse microbes with bactericides and fungicides in an effort to exterminate the pathogens, but in doing so many beneficial are also depleted in numbers.

Some helpful bacterial species include *Pseudomonas*, *Bacillus*, *Cellumonas*, *Corynebacterium*, *Rhodococcus*, and other member of the *pseudomonads actinomycetes*. Some species of *Pseudomonas* help break down urea-based fertilizers enzymatically and convert urea to ammonia. Ammonia is then converted by nitrifying soil bacteria to nitrates, a form of nitrogen readily available for plant utilization

Certain species of *Bacillus* produce insect toxins as well as antibiotics that inhibit fungal growth. *Bacillus* is an extremely hardy microbe. These bacteria form endospores under adverse conditions, enabling the microbe to withstand drought, high heat and adverse pH conditions. *Cellumonas* is extremely helpful in producing substances that break down dead plant material in the soil, thus helping remove thatch buildup. *Rhodococcus, Bacillus* and *Pseudomonas* are proven pesticide and herbicide degraders; therefore much of the excess biodegradable pesticides are devoured by microbes.

One important factor to remember is that just one bacterial type cannot do a job alone.

This is the reason why it is important to maintain a highly diverse population of microbes. What one bacterium can break down in the soil may not be an available food source for another microbe. As mentioned above, several different bacteria are needed in the first step of converting urea to ammonia. Without a team of ureadegrading organisms, the nitrifiers would be powerless to use urea-based fertilizers as a nutrient and thus produce nitrates for plants.

Without many different bacteria and fungi in the soil, a healthy environment cannot be maintained. In the decomposition of dead plants and animals, certain inorganic elements such as phosphates are released and made available for plant nutrition. Many different bacteria utilize common organic wastes and produce carbon dioxide and water.

Without microbial diversity, nutrient