

Organic-Gro™

On top of the leader board for Organics!

Organic-Gro™ PRO 20-2-6 with UFLEXX™ works as nature intended by supplying nutrients to your turf while restoring proper microbial and earthworm activity for more colorful, healthier and stronger fairways and greens.

Organic-Gro™ PRO is a special blend designed specifically for golf courses. It contains 80% all-natural organic fertilizer and 20% UFLEXX™, a slow-release nitrogen. For greens, ask for SGN90, for fairways and rough, ask for SGN150.

To order call (866) 815-4839 or visit www.organic-gro.com

- Safe for golfers, wildlife and the environment
- Perfect for use by rivers, lakes, ponds and oceans—right to the water line!
- Lasts up to 10 weeks—reduces number of applications and labor
- Won't burn—even when applied in extreme heat
- 80 percent insoluble and non-leaching—the nutrients stay where you put them
- One application provides 18.75% slowly available nitrogen per 1,000 sq. ft. when applied as directed
- Available in 50 lb. bags, 40 bags/pallet. Also 1 ton bulk sack.



Regenerated Resources™

©2004 Regenerated Resources LLC


Visit us at GCSAA Booth # 5052

TRIAL OFFER!

Buy one Pallet of Organic-Gro™ PRO

with UFLEXX™ and receive a 40 lb. bag of Organic-Gro™ 4-6-4 Flower & Garden Fertilizer FREE!

Offer ends 4/30/04.



Golf Course Pesticide Label Update

Brand Name/ Manufacturer	Active Ingredient/ Formulation(s)	Label Change/ Date Made	Geography	Benefit/Purpose
Carbait 5 insecticide Agrisel USA (877) 480-0880 www.agrisel.com	Carbaryl (5% granular)	New product registration (July 2003)	U.S.	Control of mole crickets and 120 insect pests with broad labeling.
Insignia fungicide BASF (800) 545-9225 www.turffacts.com	Pyraclostrobin (20% WDG)	New EPA registration (October 2003)	U.S./state registrations pending	Extended control of 15 turfgrass diseases and suppression of dollar spot.
Emerald fungicide BASF (800) 545-9225 www.turffacts.com	Boscalid (70% WDG)	New EPA registration (July 2003)	U.S./pending registrations in CA, HI & SD (no active registration in NY)	Preventive control of dollar spot and new chemistry for resistance management.
Pedulum AquaCap BASF (800) 545-9225 www.turffacts.com	Pendimethalin (EC)	Added state label (July 2003)	California	Now registered for non-residential use in all states (except Alaska) for preemergence control of 40 grassy and broadleaf weeds.
Proxy PGR Bayer Environmental (800) 331-2867 www.bayerprocentral.com	Ethephon (21.7% Liquid)	Tank mix with Primo MAXX (Syngenta) (November 2003)	U.S./state registrations pending	Slows cool season growth to reduce mowing and suppress <i>Poa annua</i> and white clover seed head development
Compass fungicide Bayer Environmental (800) 331-2867 www.bayerprocentral.com	Trifloxystrobin (50% WDG)	Label amendment (July 2003)	U.S.	Suppression of dollar spot on bentgrass greens, tees and fairways.
Chipco Signature fungicide Bayer Environmental (800) 331-2867 www.bayerprocentral.com	Aluminum tris (80% WDG)	Tank mix with Bayleton fungicide (September 2003)	U.S.	Broad spectrum disease control.
Revolver herbicide Bayer Environmental (800) 331-2867 www.bayerprocentral.com	Foramsulfuron (2.25% SC)	New EPA registration (February 2003)	U.S./state registrations pending	Postemergence selective removal of cool-season grasses from warm-season grasses.
Curfew soil fumigant DowAgroSciences (800) 255-3726 www.dowagro.com/turf	1,3 dichloropropene (EC)	Added state labels: (FL February 2003) (SC March 2003) (GA November 2003)	FL, GA & SC	Nematode control for improved playing surface and healthier turf.
Dimension herbicide DowAgroSciences (800) 255-3726 www.dowagro.com/turf	Dithiopyr (EC & WSP)	Remove golf green use (2003)	U.S.	Turf safety.
Dimension herbicide DowAgroSciences (800) 255-3726 www.dowagro.com/turf	Dithiopyr (EC) (Ultra 40WP)	New use direction for <i>Poa annua</i> (EC February 2003) (WP April 2003)	U.S.	Extended control of <i>Poa annua</i> .
Spotlight herbicide DowAgroSciences (800) 255-3726 www.dowagro.com/turf	Fluroxpyr (EC)	New EPA registration (November 2003)	U.S./state registrations pending	Improved postemergent broadleaf control, especially on clover.
Eagle fungicide DowAgroSciences (800) 255-3726 www.dowagro.com/turf	Myclobutanil (20EW)	New formulation (June 2003)	U.S./state registrations pending	Dollar spot and brown patch control in 2.5 gallon container.
Power Zone herbicide PBI Gordon (800) 821-7925 www.pbigordon.com	Carfentrazone, MCPA, MCPP & Dicamba (EC)	Reduced rates/added weeds (July & September 2003)	U.S.	Lower rates for sensitive weeds and 22 weeds added.
Monument herbicide Syngenta (800) 334-9481 www.syngenta.com	Trifloxysulfuron-sodium (75WG)	New EPA registration (November 2003)	U.S./state registrations pending	Sedge control in warm season turfgrasses, overseeding management in the transition zone, and selective broadleaf control.
Primo MAXX Syngenta (800) 334-9481 www.syngenta.com	Trinexapac-ethyl (MEC)	Tank mix with Proxy PGR (Bayer) (November 2003)	U.S./state registrations pending	Slows cool season growth to reduce mowing and suppress <i>Poa annua</i> and white clover.

The *Art* of *Water Management*



Turf management is a fine art and your golf course is the canvas. Every great artist demands the highest quality tools to create and maintain that work of art.

Aquatrols has a range of innovative products that help you achieve success through the more effective use of your resources to meet the specific needs of your turf.



*We have the tools
you need for
your masterpiece.*



5 N. Olney Ave. Cherry Hill, NJ 08003 • USA • (800) 257-7797 • www.aquatrols.com

USE READER SERVICE #26

SUMMARY

Annual ryegrass and intermediate ryegrass transitioned faster than most perennial ryegrass varieties.

DATA

2000-2001 and 2001-2002 from 10 southern and western sites. Sponsored by USGA, NTEP and GCSAA.

SOURCE

Kevin N. Morris, Executive Director of the National Turfgrass Evaluation Program.

MORE INFORMATION

www.netp.org/onsite/ost./htm or kmorris@netp.org.

On-site testing of grasses for overseeding bermuda

Overseeding bermudagrass fairways is a common practice throughout the southern half of the United States. This project evaluated new cultivars on bermudagrass fairways at ten (10) golf courses in the Southern and Western U.S.

The evaluation trials were jointly sponsored by the Golf Course Superintendents Association of America (GCSAA), the United States Golf Association (USGA) Green Section and the National Turfgrass Evaluation Program (NTEP). Trials were positioned strategically in the following areas: southern California; Arizona; Houston, Texas; Dallas, Texas; Mississippi; central Florida; Myrtle Beach, S.C.; Virginia; Atlanta, Ga.; and St. Louis, Mo.

The trials were located on active play sites where golfers hit fairway golf shots and/or drive golf carts. The forty-two (42) entries were established in fall 1999 and then again, in exactly the same physical location, in fall 2000. Grass species entered included perennial ryegrass, intermediate ryegrass, annual ryegrass, *Poa trivialis* and blends and mixtures of these species.

Data from 1999-2000 and 2000-2001 was compiled and published via hard copy, and posted on the NTEP web site (www.netp.org/onsite/ost.htm). Variety performance varied from location to location, however, a number of trends emerged:

Perennial ryegrass entries, in general, provided the highest quality turf averaged over the entire season.

Poa trivialis entries and perennial ryegrass/*Poa trivialis* mixtures were slower to establish, reducing their quality ratings at most locations. However, at three locations, due to other factors, the *Poa trivialis* entries finished on top, complicating the ability to predict where *Poa trivialis* may be used effectively.

Annual ryegrass and intermediate ryegrass entries transitioned faster than most perennial ryegrass entries.

At some sites, the entries that contain *Poa trivialis* transitioned back to bermuda faster than perennial ryegrass, while at other sites, the opposite was true. This leads us to believe that the transition phenomenon is highly weather and management-related.

SUMMARY

Update on second year of low cut trials for Kentucky bluegrass cultivars.

DATA

2000-2002 National Kentucky Bluegrass Test. Sites in Colorado, Maryland, Michigan, Nebraska, New Jersey, New York and Wisconsin (study to continue through 2005).

SOURCE

Kevin N. Morris, Executive Director of the National Turfgrass Evaluation Program.

MORE INFORMATION

www.netp.org/onsite/ost./htm, or kmorris@netp.org.

Low cut trials of Kentucky bluegrass

There is increased interest again in the use of Kentucky bluegrass for fairways and tees. To address this need, several locations of the 2000 National Kentucky Bluegrass Test are being maintained with a low height of cut. Seeded in fall 2000, these trials are mowed at 1 inch or less (most are maintained at 1/2 - 3/4") with 3 - 4 lbs. of nitrogen applied per 1,000 sq. ft. per year and irrigated to prevent dormancy. Trial locations include universities in Fort Collins, Colo.; College Park, Md.; E. Lansing, Mich.; Lincoln, Neb.; New Brunswick, N.J.; Ithaca, N.Y.; Brookings, S.D. and Madison, Wis.


Turfgrass quality data collected in 2001 reflected establishment rate as well as the ability to tolerate a low height of cut. In data averaged over seven of the

(continued on page 56)

TABLE 2.

Kentucky Bluegrass Cultivars in the top statistical grouping in both 2002 and 2003 at low height of cut NTEP locations

Bedazzled
Midnight II
Impact
J-1838
Award
Princeton 105



21 YEARS ON AMERICAN TURF

Over the past 21 years, innovative design, craftsmanship and heavy duty construction make Redexim Charterhouse products the standard of quality in the sports turf marketplace. By building relationships with turf care professionals, Redexim Charterhouse has listened, learned and produced the best machines money can buy. Redexim Charterhouse has not only gained a reputation of being an industry leader, but we have also earned your trust.



**21 YEARS IN AMERICA
AND STILL GROWING**

1-800-597-5664 • Fax: 570-602-3060 • www.redexim.com

USE READER SERVICE #27

SUMMARY:

New selective herbicide options for rough bluegrass control

DATA:

Summer 2002 and 2003 studies on several cultivars of *Poa trivialis* and creeping bentgrass in West Lafayette, Ind. Studies to be expanded in 2004.

SOURCE:

Dr. Zac Reicher, turfgrass Extension specialist, Purdue University

MORE INFORMATION:

www.agry.purdue.edu/turf (under annual research reports), or zreicher@purdue.edu

New control for *Poa trivialis*

There are two basic types of golf courses — those that already have *Poa trivialis* and those that will eventually get it. To find solutions to this significant golf course problem a number of compounds are being tested to determine control.

Poa trivialis is often confused with *Poa annua*, but the difference is important because controls for one do not always work for the other. The following are keys to differentiate the two: *Poa trivialis* is a gasoline or metallic green in color, goes dormant in the summer and has no visible seed heads. In contrast, *Poa annua* is an apple green, dies in the summer, and seed heads will be visible.

Field tests were conducted in the summers of 2002 and 2003 for control of *Poa trivialis*. The most promising compound was Battalion, a Monsanto product with the active ingredient sulfosulfuron. Over the course of the summer of 2002, Battalion gave more than 70 percent control.

The next two most positive controls in 2002 tests were TranXit, by Griffin, with the active ingredient, rimsulfuron, which gave 65 percent control; and a single application of Roundup Pro which gave 60 percent control.

Results in 2003 were disappointing due to record rainfall and very cool weather. In these tests Battalion still gave the most effective control, but the control level was only half the 2002 control level, and that was

achieved only after using a 2X rate from the year before. Another factor being investigated is the possible effect of the grass cultivars.

While Battalion is already a registered and labeled product, the manufacturer is delaying commercial sales until further field tests have been done regarding rates, timing and evaluation of a number of other grassy weeds. This summer field tests with Battalion will be done at more than 50 cooperating golf courses. In addition to control of *Poa trivialis*, the evaluations will include control of yellow nutsedge, tall fescue, quack grass and several other grassy weeds. Battalion has not been shown to be effective on established *Poa annua*.

Field tests to date indicate the following program has been the most effective for Battalion: Four-plus applications at 0.02/LB per AI applied at two-week intervals. Three-week intervals could be used if there is concern about turf safety for bentgrasses. There is slight phytotoxicity that must be tolerated, and the reduction of *Poa trivialis* will be gradual. Timing of applications could begin in late April or early May and continue through mid-June.

The active ingredient is also very active on creeping bentgrass in cool weather, such as applications done in mid-October.

Overseeding with creeping bentgrass can begin three weeks after final application, which would enable seeding to begin in August.

(continued from page 54)

above locations, twenty-nine entries finished in the top statistical grouping (see www.ntep.org/data/kb00/kb00_02-1/kb0002t01.txt). This first year of data showed some trends and some cultivars to watch, but is not enough information to make an informed cultivar decision.

Data from 2002 showed more cultivar separation as all entries were well-established and mature turf stands. Only eleven entries fell into the top statistical grouping in 2002 turfgrass quality data with the commercial cultivar 'Bedazzled' and the experimental entry 'PP H 6366' tied at the top (rating = 6.2) averaged over eight locations (see www.ntep.org/data/kb00/kb00_03-2/kb0003t01.txt).

The top statistical group also included 'Impact', 'Princeton 105', 'Award', 'J-1838', 'Langara', 'Midnight II', 'North Star', 'Nu Destiny' and 'Serene' (see Table 1). Only six entries finished in the top statistical group for turfgrass quality averaged over both years of the low-cut trial locations (see Table 2). This trial will continue through 2005, and 2003 data will be released this spring. Consider at least three years of data before making cultivar decisions.

TABLE 1.

Mean turfgrass quality ratings of Kentucky Bluegrass cultivars grown at eight locations in the U.S.

(Low Height of Cut - Top Statistical Group Only)	
Cultivar Name	Turf Quality Data Mean of 2002 (1-9 Scale; 9=Ideal Turf)
Bedazzled	6.2
PP H 6366	6.2
Impact	6.1
Princeton 105	6.1
Award	6.1
J-1838	6.1
Langara	6.0
Midnight II	6.0
North Star	6.0
Nu Destiny	6.0
Serene	6.0
LSD Value	0.2



More than

simply a

traditional fertilizer, Perk is a true turf revitalizer. Our proprietary slow release organic iron combines iron humate with potash, nitrogen and magnesium to produce the only organic granular controlled-released iron source available. Perk provides deep, long-lasting greening and increased resistance to wear, drought and disease without stimulating excessive growth. Plus, the 1.5% magnesium in Perk even promotes the production of chlorophyll. You'll find Perk perfect for greening up fairways, fields and lawns. For more information, visit www.LebanonTurf.com. Or, for the name of your nearest distributor, call 1-800-233-0628.

And give your turf a little perk-me-up.

LebanonTurf

1-800-233-0628 • www.LebanonTurf.com

USE READER SERVICE #32

Not just green, Perk green.

Perk makes turf green up quickly without excess growth.



SUMMARY

Recycling clippings vs. removing them reduces P fertilization requirements.

DATA

Tracking soil P over four years on stands of Kentucky bluegrass.

SOURCE

Dr. Wayne R. Kussow, Professor of Soil Science at the University of Wisconsin - Madison.

MORE INFORMATION

wrkussow@factstaff.wisc.edu.


Managing soil phosphorus

The floodgates are open. Regulation of P fertilizer use on golf courses is no longer a matter of if, but rather when. When this happens, you'll want to be in the position of being able to demonstrate that you're managing soil P wisely.

The key to managing soil P is in knowing how fertilizer P rates influence soil test levels of the nutrient. We don't have this type of information yet for all the conditions one might encounter on a golf course, but I do have some data that should get you started in the right direction.

The first piece of information comes from a stand of Kentucky bluegrass. I tracked changes in soil test P over four years when different quantities of fertilizer P were applied. The relationships I observed between fertilizer P rate and change in soil test are shown in the figures below. Note that in one case clippings were removed and in the other the clippings were recycled.

The key thing to note in these figures is the annual fertilizer P rate that resulted in no change in soil test. This represents the maintenance rate of P — that required to keep soil test P near its current level. As shown, the maintenance rates of fertilizer P were found to be 0.55 lb. P₂O₅/M/yr when the





clippings were removed, but only 0.22 lb when the clippings were recycled. The difference, 0.33 lb P₂O₅/M/yr represents how much the clippings contributed to soil test on an annual basis.

Knowing the maintenance rate of fertilizer P allows you to make adjustments in your soil test P. Exceed this rate and soil test P can be expected to increase. If your soil test P is excessive, you may elect not to apply any fertilizer P for a while. How long you can go without applying fertilizer P depends how far your soil test P is above the optimum level. You can gauge this by using the equations in the figures. Plug in "zero" fertilizer P₂O₅ for "X" and you get the expected annual reduction in soil test P. It's essentially a reduction of 4 lb soil test P per year when clippings are removed and 1.8 lb. when they're not removed.

Turfgrass uptake of soil P is regulated by clipping production. The more clippings produced, the more soil P the grass removes. During the growing season we typically encounter periods of heat or moisture stress that curtail turfgrass shoot growth. But over an entire growing season, the controlling factor is your annual fertilizer N rate. This is vividly reflected in the diagram below, where annual reductions in soil test P in a bentgrass fairway were strongly dependent on the annual N rate. This, then, is something you need to factor in when trying to gauge what might be a maintenance rate of fertilizer P for your turf. According to this figure, at an annual N rate of 2 lb/M, where the clippings were removed and no fertilizer P applied, soil test P decreased about 2 PPM. For every additional pound of N applied, soil test P declined another 0.72 PPM.

These numbers give you some idea of how to arrive at ballpark estimates of what may be happening to your soil test P and how you can responsibly use fertilizer P in your management program. I do need to caution you that the numbers given here are for the Bray #1 test for soil phosphorus and for turfgrass grown on a silt loam soil. Applying these numbers to soil P extracted by a different method or to soils of very different texture is not advised. GCN



www.sisis.com



sisis®

DESIGNER & MANUFACTURER OF TURF MAINTENANCE MACHINERY SINCE 1932



For full details of the SISIS range and your local dealer please contact:

SISIS INC., PO Box 537, Sandy Springs, SC29677

Phone: (864) 843 5972 Fax: (864) 843 5974 Email: idc@sisis.com

Web site: www.sisis.com

USE READER SERVICE #28

What's new in fairway mowers

MANUFACTURERS ANSWER DEMAND FOR GREENS-QUALITY CUT ON FAIRWAYS

More and more, fairway conditions are getting as much attention as greens. On some courses superintendents are aiming for a greens-quality cut and a striking surface appearance. To help accomplish this, mower manufacturers are introducing specialized fairway mowers or enhancing current products.

Jacobsen 'super lightweight' fairway mower

The Jacobsen Super LF-1880 aims to

deserve its "super lightweight" designation. At approximately 2,000 pounds, it weighs 400 pounds less than comparable two-wheel-drive models and 500 pounds less than four-wheel-drive mowers, reducing turf compaction.

Jacobsen fairway product manager Larry Jones offers some insight into the concept and design philosophy of the Super LF-1880: "Many courses that want the best quality fairway conditions use triplex greens mowers," he says. "These units are also used to mow landing areas that are

becoming larger and more contoured.

One of the challenges with triplexes in this application is that productivity is reduced because of their 62-inch cut, compared to 100 inches on a standard five-gang fairway mower. Also, triplex mowers and conventional five-gang fairway mowers are outfitted with 22-inch-wide cutting units, which occasionally limit their ability to follow fairway undulations under contoured situations."

With an 80-inch cut, the Super LF-1880 increases productivity compared to



Photo: Jacobsen

Jacobsen's Super LF-1880 has been designed for affordability and versatility. Engineered for fairways, the mower also handles green surrounds and long tee boxes.