Another item of importance was the mowing of the fairways. This gave me food for thought. How was I going to cut the fairways without interrupting the play? I finally decided to, cut them at night; so I had the tractors equipped with lights and the men cut the fairways every night, starting about 5:30 p.m., finishing about 2 a.m.

It is significant that on the seven greens which were hit so hard in July, I did not see a sign of brown-patch the remainder of the season, but on the greens which were *not* hit in July, the small or dollar patch was much in evidence.

## Planning Season's Maintenance Campaign

## By FRED A. BURKHARDT Westwood Country Club President, Cleveland District Association of Greenkeepers

THE greenkeeping profession is advancing by leaps and bounds and what I might consider the right thing to do this year maybe all wrong next year. Like any other scientific research work some things are stumbled into and others very carefully thought out.

Following is the program I intend to follow this season, weather conditions permitting. I do not claim that this can be used on other courses but to my particular case it seems to fit and work very nicely. The first part of April I shall use about 15 Ibs. of sulphate of ammonia to greens that average about 6000 square feet. I will follow with applications every week for the rest of the summer but reduce the amount each week until I get to about 5 lbs. per 6000 square feet and run through the hot summer months with this amount. In the fall I raise the amount again until I reach 10 Ibs. per 6000 square feet. My greens get two top-dressings a year, one in April and one in September. The topdressing contains top-soil, sand, and char coal. Before the greens are top-dressed I give them a light covering of tobacco dust, about a hundred lbs. to a green about 6000 square feet. The method I have just told about has been very successful on my course and that is as much as I will say a about it.

I have found out through sad experience that it does not pay to use new compost with a great amount of manure in it, because I have had a great deal of trouble with it. The difficulty I had last fall made me go back to lime and I intend, to use some this year. The amount I use will be very small.

No doubt you have heard about fertilizers causing a great deal of trouble. Job do not think it is fair to condemn the fertilizer itself but the way it is used. You can over-feed and under-feed just as easily as feed the right amount. I think that if you can find the right proportion of fertilizer for your grass and not go to extremes but stay at a happy medium you will not be able to blame fertilizers for other diseases.

## Scald Is Real Horror.

Brown-patch of course is the terrible enemy of the greenkeeper! But give me brown-patch instead of scald. I can bring \* my grass back in a short time with brownpatch but it is a different story with the scald. Last spring we had some unusual weather. A cloud burst of rain in the morning followed by a sun that would ð cook eggs, and what did we get? A blistered grass that was hardest in the world to bring back to life. I have no plan to"\* offer for scald. For brown-patch a light application of lime might prevent it. If you have brown-patch start a little raking, and some nitrate of soda will bring them back if you can start at the work the, minute it turns up.

The golf course today is getting motor, ized. I might spend just as much money this year, with all my equipment motorized, but that is not saying I am spending it for the same amount of work. I am, absolutely sure that I can give more time and attention to things that were impossible when we had hand equipment.

Watering is an important factor but , here again conditions of course say whether you should water one day a week or two. I water each green twice a week during the dry season. The watering is done in the evening in about two and half hours. We cover our greens with about \* ¼ inch of water per night. This will change according to conditions of the green.