



GRAU'S Answers to Turf Questions

BY FRED V. GRAU

Food Reserves for Survival and Re-growth

During the waning days of summer and the early days of autumn nature provides conditions whereby plants start to prepare for winter. Some plants will go into dormancy so deep that no growth will be evident. Others simply slow their growth processes while retaining green color in leaves. One of the important factors that influences winter survival and successful re-growth in the spring is that of *food reserves* in various plant parts.

Scientists freely admit to an incomplete knowledge of how and in what form the food reserves are stored in various plants. One point of agreement is found in the role played by nitrogen. Every known organic compound found in food reserves in plants is associated with the nitrogen supply, supplemented by other essential nutrient elements.

Encourage Fall Fertilization

For many years, agronomists have encouraged fall fertilization of turfgrass areas even though they have not always spelled out how the plants benefit. Basically, we try to provide *adequate* supplies of nitrogen, phosphorus and potash at the time when grasses most effectively can use these "building blocks" for the creation of food reserves. Nature, in her own way, builds the proteins and amino acids which will assure winter survival and vigorous new growth in the spring. Ap-

parently the process is virtually identical in both warm-season and cool-season grasses.

Weeds Can't Use Nutrients

Aside from the highly-essential factor of building needed food reserves, fall fertilization is good practice because annual summer weeds have run their course and thus can't utilize the nutrients designed for the grass. Cool nights and warm days favor food formation in leaves and subsequent transfer to storage organs. Rainfall can be expected to be more favorable for grass growth and utilization of fertilizer. Most turf species tend to produce less topgrowth and denser bottom growth during the fall period. Seedhead formation has occurred weeks before so there is less tendency to produce upright growth for seedhead support.

When we stress fall fertilization we do not mean to imply that some feeding at other times of the year is unimportant. The high rate of nitrogen consumption by many turfgrasses demands fertilization other than in the fall.

May Need Complete Feeding

On some turfgrass areas soil tests indicate that complete feeding is required. By this is meant N-P-K. The proportions of N to P to K should be adjusted according to the need as indicated by the soil test results. The guiding principle should be adequacy, not excess.

Soil tests may indicate that on some areas only one or two nutrient elements

Mercury: Symbol, Hg. (Often called quicksilver.) Atomic weight 200.61 (compared to hydrogen, atomic weight 1.008). It is said to be the most unpredictable element in kinds and properties of compounds it forms. Hg played an important part in the alchemy of the Middle Ages. It is a potent poison. Pure Hg is a silvery liquid at ordinary temperatures. "Quick-silver" means "alive and moving silvery metal." It conducts electricity, is used in thermometers, barometers and in alloys with other metals. Gold and silver dissolve in mercury.

In nature, Hg is found as cinnabar, a red sulfide associated with drops of native metal. Most mercury is produced in Italy, Spain, Calif. and Tex. It is marketed in iron flasks of 75 lbs. each. Annual production is about 24,000 tons. The United States produces 9 per cent of the world total.

Mercury long has been used as a fungicide. Bichloride of mercury (HgCl_2) is used as a vermicide (worms) as well as a control for brown-patch. Calomel (HgCl) is used in human medicine and is a widely-used fungicide (dollarspot). Calomel goes into water solution with great difficulty, is helped by the addition of an equal amount of common table salt.

Solubilized organic forms of mercury are used to reduce bacteria, slimes, algae and other organisms in paper making. Phenyl-mercury acetate is used for certain weed as well as disease control.

Best-known antidote for mercury poisoning is white of egg (albumin) with which it combines directly. Action of Hg may be so fast that no antidote may be entirely successful.

Primers for cartridges are prepared with mercury fulminate, an extremely explosive substance, made from mercury and nitric acid in the presence of alcohol.

are needed. If, for example, P and K are shown to be adequate, then nitrogen only need be applied for satisfactory preparation for winter. If P is adequate, with K deficient, then N and K should be used. Potash is especially important in preparing for winter because of its influence on the winter hardiness of plants.

Kind, Amount Important

Fall use of nitrogen deserves careful consideration. Both the kind and the amount of nitrogen are important. Soluble sources (urea, nitrates, sulfates) must be used with discretion since their complete, immediate availability may force grasses into undesirable rapid growth, causing them to become so "soft" at the onset of winter that winter survival will be poor. One wonders if this may not have been partially responsible for recent wide scale loss of many warm-season grasses.

Organic sources of nitrogen (ureaform, natural organics) do not force rapid growth by virtue of their manner of release to the plants. Nitrogen release from organics is controlled by soil organisms which are extremely sensitive to environmental changes. In effect, the grass plants will be furnished nitrogen in amounts that closely parallel the needs of the plants.

For this reason then, fall fertilization should be concerned primarily with organic sources of nitrogen (plus P and K as needed) to give grasses the best chance to develop winter food reserves in accordance with their needs. Another sound reason is that fall applications of insoluble nitrogen create a reserve of residual nitrogen which, unleached, unused and unchanged, will be there in the soil ready to be released as soon as microbial

(Continued on page 73)

Grau's Answers

(Continued from page 50)

activity starts in the spring. Start the program early enough in the fall to allow plants to "harden off" for maximum winter survival.

How to Convert

Q. We have a few greens that originally were seeded to Colonial and Highland bents. During a recent attack of dollarspot we nearly lost these greens in spite of a good nitrogen feeding program. Each time we changed cups we would throw away the old plug and use a fresh Penncross plug from the nursery. Not one of those Penncross patches had a single dollarspot. My chairman and I agree that we should convert to Penncross but we are not agreed on the method. We do not want to rebuild and keep the green out of play. Can you help us? (Michigan)

A. To keep the greens in play and to preserve contours, I suggest multiple spiking followed by hydroseeding $\frac{1}{2}$ pound of Penncross seed to 1,000 sq. ft. You can do this twice a year at low cost, using your power sprayer, and gradually convert to Penncross. Space prohibits giving detailed instructions here. Send a self-addressed, stamped envelope for a mimeographed

sheet on hydroseeding to GRAU, College Park, Maryland, 20740.

"Seeding" with Algae


Q. Our lakes on the course constitute our water supply. There has been no rain for weeks and the water is low and green with algae. We know that we are "seeding" our greens with algae every time we water but we can't help it. We don't dare try to kill the algae in the lakes because of our wild life. Is there anything we can do to counteract the algae? (Texas)

A. Yes, you can irrigate heavily at the longest possible intervals to keep the greens surfaces as dry as possible. Algae can thrive only with continuous ample moisture. Spiking the dry greens surfaces will let air through the algae crust. Periodic light dusting or spraying with hydrated lime will effectively reduce algae with no harm to the grass. Use one-quarter to one-half pound hydrated lime to 1,000 sq. ft. Apply in late afternoon and allow lime to remain on leaves overnight. Pray for rain. Hook up to city water.

Change in Fertilizer

Q. "For the last several years we have used ureaform (38%) as the principal source of nitrogen on our putting greens. Recently we were advised by some students that this is wrong and that it would be better to make weekly applications of soluble urea (45%). What is your opinion?" (Indiana)

NEW BAG RACKS



INCREASE EARNINGS AND SAVE SPACE

*Service is in sight
with vertical bag racks*

Your club members will appreciate the extra convenience and service that only Stafford Vertical Bag Racks can offer. And, like pro shops across the nation, you can profit from the 40% additional space (compared to other storage systems) this rack makes available. Additional benefits are:

- Wear and tear of golf bags is completely eliminated
- Damp bags dry faster—no mildew
- Faster, easier storage
- Provides a neat appearance
- Easy to install

PLAN TODAY TO CONTACT

THE A. G. STAFFORD CO.

2006 Allen Ave S. E. • P. O. Box 287 • Canton, Ohio
Phone 453-8431