



FOR  
GREENS AND TEES

**PENNCROSS BENT  
AND  
TORONTO C-15 BENT  
SOD or STOLONS  
ALSO  
FYLKING & WINDSOR**

Quality Growers for 22 years



4301 WEST FLOSSMOOR ROAD  
TINLEY PARK, ILLINOIS 60477  
312-798-2210

For more information circle number 273 on card



**COMPLETE  
GOLF  
COURSE  
CONSTRUCTION**

- Rebuild Greens
- Rebuild Tees
- Install Watering Systems
- Move Tees

**IBERIA EARTHMOVNG  
SERVICE, INC.** Iberia, Ohio 43325  
Phone (Galion) 419/468-5454

For more information circle number 198 on card

If  
it's an  
Award  
... We  
Have  
It!

*Golf Prizes*




TROPHIES • PLAQUES • JEWELRY  
HOLLOWARE • TOURNAMENT FAVORS  
NOVELTIES • FUNCTIONAL PRIZES

Custom Prizes Our Specialty  
For Pro Discounts . . . FREE Catalog  
Write Dept. GD Phone (312) 782-2317

**LANE**

EDWIN W. LANE CO.  
32 W. Randolph St.  
Chicago, Ill. 60601

For more information circle number 169 on card



by Dr. James B. Beard

**TURFGRASS  
RESEARCH  
REVIEW**

**Rate of evapotranspiration  
from turf**

Turfgrass evapotranspiration. R. Tovey, J. S. Spencer and D. C. Muckel. 1969. *Agronomy Journal*. 61:863-867. (from the University of Nevada, Reno, Nev. 89507).

The lysimeter technique was utilized to measure the rate of evapotranspiration from two types of turf grown on two different soil textures. The two types of turfgrass were (a) a cool season turf composed primarily of Kentucky bluegrass and red fescue and (b) two varieties (Tifway and Tifgreen) of the warm season species, bermudagrass. Cutting height, mowing frequency, fertilization and pest control were typical of the cultural practices used to maintain Kentucky bluegrass-red fescue and bermudagrass sods, respectively, under Nevada conditions. The two soil textures were a loam and a sandy loam. The irrigation treatments were (a) twice weekly, (b) a seven day interval and (c) a 10 day interval. The total amount of water applied in each case was equivalent to two inches a week. Each irrigation and soil treatment was replicated three times.

Measurements taken included (a) amount of water draining

through the lysimeter, (b) amount of water applied and (c) total quantity of water lost by evapotranspiration from the turf. The latter measurement was compared with the pan evaporation rate and net radiation measurements. Visual ratings of shoot density and color of turf were also taken.

Based on the turfgrass density and color ratings, twice weekly irrigations maintained excellent turfgrass quality throughout the growing season including the hot, dry periods. A seven day irrigation interval produced a good quality Kentucky bluegrass-red fescue turf when grown on the loam soil, but was not adequate for the sandy loam soil. A 10 day irrigation interval resulted in thinning of the Kentucky bluegrass-red fescue turf and the invasion of weedy species. Irrigation at a seven day interval was adequate for maintenance of a high quality bermudagrass turf on both the loam and sandy loam soils. The favorable turfgrass responses to the various irrigation frequencies were only achieved when sufficient water was applied to bring the soil root zone to field capacity at each irrigation.

The rate of evapotranspiration was positively correlated with the rate of pan evaporation and net radiation. Basically, the rate of evapotranspiration from irrigated turfs which have an adequate supply of available soil moisture is governed primarily by the amount of radiant energy received. The rate of evapotranspiration from the turfs grown in Nevada ranged from a 0.13 to 0.16 inch a day during the high temperature periods of midsummer.

Examination of the root distribution of the cool season species showed the majority of the roots in the upper six to eight inches of the soil profile with some roots extending down to 12 inches. The warm season bermudagrass varieties had a considerably deeper rooting capability.

The authors stress that these results were obtained under controlled experimental conditions in Reno, Nev., and that this should be taken into consideration when utilizing this information in the design of irrigation systems.

(Continued on page 26)

## PUT WINTER TRENDS ON YOUR GREENS

Highland Colonial Bentgrass—one of the most economical of the fine-bladed varieties—offers a distinct advantage in an overseeding program. It shows uncommon strength during that crucial spring transitional period when your dormant grasses are coming on. While many of the cool-season grasses in your mixture may fold sooner than you like, you can count on Highland to hang in there until the native grasses recover.

With 8 million seeds per pound, your potential number of plants is exceptionally high. Highland holds its color in the winter and you can close-cut it to 1/4 inch.

Write for free brochures Dept. B

### Cultural Practices

Answers to Questions About Highland Bent

Highland Bent on the Golf Course



## HIGHLAND COLONIAL BENTGRASS COMMISSION

SUITE 1, RIVERGROVE BUILDING

2111 FRONT STREET N. E.

SALEM, OREGON 97303

For more information circle number 205 on card

## PAR GOLF MANUFACTURING CO.

MINIATURE

PAR 3



DRIVING

FINEST RESALE EQUIPMENT

Let Par Golf put you in the Golf Business with their unique and complete portable Driving and portable Miniature courses. For further information contact

Par Golf Mfg. Co.

1117 46th Ave., Rock Island, Ill. 61201

Phone: (309) 788-8461

For more information circle number 239 on card

U.S. PAT. #2989247

# TURFTOP...

LARCHMONT'S "TURFTOP" A PLAYERS DREAM  
A NEW INNOVATION IN SPRINKLERS



Call or write on your letterhead for details and specifications

LARCHMONT  
ENGINEERING  
& IRRIGATION INC.

LEXINGTON, MASS. 02173  
(617) 862-2550



We use

Johns-Manville Transite Pipe  
for all permanent installations

Johns-Manville  
TRANSITE  
IRRIGATION PIPE



For more information circle number 226 on card

## BEARD

(Continued from page 24)

Comments: Evapotranspiration is defined as the combined loss of water from the soil by evaporation and by transpiration from the turf growing on the soil. Transpiration accounts for most of the water loss by evapotranspiration from turfgrass areas. The transpiration rate of a turf will vary seasonally, diurnally and even from hour to hour. As shown in this study, the evapotranspiration rate is typically highest in midsummer. On a diurnal basis, evapotranspiration is normally at a maximum at or slightly after midday, when the light intensity is highest.

Factors affecting the transpiration rate include (a) duration of daylight, (b) temperature, (c) atmospheric vapor pressure, (d) wind movement, (e) rate of water absorption by the turfgrass roots and (f) the soil moisture content. High temperatures, long daylengths, low atmospheric water vapor contents and high wind velocities are environmental factors which can increase the rate of turfgrass evapotranspiration.

The frequency of irrigation will vary with the rate of evapotranspiration and can be influenced significantly by the texture of the soil as illustrated in the above study. Coarser textured soils will require more frequent irrigation because of the lower water holding capacity. However, the total quantity of water applied per irrigation will be less on sandy soils.

Proper frequency of irrigation is particularly important. If water is not applied frequently enough the turf will wilt and damage may result from desiccation or high temperature stress. On the other hand, excessively frequent irrigations which cause waterlogging of the soil will result in a reduction in shoot and root growth plus a decline in the overall vigor and quality of the turf. Over watering is usually more of a problem on golf turfs than under watering.

With water supplies becoming more limited and expensive, it is important to adjust irrigation practices to utilize the water as efficiently as possible in maintaining high quality turfs. An understand-

ing of the evapotranspiration process and the environmental, soil and cultural factors affecting it is quite important to the professional turfmen in planning irrigation systems and executing proper irrigation practices.

Influence of fungus isolate and grass variety on *Sclerotinia* dollar spot development. H. Cole, J. M. Duich, L. B. Massie and W. C. Barber. 1969. *Crop Science*. 9(5): 567-570. (from the Pennsylvania Agricultural Experiment Station, University Park, Pa. 16802).

The relative susceptibility and resistance of Kentucky bluegrass and red fescue varieties to four isolates of dollar spot (*Sclerotinia homoeocarpa* F.T. Bennett) were investigated. The four isolates were obtained from Virginia, Rhode Island, central Pennsylvania and southeastern Pennsylvania. The individual isolates were cultured and inoculations applied to turfs maintained under field plot conditions. Thirty-seven bluegrass varieties and experimental selections which had been maintained at a 1.0 inch cutting height were included in the study along with 18 seeded and vegetatively propagated creeping bentgrass varieties and experimental selections maintained at 0.25 inch.

The turfs were mowed immediately prior to application of the inoculum of the four isolates. Subsequently, a light irrigation was applied for 10 minutes of each daylight hour for four days. This practice insured ideal conditions for penetration and infection of the casual organism. Data taken included the number and severity of dollar spot infection centers developed in the treated areas.

Dollar spot leaf lesions became visible on the bluegrass within two days after inoculation and visible infection centers were evident after five days. In the case of bentgrass, leaf lesions were visible within five days and infection centers appeared within seven to 10 days after inoculation. Temperatures during the bentgrass inoculation experiment were generally cooler than during the bluegrass experiment which contributed to the differential rate of visual injury

(Continued on page 28)



# Aspon<sup>®</sup> Insecticide lets your grass grow free

With Aspon protection, lawns and turf grow free from chinch bug damage. Free from unsightly brown patches of grass caused by their feeding.

Chinch bugs can damage grass before you know they are there. These tiny insects work just at ground level, at the base of the plant, and suck the life out of healthy grass.

Irregular dying areas that appear in warm weather is one sign of possible infestation.

Aspon is one of the most outstanding chinch bug control products. Apply Aspon as soon as you detect their presence.

You get quick initial action plus long-lasting control.

Aspon is recommended by leading grounds maintenance experts... has been successfully used by homeowners, golf course superintendents and commercial lawn care specialists.

Aspon is easy to handle and use. It is not highly toxic and does not persist in the environment. Check instructions concerning watering before and after treatment. Usually, one to two applications a year will do the job.

This season, save your lawns and turf from chinch bug damage with Aspon insecticide.

For best results, use Aspon correctly by following label directions carefully Stauffer Chemical Company, Agricultural Chemical Division, Dept. HD, 299 Park Avenue, New York, NY 10017.

Aspon from 

## BEARD

(Continued from page 27)  
symptom development.

The four dollar spot isolates varied significantly in severity of disease infection on both the Kentucky bluegrasses and creeping bentgrasses. Substantial variations in relative susceptibility among the varieties of bluegrass and bentgrass were also evident. For example, the dollar spot isolate,

which was most virulent on bentgrass, was least virulent on Kentucky bluegrass. Similar pathogenic variations and reversals were evident among the turfgrasses.

In evaluating the relative susceptibility to all four dollar spot isolates, the Kentucky bluegrass varieties Merion, Newport and Pennstar were the most susceptible whereas Delft and Campus were the most resistant.

*Comments:* During sexual repro-

duction a given disease causing fungus organism can produce isolates or strains of the fungus which have distinctly different properties in terms of mycelial characteristics, pathogenicity and tolerance to fungicides. Investigations such as the one reported in this study are needed in order to survey the prevalence and severity of isolates of a given fungus so that resistance to the particularly virulent isolates can be incorporated into the new varieties being developed by the turfgrass breeding program.

The variability in virulence among isolates of a disease causing organism is the reason that a given turfgrass variety may appear relatively resistant to a disease in one region. Similarly, a certain fungicide may be effective in controlling the disease in one area but not in another. The reason for this differential response is that the separate regions have distinctly different isolates of the same fungus. Similarly, the optimum temperature and moisture conditions for disease development may vary among the isolates. Variations in isolates of a fungus also exist in a number of other turfgrass diseases including brown patch, rust and stripe smut.

The variability in turfgrass varietal susceptibility, fungicide control and optimum conditions for fungal development which occur from isolate to isolate within a given disease causing organism illustrates the importance of conducting turfgrass research in the various climatic regions throughout the country rather than at one individual site. It is also equally important for the professional turfman to be up-to-date and informed on the latest turfgrass research results at his own state agricultural experiment station. □

### Other papers of interest:

1. Composted municipal refuse: Its effect on carbon dioxide, nitrate, fungi, and bacteria in arredondo fine sand. *D. F. Rothwell and C. C. Hortenstine. 1969. Agronomy Journal. 61(6): 837-840. (from the Florida Agricultural Experiment Station, Gainesville, Fla. 32601).*

has never been topped!

# CALO-CLOR®

TURF FUNGICIDE



## Still the GREATEST after 40 years!

It's tops "as is" for outstanding turf disease control . . . tops as the mercurial additive for that "extra" effectiveness in any turf fungicide program! Be ready before disease attacks . . . Always have at least two drums on hand! Order from your Mallinckrodt distributor today.

Always spray turf chemicals with the **SPRAY-HAWK (TM)** mobile turf sprayer.



**Mallinckrodt**

**MALLINCKRODT CHEMICAL WORKS**  
ST. LOUIS  
Jersey City • Los Angeles • Montreal

For more information circle number 170 on card