

Gene Flow in Genetically Altered Crops Helps Progress Transgenic Turfgrass

By David Gealy

Genetic improvement of plants through the introduction of a variety of traits — such as tolerance to insects, disease, chemicals, drought or fewer nutrients — is common in agriculture throughout the world. Traditional approaches, such as classical breeding, induced mutagenesis or wide crossing, have a long track record of success. Transgenic technologies, in which genes conferring useful traits of interest are transferred between different species, have been used for genetic improvement for little more than a decade.

Gene flow is the successful transfer of genetic information between different individuals, populations and generations (to progeny) and across spatial dimensions. It is common in nature and maintains the biological diversity that helps ensure long-term survival of populations and species in variable environments. Typically initiated by pollen dispersal in plants, gene flow is a natural and not inherently adverse phenomenon that occurs in non-transgenic as well as transgenic plants and crops.

Gene flow potential in plants is largely determined by reproductive biology (Table 1, p. 54). In some species, flowers attract pollinators such as insects or pollen grains are able to travel long distances by wind. In other species, gene flow potential is restricted because their flowers self-fertilize before pollen dispersal. Thus, the reproductive biology of crops largely governs the frequency of outcrossing. Crop reproductive biology can range from almost exclusively self-pollinating to 100 percent outcrossing. Outcrossing frequency typically decreases with distance between the pollen-producing plant and the recipient plant and with the time period between pollen dispersal and fertilization on the recipient plant. Gene flow potential tends to be higher in crop species that have high natural levels of outcrossing. Pollen dispersal, however, leads to true gene flow only if the dispersed pollen fertilizes flowers leading to seed formation and seeds that germinate to produce plants that are competitive enough to reproduce over

time. Traits such as herbicide resistance can impart a sizeable selective advantage in locations where the herbicide is used. Under this scenario, there is likely to be a greater degree of gene flow success following the initial pollen dispersal and outcrossing event than in locations where the herbicide is not used.

Physical movement of genetic traits via seeds — such as shattering, wind, water or distribution by animals — or vegetative propagules, including rhizomes and stolons, are not considered true gene flow, but it can be economically important depending on the transgenic crop and production practices. Typically, seed traits such as shattering or movement with air currents that encourage dispersal are not found in most domesticated crops. A number of turfgrasses are propagated by stolons or rhizomes.

Corn, soybean, canola and cotton represent about 99 percent of all commercialized transgenic crop production worldwide. Seed dispersal occurs at a minimal level in all these crops. Outcrossing is limited in soybean, wheat and rice. Canola and corn have greater outcrossing.

The United States also grows transgenic papaya, squash and alfalfa commercially. Transgenics have been produced in other crops/species, but few are presently proceeding through the regulatory process toward commercialization. These include turfgrasses as well as grain crops, vegetables, fruits, ornamental plants, forage crops and trees. Additional species granted nonregulated status or in the process of deregulation or approval include creeping bentgrass, carnation, tobacco, tomato, plum, potato, beet and sugar beet.

Gene flow in a larger context

Gene flow is only one of several mechanisms by which plants or seeds become intermixed at low levels with other plants or seeds where humans did not intend them to be. This type of unintended mixing occurs with both transgenic and nontransgenic plants and crops. Other mechanisms include inadvertent physical dispersal or mixing of seed, quality-control failures and inevitable human mistakes. Low or trace levels of commingling via all of these mechanisms is virtually unavoidable in

Continued on page 56

TABLE 1

Examples of useful traits being imparted to plants using all available approaches, and estimates of the probable consequences from gene flow¹

New characteristics being imparted to crop plants ¹	Potential for natural selection process to transform crop into unmanageable "weed" or "volunteer" ²	Potential for natural selection process to worsen existing problem or create new agronomic problems resulting from gene flow to wild/compatible relative ²	Potential that off-site gene flow would create significant adverse human health/nutrition impact ²
Potential grower/agricultural problems			
Potential nonagricultural/human safety problems			
•Herbicide tolerance	med ³	low-med ³	neg ³
•Insect tolerance	low-med	med	neg
•Disease tolerance (fungal, bacterial, viral)	low-med	med	neg
•Nematode tolerance	low-med	low-med	neg
•Salt tolerance	low-med	low-med	neg
•Drought tolerance	low-med	med	neg
•Cold/freezing tolerance	low-med	low-med	neg
•Improved nitrogen use efficiency	low	low-med	neg
•Early season vigor	low	low-med	neg
•Increased growth rate	low	low-med	neg
•Reduced plant height	low	low	neg
•Seed shattering resistance	neg	neg	neg
•Altered flowering time	low	low	neg
•Male sterility in crop plant	neg	neg	neg
•Yield/yield components	neg	neg	neg
•Altered maturity date	low	low	neg
•Altered fruit shape/flavor	neg	neg	neg
•Seed protein or oil content or quality	neg	neg	low-med
•Plant-made pharmaceuticals (PMPs) (many traits)	neg	neg	med-high
•Industrial compounds	neg	neg	med-high

1. Approaches include transgenic as well as traditional, nontransgenic classical breeding approaches that use long-established methods to introduce and select for desirable traits in crops. All traits are eventually manipulated through classical breeding methods before commercialization.

2. Estimates compare the most probable outcomes relating to gene flow. Outcomes of gene flow listed here may differ among transgenic crops; some crop/weed species combinations are more likely than others to have high outcrossing or gene flow; and some weed species are inherently more aggressive than others.

3. Key for degree of problem expected (author estimates). Negligible, "neg"; low, "low"; medium, "med"; high, "high." The degree of the problem is likely to vary for a given characteristic or trait depending on the reproductive biology of the species, the trait itself and the cropping system in which it is used.

Source: Council for Agriculture Science and Technology



QUICK TIP

Summertime is barbecue season, and hopefully we're talking ribs and hamburgers and not ultra-tanned bodies. Also true to summer and hot weather is increased insect activity. One common aspect insects and fertilizers have with turfgrass is the importance of best management practices. BMP's are innovative, dynamic and improved environmental protection practices applied to proper turfgrass management. A good example would be the application of both fertilizers and insecticides in the proper timing to reach the intended result with reduced impact to the environment. Of course, other factors are important as well, such as product selection; so make the right choice, and stay cool.

Continued from page 53

commercial-scale plant production. Thus, requirements for unrealistically low occurrence of unwanted substances, such as transgenic seed, can lead to progressively higher overall costs to industry and consumers without a measurable benefit.

Gene flow from transgenic crop plants to weedy or wild relatives can be an issue if they are sexually compatible. Thus, one way to view gene flow issues in a global context is from the perspective of the important worldwide food/feed crops and the weedy or wild species with which they are sexually compatible. About 200 plant species account for essentially all of the significant economic activities for humans worldwide, and approximately 10 percent of these account for almost all of the human caloric consumption. Among the world's worst weed species, only five groups — related weeds of rice, sorghum, rapeseed, sugarcane and oats — are sexually compatible with the most important crops. The potential for gene flow between crop and weed species might not become actual gene flow due to differing habitats or geographic distribution, genetic barriers to outcrossing, etc. The number of combinations of transgenic crop/weed or transgenic crop/nontransgenic crop that are likely to develop highly troublesome gene flow problems is quite small globally. Problems with particular weed or crop species, however, could occur locally or regionally for certain traits conferred by transgenes.

The biology of the transgene trait will largely determine consequences of gene flow to sexually compatible nontransgenic crops, weeds and wild relatives. Outcrossing from herbicide-tolerant transgenic crops can produce progeny that is highly favored in areas where the particular herbicide is used, but not in other areas. However, ordinary selection pressure from repeated herbicide use in herbicide-resistant crops may lead to a greater development of weed resistance (in the absence of gene flow) than gene selections attributable to gene flow between the crop and weed.

Importation of weeds or other plants

from foreign lands is an historic problem, but instances in which fully domesticated crops have become weed nuisances are rare. The chances that transgenic modification of a single trait in a domesticated crop would change the crop into a successful weed also is low, and gene flow from the majority of transgenic crops probably will have minimal ecological impacts outside of agricultural areas.

Transgenes that confer tolerance to stresses — such as drought, diseases, insects and salty soil — might require different or additional evaluation because gene flow into nontransgenic crops or weedy relatives can provide selective advantages in both agricultural and nonagricultural settings.

Risk assessment for transgenic crops

Before transgenic crops are approved for commercial use, law requires that they pass extensive testing addressing any significant food safety, environmental protection and human/animal health issues. Once approved, concerns about low-level presence of transgenes from gene flow or other mechanisms mostly revolve around economic issues, such as contract specifications and consumer expectations.

Risk assessment is an early step in the approval process for transgenic crops. Risk is the possibility of harm occurring. For a risk to be realized, there must be something harmful or adverse and also a likelihood that it will occur. Hazards do not pose risks unless there is significant exposure. Likewise, significant exposure does not constitute a risk if it poses a very low hazard. Gene flow fits the exposure part of the risk equation. Risk assessments of transgenic crops in the United States use accepted scientific methods and analyses. They are case by case; conclusions are examined in light of new and established information, and they are comparative, using the nontransgenic crop or plant as the basis for characterizing risk. The vast majority of gene flow events between transgenic crops possessing a wide array of transgenic traits and nontransgenic crops or weeds likely will not lead to a hazard. The transgenic crops approved for commercial

Continued on page 60



QUICK TIP

Insect control begins with healthy turf. Pests of all sorts are always present, and the ability of the turf to grow through the invasion is what differentiates fine turf from unacceptable turf. While times might warrant the use of a pesticide to kill the invading pest, far too often the grass is still left in a compromised state that is in dire need of the nutrition necessary for its recovery. Having linear uncompromised growth in place at all times is the best defense against any pest invasion by allowing the plant to out-compete the pest's activity. Read more at www.floratine.com.

Continued from page 56

use in the United States to date have been shown to pose no or minimal risk to health and the environment.

Developing uniform seeds for planting is a major economic concern for producers and suppliers of transgenic and nontransgenic seed. For many decades, procedures for maintaining genetic integrity in seed production have been in place for nontransgenic crops, but some amount of admixture — via pollen flow or mechanical mixing — occurs with all seed-production operations. Seed certification programs, which began in the United States in the early 1900s, typically use a pedigree system that begins with the breeder and progresses to foundation seed, registered seed and certified seed, which is then sold to farmers for planting the commercial crop. Despite numerous precautions, seed laws recognize that achieving zero-tolerance thresholds is not possible.

Isolation of production fields to varying degrees has been used to minimize the inflow and outflow of pollen to and from these fields. This principle can be used with seed production fields, transgenic crops or both. Although this practice can restrict

pollen flow substantially, it cannot assure zero gene flow on a landscape level. Novel molecular approaches to further prevent unwanted transfer or spread of transgenes are being developed and could be implemented where warranted. Among these are: 1) insertion of transgenes into organelles — such as chloroplasts — that do not pass along their genes through pollen; 2) adding a male sterility trait that prevents seed production in crop/weed plants that receive transgenic pollen; 3) genetic use restriction technologies, which cause sterility in second-generation seeds; and 4) tightly linking desired transgenes to a gene that is deleterious to reproductive fitness so transgenic weedy or volunteer populations cannot build up after an outcrossing event.

David Gealy is a plant physiologist with the USDA-ARS Dale Bumpers National Rice Research Center, where he specializes in biology control and gene flow of weeds in rice. He previously worked as a plant physiologist at the USDA for 12 years. Gealy has been a member of the board of directors for the Weed Science Society of America since 2006.



JOHN DEERE

QUICK TIP

Summer months bring blazing temperatures and drought-like conditions, which mean you'll be relying even more on your irrigation system. Be sure to optimize water conservation while still preventing turf damage by replacing worn rotors, nozzles and internal assemblies immediately. For more information on irrigation rotor replacement components that will retro fit most manufacturer models, contact your local John Deere Golf distributor.

REFERENCES

Council for Agricultural Science and Technology (CAST). 2007. Implications of gene flow in the scale-up and commercial use of biotechnology-derived crops: Economic and policy considerations. Issue Paper 37. CAST, Ames, Iowa.

Bradford, K. J. 2006. "Methods to Maintain Genetic Purity of Seed Stocks." Division of Agriculture and Natural Resources Publication 8189, University of California, Oakland. <http://anrcatalog.ucdavis.edu>

Bradford, K. J., A. Van Deynze, N. Gutterson, W. Parrott, and S. H. Strauss. 2005. Regulating transgenic crops sensibly: lessons from plant breeding, biotechnology, and genomics. *Nature Biotechnol.* 23:439-444.

Gressel, J. and H. Al-Ahmad. 2005. Molecular containment and mitigation of genes within crops—Prevention of gene establishment in volunteer offspring and feral strains. Pp. 371-388. In J. Gressel (ed.). *Crop Fertility and Volunteerism*. CRC Press, Boca Raton.

Information Systems for Biotechnology. 2007. A National Resource in Agbiotech Information. <http://www.isb.vt.edu/>

James, C. 2006. Executive summary of Global Status of Commercialized Biotech/GM Crops: 2006. ISAAA Brief No. 35-2006. ISAAA, Ithaca, New York. <http://www.isaaa.org>

U.S. Department of Agriculture—Animal and Plant Health Inspection Service (USDA-APHIS). 2007. Petitions of Nonregulated Status Granted or Pending by APHIS. http://www.aphis.usda.gov/bns/not_reg.html

Warwick, S. I. and C. N. Stewart, Jr. 2005. Crops come from wild plants—How domestication, transgenes, and linkage together shape fertility. Pp. 9-30. In J. Gressel (ed.). *Crop Fertility and Volunteerism*. CRC Press, Boca Raton, Florida.

Watrud, L. S., E. H. Lee, A. Fairbrother, C. Burdick, J. R. Reichman, M. Bollman, M. Storm, G. King, and P. K. Van de Water. 2004. Evidence for landscape-level, pollen-mediated gene flow from genetically modified creeping bentgrass with CP4 EPSPS as a marker. *Proc Natl Acad Sci USA* 101:14533-14538.

Wolt, J. D., Y.-Y. Shyy, P. Christensen, K. S. Dormin, and M. Misra. 2004. Quantitative exposure assessment for confinement of maize biogenic systems. *Environmental Biosafety Research* 3:1-14.

TURFGRASS TRENDS

SECTION STAFF

Managing Editor
Curt Harler
440-238-4556; 440-238-4116 (fax)
curt@curtharler.com

Golfdom Staff Contact
David Frabotta
216-706-3758; 216-706-3712 (fax)
dfrabotta@questex.com

Graphic Designer
Kristen Morabito
216-706-3776; 216-706-3712 (fax)
kmorabito@questex.com

INDUSTRY ADVISORS

Jerry Quinn
John Deere
Chris Derrick
Agrium Advanced Technologies

Scott Welge
Bayer Environmental Science
Carmen Magro
Floratite

EDITORIAL REVIEW BOARD

Dr. Rick Brandenburg
N.C. State University

Dr. Vic Gibault
University of California

Dr. Gerald Horst
University of Nebraska

Dr. Richard Hull
University of Rhode Island

Dr. Eric Nelson
Cornell University

Dr. A.J. Powell
University of Kentucky

Dr. Eliot C. Roberts
Rosehall Associates

Dr. Pat Vittum
University of Massachusetts

CONTACT US:

Web site: www.turfgrasstrends.com

Reprints: TurfgrassTrends@reprintbuyer.com