

file. Root cells are enclosed within cellulose-based cell walls. The carbohydrate polymers that comprise these walls are highly hydrated and constitute an aqueous phase from which root cell protoplasts acquire their water and mineral nutrients. Since roots grow in the soil, their surface cells are bathed in soil water. Any nutrient ions available to a root must find their way from soil water into this cell wall space before they can be absorbed within living root cells.

The environment of the cell wall space is often quite different from that of the surrounding soil. Within the apoplast, much of the water is bonded to the polymers that comprise the cell walls but free water is reasonably abundant as well.

Some of these carbohydrate polymers (pectins) contain sugar-acid units that, at a pH above 4.5, will release a H⁺ from their carboxyl group leaving a negative charge on the polymer. These negative charges will attract and bind with cations much as cation exchange sites do in the soil. Because of these cation exchange sites, water in the cell walls will contain a greater concentration of nutrient cations than will be present in the soil solution and likely a somewhat lesser nutrient anion concentration but one still greater than that of soil water.

Cell wall spaces will have an elevated H⁺ concentration because H⁺s are also attracted from the soil water and in addition, are pumped out of the protoplasts into the apoplast during normal cell functions.

For the reasons cited above, the apoplast from which nutrient ions are actually transported into root cell protoplasts is likely to be more acidic and have a greater nutrient concentration than the soil water.

The protoplasts of root cells are interconnected by tiny protoplasmic tubes called plasmodesmata and constitute a network of living protoplasts that can develop independently yet exchange materials and information among themselves and throughout the plant. This living protoplasmic network is called the symplast. Each cell's protoplast is enclosed within a plasma membrane that separates the living part of a cell from its external nonliving cell wall (apoplast). To enter into a protoplast,

all water and nutrient ions must cross a plasma membrane that is selectively permeable, allowing some ions to cross while excluding others. Thus, the protoplast compartments of a root can have a nutrient composition very different from that of the soil solution and its surrounding apoplast.

It is within the plant that nutrient balance is most critical since it is here that nutrients perform their metabolic functions. Once absorbed by root cells, nutrients are distributed throughout stems and leaves via xylem elements of the vascular system. Transpiration of water from leaves provides the driving force for upward movement of water and nutrient ions within xylem elements. Some nutrient ions can exit the leaves via sieve elements of the phloem and circulate throughout the plant. Such nutrients are likely to be sufficient for growth of meristems (growing points) but may become deficient in leaves. Others have limited phloem mobility and may become deficient in meristems while remaining adequate in leaves. These resulting nutrient imbalances within the plant can cause problems that will be discussed later, but often may be addressed through foliar fertilization.

Plant nutrients will interact with each other differently within each of these three compartments making nutrient availability to, uptake by and distribution within plants a highly complex phenomenon.

We will explore these nutrient interactions in the soil in the next article in this series.

Richard Hull is professor emeritus of plant sciences at the University of Rhode Island and adjunct professor of horticulture at Clemson University. Haibo Liu is professor of turfgrass physiology and management at Clemson University. Liu earned his Ph.D. with Hull at the University of Rhode Island and they continue to collaborate on research and publications in turfgrass physiology and nutrition.

REFERENCES

- Brady, N. C. and R. R. Weil. 1999. *The Nature and Properties of Soils*. 12th edition. Prentice Hall, Upper saddle River, NJ.
- Carrow, R. N., D. V. Waddington and P. E. Rieke. 2001. *Turfgrass Soil Fertility and Chemical Problems: Assessment and Management*. Ann Arbor Press, Chelsea, MI.
- Marschner, H. 1995. *Mineral Nutrition of Higher Plants*: 2nd Edition. Academic Press. London.

Ad Index

Advertiser	Page No.
Agrium Adv. Tech.	CV2-1
Andersons	8, 33
Aquatrols	19-30
B A S F Cover Tip, 6, 15, CV3	
Becker Underwood	13
Bell Labs	47
Champion Turf	40
DuPont	4-5, 38-39
Empire State Green Industry Show	35
FMC	16-17
Jacobsen	9
John Deere	2-3
Kochek	31
NuFarm	36, 37
Pasteuria Bioscience	32
Quali-Pro	11
Rain Bird	19-30
Sto-Cote	32
Syngenta	CV4
Toro	46
Turco Mfg.	34
Valent	18
White Metal Golf	47

This index is provided as an additional service. The publisher does not assume any liability for errors or omissions.