



by Dr. James B. Beard

TURFGRASS RESEARCH REVIEW

Use of benomyl to control dollar spot

Control of dollar spot disease on turf in New South Wales with the new systemic fungicide, benomyl. D. Reilly. 1969. The Journal of the Sports Turf Research Institute. 45:63-66. (from the Department of Agriculture, New South Wales, Aust.)

The effectiveness of a new systemic fungicide, benomyl (methyl 1-(butylcarbamoyl)-2-benzimidazole-carbamate), for controlling dollar spot (*Sclerotinia homoeocarpa* F. T. Bennet) was compared with mercuric chloride in this study. The standard recommendation throughout Australia for controlling dollar spot uses mercury or cadmium containing fungicides. The experiments were conducted on two bentgrass bowling greens in the Sydney area which had visible dollar spot damage. Similar experiments were conducted on a Penncross creeping bentgrass (*Agrostis palustris* Huds.) bowling green at the Cronulla R.S.L. Club and on a New Zealand colonial bentgrass (*Agrostis tenuis* Sibth.) bowling green at the Hills District Bowling Club. The fungicide treatments included (a) mercuric chloride applied at two ounces active ingredient (a.i.) per 1,000 square feet, (b) benomyl (50 per cent wettable powder) applied at 1.5 ounces (a.i.) per 1,000 square feet, (c) benomyl (50 percent wettable powder) applied at three ounces (a.i.) per 1,000 square

feet and (d) an untreated control plot. Surfactant F was added to the two benomyl treatments at a rate of four ounces per 100 gallons of dilute spray. The experimental area consisted of 5 by 10 foot plots with each of the treatments being replicated six times on both bentgrass species. All fungicide treatments were applied in two gallons of water per 50 square foot plot using a watering can. Two treatment applications were made at a 21 day interval. The degree of dollar spot disease control was quantitatively determined by counting the number of two-inch-squares per plot damaged by the pathogen.

Under the conditions of these two experiments both rates of benomyl were more effective in controlling dollar spot than mercuric chloride. The relative per cent control of dollar spot from the three fungicides showed approximately 97 per cent control with the three ounce rate of benomyl, 94 per cent control with the 1.5 ounce rate of benomyl and 65 per cent control with the two ounce rate of mercuric chloride. The author reported that the benomyl appeared to have an eradicant action on the dollar spot disease as well as a longer residual effect than other currently available fungicides.

No visible phytotoxicity to either the Penncross creeping bentgrass or New Zealand colonial bentgrass was observed from the benomyl applied at either rate. Actually, turfs receiving the benomyl application showed improved color, giving the impression that it was stimulating growth. In contrast the two ounce rate of mercuric chloride caused discoloration of the two bentgrasses for an indefinite period of time following its application.

Comments: Dollar spot is a major turf disease in Australia, especially bowling and golf greens. Its occurrence was first recorded from Queensland in 1934. Mercury and cadmium fungicides have been the standards for effective control of dollar spot throughout Australia. Recently, mercury and cadmium have not been as effective in controlling dollar spot disease under certain conditions. The author gave no reason for this development.

The experiments reported in this paper demonstrate that benomyl provides good control of dollar spot without causing discoloration of the turf. Also, this systemic fungicide

(Continued on page 35)

GIVE YOUR GROUNDS A BETTER CHANCE with a W-W SHREDDER GRINDER

Choose the capacity you need to handle all your soil preparation, top dressings, mulches and build compost piles. Shred, grind, mix, screen with easily interchangeable shredding rollers and grinding screen. Handle osmunda, peat moss, even phosphate rock with big capacity to cover large areas—fast!



W-W MODEL 4 with optional Detachable Elevator / Process up to 20 cu. yds. per hour with standard 35" x 36" hopper and 9-hp. engine. Independently powered elevator can be used separately when desired. Options include oversize 36" x 52" hopper for tractor loading and 12-hp. engine for increased capacity.

Interchangeable shredding bars and grinding screen are standard on W-W Shredder-Grinders. Replaceable hammer tips are reversible for double life. Solid cast-iron main frames are guaranteed for a lifetime against corrosion. Write for literature and prices.

SINCE 1910 **W-W GRINDER CORP.** 2957-S No. Market, Wichita, Kansas 67219

For more information circle number 205 on card

There are 101 reasons why a Cyclone Lawn Spreader is your best buy!



When you select a Cyclone, you are getting the benefit of 101 years of broadcast spreading equipment know-how... experience that has made Cyclone the first choice of professional turf men.

- Broadcasts material accurately—evenly
- Stripe-free performance
- Ruggedly built for years of service

Ask about the other Cyclone models that are ideally suited for golf courses.

THE CYCLONE SEEDER CO., INC.
Dept. 60-2, Urbana, Indiana 46990

For more information circle number 182 on card



Get the pipe that eliminates winter rules ...in the summer.

The strength and versatility of cast iron pipe can easily become your most valuable asset.

Even if yours is a four-season course, rugged cast iron pipe can handle the hazards. Such as water hammer, higher and higher pressures, heavily laden maintenance vehicles.

And if you're planning a revamped look for your course, cast iron pipe makes it easy. Dig it up. Move it around. Reuse it. Expand the system to suit your new layout. Add sprinklers when and where you want them—cast iron pipe takes quick, clean, leakproof taps with strip-proof threads...and no water hammer hazards.

As for long life, would you settle for a century (or two) of service? That's how long corrosion-resistant cast iron pipe has been reliably delivering water to hundreds of cities and towns nationwide.

Get set for season after season of worry-free watering. Install cast iron pipe.

Write for your free copy of Booklet A-27, "Golf Course Irrigation." It's packed with design and installation tips and helpful planning guidance.



CAST IRON PIPE RESEARCH ASSOCIATION • Executive Plaza East
1211 West 22nd Street • Oak Brook, Illinois 60521



An association of quality producers dedicated to highest pipe standards through a program of continuing research
THE MARK OF PIPE THAT HAS SERVED OVER 100 YEARS IN OVER 100 CITIES

CAST IRON PIPE


ALABAMA PIPE COMPANY • AMERICAN CAST IRON PIPE COMPANY • CLOW CORPORATION • GLAMORGAN PIPE & FOUNDRY CO. • LONE STAR STEEL COMPANY
LYNCHBURG FOUNDRY COMPANY • McWANE CAST IRON PIPE CO. • PACIFIC STATES CAST IRON PIPE CO. • UNITED STATES PIPE & FOUNDRY COMPANY

For more information circle number 253 on card

Photo taken at Palmetto-Pine Country Club, Lake Okauchee, Virginia Beach, Virginia



Lark



Hawk

Chickadee

The Red Rider is for the Birds...

The Famous Red Rider

the most-proved, most dependable turf work cart in the land—has been joined by America's best known turf birds... the LARK, HAWK, CHICKADEE and GOLDEN EAGLE... the finest sprayers and equipment ever to go to work on a golf course. Here's the first team of the turf... ready to go to work for you!



Golden Eagle Pump
The heart of the SMITHCO SPRAYERS

SMITHCO Inc.
11 WEST AVENUE, WAYNE, PA. 19087

Smithco Sprayers formerly manufactured by Thuron.

For more information circle number 166 on card

GET
GET

Cutting Height as Low as....

3/8"

Frequency of Cut as Close as..

1/2"

10 BLADE HIGH SPEED REEL: For the finer groomed creeping Bent in the North and the new improved fairway Bents and Bermudas in warmer climates.

EVEN DISTRIBUTION OF CLIPPINGS: With new scraper-deflector, grass clippings are evenly distributed, eliminating bunching, dropping and windrowing.

PREVENTATIVE MATTING AND THATCH CONTROL: Presents a preventative control of thatch and matting by cutting as low as 3/8 inch with a 1/2 inch frequency of cut.


ELIMINATES SCALPING ON MOUNDS: The 26 inch cutting swath of each unit gives greater flexibility. No "ribbed appearance." Smoother, more uniform cut.

AVAILABLE in Hollow Roller or Rear Wheel Drive. 3, 5, 7- and 9 gangs.


Gives Putting Green Appearance to Fairways, Aprons and Tees!



with the **NEW**
ROSEMAN
LO-CUT MOWER



ROSEMAN 7 1/2" DIAMETER REEL WITH 10 CHROME-NICKEL ALLOY STEEL BLADES



ROSEMAN ROLLER DRIVE and ROSEMAN REAR WHEEL DRIVE



ROSEMAN MOWER
CORP.

Nation-Wide Sales and Service 2300 W. LAKE AVE. • GLENVIEW, ILLINOIS

For more information circle number 213 on card

does possess a residual effect.

Practically all turfgrass fungicides which have been used in the past are of the non-systemic type. That is, disease control is achieved by killing the spores, mycelia and sclerotia present on the turfgrass surface at the time of fungicide application and by killing inoculum which is subsequently deposited on the turfgrass surface during the active residue period of the fungicide. Thus, effective control by a non-systemic fungicide is only achieved on those plant parts to which the fungicide is applied and only for the period of time that the fungicide remains on the plant surface in an active state. In contrast to this, a systemic fungicide is absorbed and translocated throughout the plant. It can eradicate fungi which have already infected the host plant and can also prevent subsequent infections by the pathogen. A systemic fungicide is preferred to a non-systemic type providing the disease control is comparable and there is no phytotoxicity to the turfgrasses. Benomyl is a systemic fungicide which fulfills these criteria along with

a longer residual on turfs. No doubt, the future trend in turfgrass disease control will be toward the use of systemic type fungicides as they become available.

Effect of time, rate and gallons of water applied on paraquat for the control of annual bluegrass (*Poa annua*). W.W. Huffine, J.L. Coltharp, L.B. Gillham. 1970. *Oklahoma Agricultural Experiment Station Progress Report P-632. pp. 10-11. (from the Department of Agronomy, Oklahoma State University, Stillwater, Okla. 74075).*

Various methods of paraquat application for the control of annual bluegrass were evaluated in this study. The experiment was conducted under greenhouse conditions utilizing annual bluegrass turfs grown in individual pots. Treatments consisted of (a) time of application—8 a.m. versus 8 p.m.; (b) rate of paraquat application—0.25 versus 0.5 pounds of active ingredient per acre; (c) rate of surfactant—none versus 0.001 per cent and (d) rate of water utilized—comparing 40, 80 and 120 gallons of water applied per acre. The per cent control of annual bluegrass was determined seven days after

the paraquat application.

The best contact control of annual bluegrass was obtained by applying paraquat at the rate of 0.5 pounds active ingredient per acre along with a surfactant in 40 gallons of water. The 0.25 pound per acre rate of paraquat applied in 40 gallons of water with a surfactant gave more effective annual bluegrass control than the 0.5 pounds per acre rate applied without a surfactant. Forty to 80 gallons of water per acre was the optimum amount of carrier for paraquat applications. The degree of annual bluegrass control was not affected by the time of application when the 8 a.m. "day" treatment and the 8 p.m. "night" treatment were compared.

Comments: Paraquat provides a contact nonselective post-emergence type of weed control. A contact herbicide kills only those plant parts or living cells to which the chemical is applied. The death of the plant occurs very rapidly after application. A nonselective herbicide kills all plant species or plant parts which the chemicals comes into contact with, regardless of whether desirable turfgrass species are present.

(Continued on page 38)

TURFTOP*... from LARCHMONT



THE NEW INNOVATION IN SPRINKLERS
BLENDS WITH AND BECOMES PART OF THE TURF
ELIMINATES THE HARDTOP

Larchmont SYSTEMS ARE DESIGNED BY TURFMEN

AUTOMATIC SYSTEMS - MANUAL SYSTEMS -
SPRINKLERS - VALVES - COUPLINGS - PUMPS

**LARCHMONT
ENGINEERING
& IRRIGATION INC.**



AGRICULTURE ENGINEERS
LEXINGTON, MASSACHUSETTS 02173

(617) 862-2550

LARCHMONT
AUTOMATIC SPRINKLER SYSTEM

for
Golf Courses - Cemeteries
Municipal Parks - Large Industrial Grounds
WRITE TODAY FOR LITERATURE... THERE'S LOTS MORE TO TELL

* U.S. PAT. #2989247

we use

Johns-Manville
TRANSITE
Asbestos-Cement
IRRIGATION PIPE



■ When you put in something as critical as underground sprinkling, you get only one chance to do it right. And on hundreds of courses since 1961, that has meant Toro equipment.

Toro's Vari-Time* Control System has set the standard.

We've been leaders in sprinkler head design for years. The first Valve-In-Head, the first sealed gear drive, the first 2-speed heads for fairways and greens. A few reasons great championship courses are choosing Toro.

Our equipment costs about the same as other brands cost.

Toro. Because you only do it once.



Because of better design, ours
saves money over the long haul.

With a Toro Central
System, you control the entire
course from one location. Make
instant changes in your program
as the weather shifts. Savings on
labor costs can be dramatic.
Water efficiency is maximized.

Soggy turf and puddling are
virtually eliminated.

Let us help you build and
maintain the kind of turf you've
been dreaming about.

TORO®

We Take Care

Toro Manufacturing Corporation, Moist O'Matic Division 5825 Jasmine Street—P.O. Box 489, Riverside, California 92502



A Toro Vari-Time* Central Control System keeps
the turf here at Oak Ridge Country Club and
over 300 courses installed in the last two years.

*Exclusive Trade Name Toro Mfg. Corp.



**RUGGED!
RAPID!**

That's RYEGRASS

Versatile Oregon Ryegrass is something to consider for use on any greenway.

It's rapid! Both Annual and Perennial Ryegrass will normally germinate in a matter of 6 to 14 days.

Because of this seedling vigor and its ability to stand up under heavy traffic once it is established, Ryegrass is a fine performer in problem areas.

Ryegrass also does an excellent job of stabilizing soil in disturbed areas.

Annual Ryegrass is, of course, a prime ingredient in overseeding mixtures for its early strength.

Both Annual and Perennial Ryegrass develop a strong root system which holds moisture.


For any greenway, remember Oregon Ryegrass.

WRITE FOR A FREE BROCHURE

OREGON RYEGRASS GROWERS SEED COMMISSION

Dept. 1 One, Rivergrove Building • 2111 Front Street N.E., Salem, Oregon 97303

For more information circle number 271 on card



BRING YOUR STORAGE SPACE UP TO PAR

with **Utilicom**

...the new, multi-purpose utility building by Conrad

Ideal storage for golf carts, green equipment and all maintenance supplies. Utilicom provides clean, dry, safe storage all year 'round. Rent to your customers for convenient, private storage of carts, clubs.

Heavy gauge, overlapping roof and side panels make Utilicom the strongest, sturdiest building of its kind you can buy. Made of all galvanized metal . . . won't rust. Features full 7½' x 10' floor space, 70" entrance, 7¼' height. Big 57", free-sliding double doors (available on end for golf carts). Available in tan, white,

maroon and blue-green. All bolted construction makes Utilicom easy to assemble.

Send coupon to Conrad, Inc., Dept. GD-2, Dealer Inquiries Invited.
Houghton, Iowa, 52631 for complete information and name of nearest dealer.

Name _____
Address _____
City _____
State _____ Zip _____

For more information circle number 223 on card

A surfactant is a surface active agent which increases the ability of a liquid such as water to moisten a solid substance such as a plant surface. Basically, a surfactant acts by modifying the surface tensions which exist between water and plant surface. The resultant net reduction in inter-facial tensions causes improved wettability.

Contact herbicides such as paraquat are most commonly used on turfgrass areas for complete kill or "burn off" of the turfgrass shoots and above ground plant parts of weeds just prior to renovation overseeding. Paraquat has proven to be one of the better contact herbicides for the nonselective control of weedy turfgrass vegetation. In the case of annual bluegrass, kill of individual plants is more complete than with the other commonly available contact herbicides. In using paraquat, one should be aware that it can move laterally from the area of application in surface drainage water and thus cause kill to adjacent vegetation where a direct paraquat application has not been made. This most commonly occurs for a 14-day period following application when an intense rainfall or when irrigation has occurred.

Southern chinch bug, a new pest of St. Augustinegrass in Southern California. F.S. Morishita, R.N. Jefferson and L. Johnston. 1969. *California Turfgrass Culture*. 19(2): 9-10. (from the Department of Entomology, University of California at Riverside, Riverside, Calif. 92502).

This paper summarizes results from seasonal population studies and insecticide control experiments with southern chinch bug (*Blissus insularis* Barber) on St. Augustinegrass turfs.

Studies conducted over the growing season showed that a major population build-up began during the latter part of March and declined substantially during late October. Two distinct generations of the insect developed in early May and late July. Although the nymphs were difficult to see, they constituted the dominant portion of the population counts made. In contrast, the active crawling adults were readily visible, particularly in turfs which had been severely damaged during the late summer period.

(Continued on page 40)



One type of "water hazard" Certain-teed pipe can eliminate!

Scorched, hard greens and fairways don't make for great golf games. An irrigation system can solve the problem.

There are two very important things you need to keep in mind when you plan an irrigation system. 1. Find a good contractor. 2. Use Certain-teed Pipe. Why Certain-teed? For one thing, we make both asbestos-cement and polyvinyl chloride pipe—A-C for large diameters and PVC for small. A-C and PVC pipe are compatible with one another and with other piping systems—tying in with old systems or expanding the system in the future present no complications. Certain-teed PVC pipe is white—white pipe is more dimensionally stable than colored pipe and easier to handle on the job. Our PVC pipe offers the design flexibility of either using a deep-socket solvent weld

joint or a unique leak-proof gasket coupling. Furthermore, you get the whole system from one source—deliveries are coordinated and everything works together. When it comes to an irrigation system, remember the two important points: 1. Get a good contractor. 2. Look for the bid that specifies Certain-teed pipe.



Piping and Plastics Industries

Certain-teed Products Corporation
Valley Forge, Pa. 19481
(215) 687-5000

For more information circle number 193 on card

Three granular insecticides plus an untreated check were evaluated for the control of southern chinch bug. The three insecticides used were diazinon, ethion and Akton. All three were applied at the rate of four pounds of active ingredient per acre. The turf received an extended irrigation following the application. The insecticide treatments were made in mid-June. Dursban was also included in a comparable, second study.

The diazinon, ethion and Akton provided southern chinch bug control for a period of two months. In addition to utilizing an insecticide, the authors stress the importance of good turfgrass cultural practices in reducing the damage to turfs because the pests appeared to attack turfs which were in a more weakened condition.

Comments: The southern chinch bug is thought to have been introduced into the Whittier area of Southern California in 1967 and has subsequently become a problem on St. Augustine-grass lawns in that area. The southern chinch bug develops from white,

translucent eggs having a red spot. The immature nymph stages have two distinct colorations: the earlier instars are brilliant red whereas the last instar is distinctly black. The adult, crawling form of the insect is of two distinct types. There are long-winged and short-winged forms: both are black with white wings.

Damage caused by the southern chinch bug appears as large irregular yellowish to brown patches. Detailed examination of the damaged area frequently reveals large concentrations of the insect which feeds by sucking juices from the turfgrass plant.

Diazinon and ethion, both organic phosphates, have proven quite effective for chinch bug control over the years. In utilizing these materials, one should be aware that the former has a comparatively high human toxicity rating whereas the latter ranks as comparatively less toxic. All insecticides should be utilized judiciously as needed to control a potentially damaging insect infestation. □

Other papers of interest:

1. Cytogenetic studies on

Cynodon L.C. Rich. (Gramineae). J.R. Harlan, J.M. J. de Wet, K.M. Rawal, M.R. Felder, and W.L. Richardson. 1970. *Crop Science*. 10(3):288-291. (from the Agronomy Departments, University of Illinois and Oklahoma State University.)

2. *Post-harvest drying rate and germination of Kentucky bluegrass seed.* C.L. Canode, A.G. Law and J.D. Maguire. 1970. *Crop Science*. 10(3):315-317. (from the Department of Agronomy, Washington State University, Pullman, Wash. 99163).

3. *Performance of crownvetch and selected cool season grass on roadside backslopes as affected by slope exposure and nitrogen fertilization.* A.E. Dudeck and J.O. Young. 1970. *Agronomy Journal*. 62(3): 397-399. (from the Horticulture and Forestry Department, University of Nebraska, Lincoln Neb. 68503).

4. *Sources of variation in Cynodon Dactylon (L.) pers.* J.R. Harland and J.M.J. deWet. 1969. *Crop Science*. 9(6): 774-778. (from the Department of Agronomy, University of Illinois, Urbana, Ill. 61801).

HELLO

PO-SAN^(TM)

Good-bye FAIRWAY Poa-Annua!



You can get rid of fairway Poa-Annua by stopping seedhead development and foliar growth with PO-SAN^(TM). Fairways will have more open turf! PO-SAN leaves no phytotoxic residues in soil—allows overseeding immediately with desirable perennial grasses for exceptional germination and development. One PO-SAN treatment lasts 30-45 days.

PO-SAN also eliminates clover, dandelions and other broadleaf weeds. Stock up on PO-SAN today! See your Mallinckrodt distributor.

PO-SAN^(TM)

Mallinckrodt

MALLINCKRODT CHEMICAL WORKS
ST. LOUIS
Jersey City • Los Angeles • Montreal

For more information circle number 259 on card