



The tournament's tomorrow and Harry's forecasting a cloudburst.


Harry's the club's groundskeeper and a fine weather forecaster, too. Right now he's calling for rain even though the only thing as big as his hand in the sky is his hand.

Because Harry's also the club's rainmaker.

And by pushing the go button on a Certain-teed irrigation system, Harry can order up just the right soak needed to make the ground sparkling and ready for the next day's play.

If you're charged with keeping a brown-out out of the greens, Certain-teed can make everything pushbutton simple for you with a PVC/AC irrigation system. "PVC" and "AC" stand for polyvinylchloride plastic pipe and asbestos-cement pipe. Each does its own thing best: PVC for small, AC for large pipe diameters. Both are tough, rust- and corrosion-proof, and easy to install. You get FLUID-TITE® or deep-socket solvent-weld joints that speed connecting and never leak.

Whatever's best for your greens, you can get it all from one source—Certain-teed. Which means coordinated deliveries, good service, and everything working together. Ask your contractor to put Certain-teed into your irrigation plan specs. You'll never have another game cancelled for lack of rain.



**Pipe & Plastics
Group**

CERTAINTEED

A DIVISION OF CERTAIN-TEED PRODUCTS CORPORATION

Valley Forge, Pa. 19481

The Reliable Rain Makers

DEVELOPING A GOLF COURSE?



The easiest way to wreck a golf club is leave too many stones unturned



Greens are meticulously shaped to specifications and groomed before seeding.



An array of our modern mobile equipment, ready to move to another site.

Nothing turns a golfer off quicker than ruining his clubs on fairway stones. Nothing kills play on a golf course faster than unhappy golfers with damaged woods and irons.

That's why the fast and efficient ***Bergman Method*** of golf course preparation plows deep; turns and returns the soil to bring rocks and debris to the surface; rakes and picks the stones before leveling, contouring, fertilizing and seeding, including tees and greens. Naturally we can also install your irrigation system at the same time or serve as your consulting supervisor.


These are the reasons why ***Bergman Method*** golf courses are played by more enthusiastic golfers sooner . . . why "stony" is the description of a beautiful approach shot and not a criticism of the condition of the course. And because we are a "family" company, your golf course is assured of tender, loving care.

We go anywhere

Wm. Bergman, Jr.

1335 M-15 REESE, MICH. 48757

PHONE: CODE 517 VO 8-9812



Let
them
tear
it up...

Warren's® A-20 Kentucky Bluegrass will bring it back fast!

Warren's A-20 Kentucky Bluegrass, developed for all turf purposes, has proven to be the finest grass for tees, fairways and approaches. It is rated first in disease resistance by leading turf research organizations and universities. This characteristic makes the turf easier to maintain; lessening costs.

A-20 Kentucky Bluegrass has been "performance tested", in play, on golf courses throughout the country. A-20 is a medium-bladed grass that grows upright with a more pronounced stiffness for a better lay of the ball. It thrives under short mowings down to 1/2 of an inch.

Warren's A-20 greens up earlier in the spring and stays green longer in the fall enhancing the beauty of any course. It is drought tolerant and does well in areas previously considered too far south for bluegrass.

Write for a list of golf courses now using Warren's A-20 Kentucky Bluegrass.

Available only as Sod or Sod Plugs.

Warren's TURF NURSERY

General Offices: 8400 West 111th. St., Palos Park, Illinois 60464
Phone: 312/974-3000

**WARREN'S
TURF NURSERIES**

CHICAGO
8400 West 111th. Street
Palos Park, Ill. 60464
312/974-3000

NEW YORK
R. R. 1 — Box 240-D
(Lower & Gardnerville Rds.)
New Hampton, N. Y. 10958
914/355-6162

CLEVELAND — AKRON
Celeryville & Baseline Rds.
Box 257 — R. D. 1
Plymouth, Ohio 44865
419/687-7821

INDIANAPOLIS
6300 Donnelly Rd.
Anderson, Ind. 46011
317/378-7167

MILWAUKEE
Palmyra Rd., Box 22 — Rt. 1
Sullivan, Wisc. 53178
414/593-8815

SAN FRANCISCO
Cordella & Chadbourne Rds.
Box 347X R. R.
Suisun City, Calif. 94585
707/422-5100

LONG ISLAND
690 Deer Park Ave.
Dix Hills, N. Y. 11746
516/242-3577

BOSTON (Distributor)
Mahoney's Rocky Ledge Nursery
242 Cambridge St. — Rt. 3
Winchester, Mass. 01890
617/729-5100

GREAT BRITAIN (Distributor)
Hurst Gunson Cooper Taber Ltd.
Witham, Essex
Phone Witham 3451

**WARREN'S
ASSOCIATE GROWERS**

BOISE, IDAHO
Cloverdale Nursery
2528 N. Cloverdale Rd.
Boise, Idaho 83704
208/375-5262

WASH., D.C. — BALTIMORE
Princeton Turf Farms
Route 3
Centreville, Md. 21617
301/827-8454

MINNEAPOLIS — ST. PAUL
Green-Way Turf Nursery
170 125th. Ave. N. W.
Anoka, Minn. 55303
612/755-1319

OMAHA
Long Creek Nursery
5910 N. 52nd. St.
Omaha, Nebraska 68104
402/453-9586

WICHITA, KANSAS
M. Meyer & Son Nursery
5439 S. Hydraulic
Wichita, Kansas 67216
316/524-3413


KANSAS CITY
Princeton Turf of Kansas City
(Latta-Scholes, Inc.)
Box 9268 — Riverside Station
Kansas City, Mo. 64168
816/741-4800

PHILADELPHIA
Princeton Turf Farms, Inc.
P. O. Box 245, Old York Rd.
Hightstown, N. J. 08520
609/448-9300

BUFFALO
Country Side Turf
7229 Plank Rd.
Lockport, N. Y. 14094
716/434-5338


COLORADO
Reynolds Turf Farms
Box 595
Brighton, Colorado 80601
303/659-5760

PUTTING THE DAMPER ON WINTER DAMAGE



Water is a key to preventing winter desiccation, but how to retain it over the cold, dry period is the problem. Here are programs tailored to individual course conditions

by DR. JAMES B. BEARD



Winter desiccation, along with direct low temperature kill and the low temperature diseases, is one of three major causes of turfgrass winter injury. Winter desiccation of turfs becomes a concern if severe injury occurs to the meristematic areas of the crown responsible for initiation of new roots and shoots or to the nodes on rhizomes and stolons that initiate new turfgrass plants. Some atmospheric desiccation occurs to turfs every winter. If the injury is only to the leaf tissue, then it is superficial and causes no permanent damage to the turf, which requires reestablishment. This type of winter desiccation is frequently referred to as "windburn."

Winter desiccation is most commonly observed on cool season turfgrasses, but can also be a problem on warm season species, such as bermudagrass. Turfgrass species or cultivars that continue to maintain shoot growth later into the fall period are more prone to winter desiccation injury. Differences have been observed among the bentgrass cultivars in terms of their proneness to winter desiccation injury. Seaside creeping bentgrass is particularly tolerant of winter desiccation under Michigan conditions.

CAUSES OF WINTER DESICCATION

The basic cause of winter desiccation may be attributed to either soil or atmospheric drought. Winter desiccation caused by soil drought most commonly occurs in the plains regions of the United States and Canada where fall and winter precipitation, both rain and snow, is very low. Turfs that are subjected to extended soil drought during the winter period may develop deep cracks caused by soil shrinkage. This condition is frequently observed on greens.

Snow fence (top), as a windbreak, enhances snow accumulation on leeward side. Winter protection plots with covers (center, left) and turfgrass appearance just after removal (right). Appearance of winter top-dressing (bottom) at 0.4 and 0.3 cubic yards per 1,000 square feet.

Atmospheric winter desiccation is a problem to varying degrees throughout North America. It may occur even though adequate soil moisture is available, because the evapotranspiration rate from the turf exceeds the water absorption rate from the soil. Conditions most commonly associated or contributing to atmospheric winter desiccation injury include (a) a restricted root system, (b) droughty atmospheric conditions, including a low relative humidity and high wind velocity, (c) soil water in a frozen, unavailable state in so far as root absorption is concerned, and (d) increased viscosity of water at low soil temperatures plus decreased root membrane permeability to inward water movement.

Atmospheric winter desiccation most commonly occurs on elevated sites that are subjected to the severe desiccating effects of winds plus increased loss of precipitation by surface runoff.

PREVENTING WINTER DESICCATION

Numerous cultural techniques have been utilized over the years to prevent or minimize the degree of winter desiccation injury, particularly on intensively maintained turfs, such as putting greens. The basic methods involved are controlling the loss of water from the turf by using some sort of barrier or by applying additional quantities of water to the turf during the winter period. The particular method that is most effective in a given location of the country will be determined by (a) the extent of

golfing activity during the winter period, (b) facilities and availability of water to irrigate the area, (c) severity of desiccating conditions anticipated, and (d) budget available for use in preventing winter desiccation injury.

Synthetic Protective Covers. A winter protection cover should be considered for use on turfs subject to (a) low amounts of winter precipitation, (b) severe drying winds, (c) minimal snow accumulation, (d) negligible winter use, and (e) a history of serious desiccation injury. A number of synthetic protective covers have been developed recently for use on critical turfgrass areas, such as greens and tees. Characteristics desired in a synthetic winter protection cover include:

1. Protection against winter desiccation by functioning as a barrier to soil moisture loss;
2. Insulation against low temperature extremes that can cause turfgrass injury;
3. Partial light penetration that permits early spring green-up of the turf;
4. Sufficient temperature insulation in the spring so that excessive shoot growth does not occur;
5. A minimal "greenhouse effect" that may result in producing lethal heat buildups.

A comparison of five winter protection covers is presented in Table 1. The appropriate winter protection cover should be selected for a particular site depending on the particular problems encountered and the characteristics that best suit these needs.

continued on page 48

TABLE 1. Comparisons among five winter protection materials for use on putting greens and similar intensively cultured turfs.

CHARACTERISTICS COMPARED	TYPE OF DESICCATION PROTECTION COVER				
	Conwed Winter Protection Blanket®	Polyethylene 4-6 mil	Saran Shade® 94%	Soil Retention Mat®	Soil top-dressing (0.4-cu yd/1,000 sq ft)
Desiccation protection*	Good	Very Good	Good	Medium	Medium
Low temperature protection*	Good	Poor	Fair	Good	Poor
Rate of spring green-up*	Good	Very Good	Good	Good	Medium
Degree of shoot growth control*	Good	Poor	Good	Good	Good
Difficulty in timing removal**	Medium	High	Medium	Medium	Low
Wind displacement proneness**	Low	High	Low	Low	Very low
Lethal heat build-up potential**	Low	High	Low	Low	None
Winter disease proneness**	Medium	High	Medium	Medium	Low
Stability for reuse*	Medium	Good	Good	Poor	None

*Order of comparative rankings (Very good, good, medium, fair, poor)

**Order of comparative rankings (High, medium, low, very low, none)

See a Hahn distributor for the versatile Tournament Triplex... and a whole lot more!

The rapidly growing line of Hahn equipment can serve your mowing and turf maintenance needs from tee to green.

Mowing equipment

Our premier performer is the Tournament Triplex greensmower. So good, we guarantee it will cut equal to, or better than any other. Even on the toughest-to-mow bent greens. Attachments do four separate jobs: greens-mowing, utility mowing, de-thatching and spiking.

We are also exclusive U.S. distributors of Ransomes mowing equipment, the first choice of England's finest courses. Cut fairways and roughs with a 5- or 7-unit Hahn-Ransomes Versa-Cut hydraulic gang mower. Or, mow large areas fast with Hahn-Ransomes 3-point hitch-mounted or tractor-towed gang mowers. Use in gangs of 3, 5, 7 or 9. Choose from 16" push-type or 20" motorized Hahn-Ransomes walk-behind greensmowers, too. And a 24" Hahn-Ransomes utility riding or walk-behind reel mower!

Coming soon: Flexamatic, a five-gang specialty power-driven reel mower, with individual reel lifts and cable drive—available in the Spring of 1973.

Turf maintenance equipment

Aerifying equipment has long been one of our specialties. Choose from Grasslan and TM tractor-mounted Aerifiers,[®] walk-behind "Junior" and Verifier models, or the popular Aeri-Boy.

Remove thatch, fast, with a Hahn Verti-Cut.[®] Use a Hahn Aeri-Spiker to relieve surface compaction and prune roots. Mat in top dressing quickly, effectively with a Hahn Power Drag.

You'll also find the widest selection of ground-engaging tools for turf maintenance equipment at your Hahn Turf Products distributor. Ask him for a demonstration. Soon.





If you'd like to be a Hahn turf maintenance equipment distributor, contact Hahn, Inc., Turf Products Division, manufacturer of Hahn-West Point equipment and exclusive U.S. distributor of Hahn-Ransomes equipment. Dept. G-102 S



Hahn

Hahn, Inc., Turf Products Division • 1625 N. Garvin • Evansville, Indiana 47717

For more information circle number 176 on card



DESICCATION from page 45

Installation of the winter protection cover should be done after turfgrass growth has ceased and a high state of winter hardiness achieved, but before the soil freezes and the first permanent snowfall occurs. The following steps should be followed when installing a cover:

1. Remove all clippings and debris from the turf;
2. Apply the appropriate fungicide to prevent low temperature disease development;
3. Install the cover by securing with five-inch wire staples or with wooden laths placed at 20-foot spacings over the cover and along the (two-inch) overlapping edges. Spikes can be driven through predrilled holes in the lath and into the underlying soil. The cover should be drawn tight prior to nailing down the lath to minimize wind effects;
4. It may be necessary to place barriers around turfs covered with a winter protection cover in order to minimize damage by snowmobiles, skiers or other individuals involved in winter activities.

One additional concern in the use of winter protection covers is to recognize the importance of ensuring that the turf area has adequate surface and subsurface drainage. This is particularly important in areas that are also subject to low temperature kill. Winter injury cannot be prevented by the use of covers if the area will be subjected to periodic winter flooding followed by rapid drops in temperature to below 20° F. In this case, the potential for direct low temperature kill is quite high unless adequate surface drainage and internal soil drainage are provided.

One of the most difficult decisions in effective utilization of synthetic winter protection covers is the proper timing of removal in the spring. Success will no doubt be increased as experience is gained. General guidelines concerning removal include the following:

1. Climatic conditions should be such that there is a relatively low probability of severe desiccation or low temperatures occurring;
2. The turf be dark green without an excessive quantity of shoot growth;
3. The timing of removal is dic-

tated by the quantity of leaf growth that has occurred. The cover should be removed early enough so that the initial mowing does not scalp the turf. The first mowing after removal of the cover should be delayed for at least three days to permit hardening of the turf and to improve the initial mowing quality.

Topdressing. A topdressing applied in late autumn, but not matted into the turfgrass surface can be utilized for winter desiccation protection. The most effective topdressing rate under Michigan conditions has been between 0.3 and 0.4 cubic yard/1,000 square feet (page 45). The soil mix used in winter topdressing should be the same as that on the green to prevent a soil layering problem. Topdressing offers the advantage of being one of the cultural practices frequently used in smoothing greens as well as in preventing excessive thatch accumulation. A late fall application that is not matted in will serve the additional benefit of protecting against winter desiccation. Thus, the cost for this type of practice is relatively low. There is also no problem in timing the spring removal and few difficulties in removal as are encountered with the mulch-type materials. However, topdressing has not proven as effective a winter protection as some of the synthetic protective covers listed in Table 1. It has almost no benefit from the standpoint of low temperature insulating effects. As is the case with windbreaks that enhance snow accumulation, the winter topdressing practice should not be utilized until a preventive fungicide has been applied to protect against the low temperature diseases.

Mulches. In areas where an adequate snow cover can not be guaranteed, with or without the use of windbreaks, the use of alternative and locally available materials, such as straw, pine needles or a similar organic mulch is a possibility. This has been one of the older practices that can be used in preventing winter desiccation.

Windbreaks. Windbreaks can be utilized to minimize the desiccating effects of strong prevailing winds. An additional benefit in areas where snowfalls occur is the

enhancement of snow accumulation on the leeward side of windbreaks; thus providing even greater protection against desiccation. The windbreaks may involve (a) planting a permanent vegetative screen of trees and shrubs, (b) strategic placement of snow fence during the late fall or early winter period or (c) piling brush or tree limbs on the site during the late fall period. Increased severity of low temperature fungi may occur when utilizing windbreaks to enhance snow accumulations on greens. Thus, steps should be taken to ensure that an adequate preventive fungicide program has been instituted prior to the first snowfall. Using a snow fence to surround greens has the additional benefit of preventing potential damage from winter sports activities.

Irrigation. Winter irrigation of dormant or semidormant turfs is most commonly practiced in those portions of the country where the winters are less severe. Under these conditions it may not be possible to utilize windbreaks, mulches or synthetic protective covers because of periodic winter golfing activity. The alternatives under this condition are to correct the desiccating conditions by applications of water or to establish a temporary green and place the appropriate cover or windbreak on the permanent green in order to ensure an adequate turf for the prime golfing season.

Dormant turfs that have been without a snow or organic protective cover for an extended period of time may have to be irrigated to prevent winter desiccation. If the soil and environmental conditions do not permit opening up of the irrigation system, the alternative is to haul water to the site by tank trucks or similar equipment.

Additional irrigation practices that will assist in minimizing winter desiccation by soil drought include ensuring that the turf is adequately irrigated in late fall before the soil freezes. The soil moisture level should be near field capacity at the time the soil freezes or the first permanent snowfall occurs. Saturated soil conditions should be avoided, because this will increase the proneness to low temperature kill or winter diseases. □

Golf course superintendents and other turf experts throughout the country rely on these outstanding Chemagro Blue Bullseye chemicals to keep their grass in first-class condition throughout the year:

***DYRENE** 50% Wettable Powder turf fungicide provides outstanding protection against leaf spot, melting-out, copper spot, rust, *Sclerotinia* dollar spot, brown patch and snow mold. Will not clog or corrode spray equipment. No unsightly deposit. Will not stain fabric or shoes when dry.

***DASANIT** nematicide provides fast contact and long-lasting residual control of nematodes. Merely apply with a granular applicator and drench the soil with water.

***DYLOX** 80% Soluble Powder insecticide provides outstanding control of sod webworms. Three spray applications at monthly intervals are recommended.

***BAYGON** 70% Wettable Powder insecticide provides economical, long-lasting control of certain destructive insects attacking turf. Sprays are easy to mix and apply.

***DEXON** 35% Wettable Powder fungicide is unsurpassed for long-lasting control of *Pythium*. Will not cause leaf burn when applied as directed. Compatible with other turf pesticides.

Contact your Chemagro supplier *now* for full information about these time-tested turf chemicals.




KANSAS CITY, MISSOURI 64120


Everybody has turf problems. Chemagro has turf solutions.



7278



BOSTON '73



You are cordially invited to attend the GCSAA 44th Annual International Turfgrass Conference and Show in Boston, Massachusetts, January 7-12, 1973.

Good turf is just as important as good golfing techniques in keeping scores low. This is your once-a-year opportunity to learn how to make your club a real winner.

At this year's Annual GCSAA Conference over 50 well-known experts will set a busy pace for the 4½-day educational program. Golf course superintendents will hear about the latest turfgrass developments and management practices, participate in discussions on timely subjects of the profession, receive reports on practical day-to-day, problem-solving solutions.

You'll also find answers firsthand to new time- and labor-saving equipment directly from the suppliers. More than 100 exhibitors will display the latest and best in materials, machines and supplies.

More important, this is the unique occasion when you can get together with leading golf superintendents from the top clubs to compare notes, exchange ideas, "talk shop."

44th INTERNATIONAL TURFGRASS CONFERENCE AND SHOW

JOHN B. HYNES CIVIC AUDITORIUM - BOSTON, MASS.

JANUARY 7-12, 1973

Sponsored by the

GOLF COURSE SUPERINTENDENTS ASSOCIATION OF AMERICA

3158 Des Plaines Avenue, Des Plaines, Illinois • 312/824-6147

Registration Fee:

GCSAA Members \$15 • Nonmembers \$35 • Ladies \$15

Send in for your registration form now. By registering in advance you will save time at the registration desk in Boston, where a distinctive badge and your other conference materials will be awaiting your arrival.

Last name First Name

Mailing Address

City & State Zip Code

Club or Firm