



Seeds of Change.

Introducing a series of seed blends that produce the highest quality turf...from Scotts®/Landmark, the industry's most widely known and respected turf science expert. Double Eagle® brings you—

- Blends or mixtures that contain NTEP highly rated varieties
- Available in Platinum Tag® quality—the highest standard of purity in the industry
- Available in Premium Tag® quality – a higher standard than Blue Tag certified
- Superior performance in any application
- Superior turf with excellent color
- Rapid establishment, tolerance to low mowing and excellent disease resistance
- Unmatched technical support. From now on, think Double Eagle. Because no other seed brand can benefit you more. Contact your local Scotts or Landmark distributor for more information or contact Scotts or Landmark at 937-644-7270 or 1-800-268-2379.



DOUBLE EAGLE™
PREMIUM SEED

* Scotts Platinum Tag® is the highest standard of purity available—four times as stringent as Gold Tag sampling.

How Do Turfgrasses Use Nickel?

After all, it's the second least-abundant micronutrient

By Richard J. Hull

Nickel (Ni) is the latest element to be added to the list of nutrients that are essential for plants.

While suspected to be essential since the 1970s, Ni was experimentally demonstrated to be required for legumes in 1983 by Dr. Ross Welch and his associates of the USDA Plant, Soil and Nutrition Laboratory at Cornell University (Eskew et al., 1983). Four years later, the same investigators reported Ni to be generally essential for all plants (Brown et al. 1987). In 1992, the Agricultural Research Service officially added Ni to the list of plant-essential elements.

Although Ni is the last essential nutrient element to be discovered, it is not really the least abundant in plants. Plant leaves require approximately .1 mg Ni/kg dry tissue in parts per million (ppm), a little more than the amount of molybdenum (Mo) required.

To put this in relative terms, for every atom of Mo required by plants, two atoms of Ni and one million atoms of nitrogen are needed (Hull, 2003). Clearly Ni is a micronutrient.


I am aware of no Ni analyses having been conducted for turfgrass leaves but like Mo, they would likely average about 2.5 ppm. In most plants, the Ni concentration is in the range of 1 ppm to 10 ppm of dry tissue (Marschner, 1995). Given those values, the Ni content would be sufficiently high to preclude any possibility of deficiency symptoms in field-grown plants.

The low probability of Ni ever being of concern to the turfgrass manager may explain the virtual absence of published research on Ni in the turf literature. Since Ni is an essential part of an enzyme required for the proper metabolic cycling of nitrogen (N) in all plants, it should spark at least indirect interest among turfgrass professionals.

Ni: essential component of urease

The only known function of Ni in higher plants is its role in the enzyme urease. There it is built into a complex protein structure by becoming coordi-

ated between three N and three oxygen (O) atoms. During enzymatic activity, one of the oxygen atoms is displaced by a water molecule which is then inserted into a urea molecule, hydrolyzing it into two ammonium ions (NH_4^+) and a carbon dioxide (CO_2).



The primary purpose of the urease enzyme is to recycle the N in urea that is a waste product of N metabolism. Urea is a byproduct of several biochemical pathways through which N is cycled during biosynthetic and degradation processes (Figure 1). These include the ornithine cycle by which the essential amino acid arginine is made and metabolized into probably the greatest source of metabolic urea.

Metabolism of the ureides, allantoin and allantoic acid, that in many plants are transported from roots to leaves through the xylem, produces urea when their N is assimilated into proteins. Ureides can also be derived from the recycling of nucleic acids when purines are metabolized.

In short, urea is an inevitable product of normal nitrogen metabolism that if allowed to accumulate in plant tissue will cause substantial damage. To prevent such damage and reutilize its N, urea is hydrolyzed by urease and the resulting NH_4^+ is reassimilated in the same way as that derived from nitrate metabolism (Hull, 1996). Since plants lack the elimination system of animals by which urea can be released in urine and its N wasted, plants have evolved a more conservative approach to urea removal and N recycling. To do this, Ni is essential.

Ni deficiency symptoms can be linked directly to the failure of urease to eliminate urea. The initial deficiency symptom is leaf tip burn resulting from urea accumulation in this oldest portion of a grass leaf.

Injury probably occurs as a salt-like effect as urea accumulates to high levels. Root and shoot growth is suppressed, and leaves lose their green

Continued on page 54



QUICK TIP


Nothing is more important to your chemical applications than actually applying the amount you intended. The EPA requires it, your course and community counts on it. Toro can help enhance your control. Multi Pro® sprayers feature the industry's most accurate control system that delivers a constant rate of application regardless of speed. In addition, check with your Toro distributor for sprayer inspection and calibration services or visit toro.com/calibrate for a useful tool to help accurately calibrate any sprayer.

Just because you use liquid fertility, don't assume you're getting True Foliar nutrition.

Top 10 reasons to use the CPR— NutriRational™ True Foliar Nutrition Program

1. Increase turf quality and density.
2. Increase root mass and depth.
3. Increase resistance to disease especially anthracnose.
4. Increase stress tolerance.
5. Reduce thatch accumulation.
6. Increase nutrient absorption.
7. Reduce clipping removal and loss of applied fertilizer.
8. Reduce nutrient leaching.
9. Increase photosynthesis and phytochemical efficiency.
10. Reduce potential for diseases associated with excess Nitrogen.

The CPR-True Foliar Program provides the complete, balanced nutrition and seaplant extract necessary for turf physiological fitness. The program produces controlled, consistent growth necessary to meet today's demands for speed and quality.




Emerald Isle Solutions™



MILLIKEN

(800) 845-8502 www.millikenturf.com

TABLE 1

Sources of Urea in Plants and N_4H Assimilation

Continued from page 52

color due to interveinal chlorosis and eventually necrosis (tissue death). In younger leaves, the tip region fails to unfold properly. Because seed germination is accompanied by rapid nitrogen metabolism, the seeds from Ni-depleted plants often fail to germinate likely because of urea accumulation. These observations have been made on laboratory-grown plants since no Ni deficiency has ever been reported in the field.

Urease key to N fertilizer

Urease allows urea to be used as N fertilizer. Urea contains two N atoms per molecule (40 percent N), is easily and inexpensively manufactured and is safe to handle, making it a popular source of fertilizer N. However, when applied to soil or plant leaves, urea is not absorbed directly but rather hydrolyzed to NH_4^+ and CO_2 with the NH_4^+ , then absorbed by plant cells and assimilated into amino acids.

It is urease derived from the plant or soil micro-organisms that releases NH_4^+ from urea making its N available to the plant. Urease, regardless of its source, makes urea-N available to plants, and it depends on Ni for its hydrolytic function. Thus, urea is an effective N fertilizer due to the action of Ni.

Urea probably can be absorbed by plant cells because Ni-starved plants exhibit greater deficiency symptoms when urea is provided as their N source. Since urease is less active in Ni-deficient plant, any urea absorbed will accumulate within cells and reach toxic concentrations. However, urease is so abundant in the plant-soil environ-

ment, it is generally believed that urea-N is actually absorbed as free NH_4^+ or as its nitrification product, NO_3^- . If urea is applied to wet leaves or thatch under high temperatures, the rate of urea hydrolysis may exceed the rate of NH_4^+ absorption. The excess NH_4^+ may be lost as ammonia (NH_3) gas.

Several researchers have determined that as much as 40 percent of urea-N applied to turf can be lost as NH_3 when conditions are favorable for rapid hydrolysis (Turner and Hummel Jr., 1992). Such N losses can be prevented by irrigating turf immediately after applying urea.

Ni accumulated in seeds

Although absorbed from the soil solution as the divalent nickel ion (Ni^{2+}), within plant cells Ni readily combines with organic anions such as citrate and cysteine. It is likely that Ni is transported within plants as its citrate chelate. Unlike some cationic micronutrients (e.g. iron or manganese) that exhibit substantial xylem mobility but are virtually immobile in the phloem, Ni^{2+} is readily transported in both xylem and phloem (Kochian, 1991). As a result, Ni often accumulates in developing seeds to concentrations greater than most other plant organs. This is one reason why Ni deficiency symptoms were not demonstrated until recently. There can be sufficient Ni within the seeds of some annual plants to meet their needs for one or more generations. This contributes to the low probability of observing a Ni deficiency under anything but carefully controlled laboratory conditions.

Toxicity greater concern than deficiency

In many situations, Ni toxicity is a reasonable possibility especially where soils have been supplemented with sewage or industrial sludges contaminated with heavy metals.

Plants vary in their sensitivity to Ni with some capable of tolerating tissue concentrations of more than 1,000 ppm while many plants exhibit toxic symptoms when their tissues exceed 10 ppm Ni. Most soils contain less than 100 ppm Ni but only a small amount of that is exchangeable and available to plants (about 1 ppm). Much Ni in soils is present as insoluble mineral fractions and the remainder is bound into organic complexes and is sparingly available to plant roots. Thus, Ni toxicity is not normally a problem for the turf manager.



QUICK TIP

Look to The Andersons Advantage when using granular fungicides. Better carriers, expert formulations and years of experience all add up to dependable products you expect from The Andersons. For more information, contact your Andersons Golf Products distributor.

List of articles in the micronutrient use series as published in *TurfGrass Trends*

- Hull, R.J. 1996. "Nitrogen usage by turf." *TurfGrass Trends* 5(11):6-15.
- Hull, R.J. 1997. "Phosphorus usage by turf-grasses: The energy nutrient often neglected by turf managers." *TurfGrass Trends* 6(5):1-12.
- Hull, R.J. 1997. "Calcium usage by turfgrasses: The nutrient forgotten by turf managers." *TurfGrass Trends* 6(10):6-13.
- Christians, N.E. 1998. "Potassium fertilization." *TurfGrass Trends* 7(3):9-13.
- Hull, R.J. 1998. "Sulfur usage by turfgrasses." *TurfGrass Trends* 7(2):1-9.
- Hull, R.J. 1998. "Magnesium usage by turf-grasses." *TurfGrass Trends* 7(8):7-14.
- Hull, R.J. 1999. "Iron usage by turfgrasses." *TurfGrass Trends* 8(2):1-11.
- Hull, R.J. 2001. "Zinc usage by turfgrasses." *TurfGrass Trends* 10(7):7-11.
- Hull, R.J. 2001. "Manganese usage by turf." *TurfGrass Trends* 10(9):6-13.
- Hull, R.J. 2002. "Recent research offers clues to boron's purpose." *TurfGrass Trends* 11(3):11-16.
- Hull, R.J. 2002. "Copper management demands attention." *TurfGrass Trends* 11(7):12-16.
- Hull, R.J. 2002. "Copper is essential bridge for many plant functions." *TurfGrass Trends* 11(11):14-16.
- Hull, D. 2003. "Small amounts of molybdenum streamlines nitrogen uptake." *TurfGrass Trends in Golfdom* 59(1): 42-44).
- Hull, R.J. 2003. Ubiquitous chlorine performs vital tasks in turf. *TurfGrass Trends in Golfdom* 59(3):58, 60-62).

In the coastal mountains of California, serpentine soils can be found that are enriched in several heavy metals including iron, magnesium, chromium (Cr), cobalt and Ni, the latter being present at concentrations as high as 250 ppm exchangeable metal (Mengel and Kirkby, 1982). Such soils are inhospitable to most plants largely because of the magnesium content being greater than calcium (Ca/Mg about .4) but also because of Ni and Cr being present in toxic concentrations. Serpentine soils are not suitable for turfgrass growth but the Ni toxicity can be alleviated by lime applications that neutralize the soil acidity.

At higher pH values, both Ni and Cr become less soluble, making them less available for root absorption. Lime also increases the Ca/Mg ratio and may stabilize membrane transport proteins which increase the selectivity of ion uptake by roots. Those plants that can tolerate serpentine soils absorb Ni less rapidly and concentrate it to levels in excess of 200 ppm but as citrate and malate chelates that are sequestered within vacuoles and rendered less toxic. Some hyperaccumulators can achieve tissue Ni concentrations of more than 10,000 ppm again mostly as a chelated form. Other defensive measures likely are also employed by Ni-tolerant plants.

Hull is a professor of plant sciences at the University of Rhode Island in Kingston, R.I., who specializes in plant nutrition.

REFERENCES

- Brown, P.H.; R.M. Welch and E.E. Cary. 1987. "Nickel: a micronutrient essential for higher plants." *Plant Physiology* 85:801-803.
- Eskew, D.L.; R.M. Welsh and E.E. Cary. 1983. "Nickel: an essential micronutrient for legumes and possibly all higher plants." *Science* 222:621-623.
- Hull, R. 1996. "Nitrogen usage by turf." *TurfGrass Trends* 5(11):6-15.
- Hull, R. 2003. Small amounts of molybdenum streamlines nitrogen uptake. *TurfGrass Trends* 59(1): 42-44).
- Kochian, L.V. 1991. "Mechanism of micronutrient uptake and translocation in plants." *In Micronutrients in Agriculture*. (J.J. Mortvedt, ed.), pp. 229-296. *Soil Sci. Soc. Am. Book Series* No. 4, Madison, Wis.
- Marschner, H. 1995. *Mineral Nutrition of Higher Plants*, 2nd edition. Academic Press, San Diego. pages 364-369.
- Mengel, D. and E.A. Kirkby. 1982. *Principles of Plant Nutrition*, 3rd Edition. pp. 564-565. International Potash Institute, Bern, Switzerland.
- Turner, T.R. and N.W. Hummel, Jr. 1992. "Nutritional requirements and fertilization." pp. 385-439. *In* D.V. Waddington, R.N.
- Carrow and R.C. Shearman (eds.). "Turfgrass, Chapter 11." *Agronomy Monograph series* No. 32, ASA, CSSA & SSSA, Madison, Wis.

TURFGRASS TRENDS

SECTION STAFF

Managing Editor

Curt Harler
440-238-4556; 440-238-4116 (fax)
curt@curtharler.com

Golfdom Staff Contact

Frank H. Andorka Jr.
440-891-2708; 440-891-2675 (fax)
fandorka@advanstar.com

Online Editor

Lynne Brakeman
440-826-2869; 440-891-2675 (fax)
lbrakeman@advanstar.com

Senior Science Editor

Dr. Karl Danneberger
614-292-8491; 614-292-3505 (fax)
danneberger.1@osu.edu

Production Manager

Jill Hood
218-723-9129; 218/723-9223 (fax)
jhood@advanstar.com

Art Director

Lisa Lehman
440-891-2785; 440-891-2675 (fax)
llehman@advanstar.com

Publisher

Patrick Jones
440-891-2786; 440-891-2675 (fax)
pjones@advanstar.com

Group Publisher

Tony D'Avino
440-891-2640; 440-891-2675 (fax)
tdavino@advanstar.com

Corporate & Editorial Office

7500 Old Oak Blvd.
Cleveland, OH 44130-3369

FIELD ADVISORS

Rob Anthony
Southern Methodist University

J. Douglas Barberry
Turf Producers International
Agronomist

F. Dan Dinelli
North Shore CC

Merrill J. Frank
Columbia CC

Michael Heacock
Pacific Golf Management K. K.

Paul B. Latshaw
Muirfield Village CC

Kevin Morris
National Turfgrass Evaluation
Program

Sean Remington
Green Valley CC

Ken Schwark
Roddy Ranch GC

Matt Shaffer
Merion GC

Wayne Horman
The Scotts Co.

Eric Kalasz
Bayer Environmental Sciences

David Irmen
The Andersons

Chris Byrd
Milliken

Van Cline
The Toro Co.

EDITORIAL REVIEW BOARD

Dr. A.J. Powell
University of Kentucky

Dr. Eliot C. Roberts
Rosehall Associates

Dr. Gerald Horst
University of Nebraska

Dr. Eric Nelson
Cornell University

Dr. Richard Hull
University of Rhode Island

Dr. Vic Gibeault
University of California

Dr. Pat Vittum
University of Massachusetts


Dr. Rick Brandenburg
NC State University

CONTACT US:

Editorial: 440-238-4556

Web site: www.turfgrasstrends.com

I will never say a riding greensmower
I will never say a riding greensmower
I will never say a riding greensmower
I will never say a riding greensmower
I will never say a riding greensmower
I will never say a riding greensmower
I will never say a riding greensmower
I will never say a riding greensmower
I will never say a riding greensmower
I will never say a riding greensmower



The new Toro® riding greensmower cutting unit.

With its die-cast design, easy dual-point adjustment, and bedbar
an ideal cut, as low as 1/16 inch, time after time. Contact your

can't cut as well as a walk behind again.
can't cut as well as a walk behind again.
can't cut as well as a walk behind again.
can't cut as well as a walk behind again.
can't cut as well as a walk behind again.
can't cut as well as a walk behind again.
can't cut as well as a walk behind again.
can't cut as well as a walk behind again.
can't cut as well as a walk behind again.
can't cut as well as a walk behind again.
can't cut as well as a walk behind again.

Forget what you've learned.

pivot location, the new cutting unit for Toro riding greensmowers maintains distributor at 1-800-803-8676 or visit toro.com for Toro Financing™ options.



Count on it.

LaFoy, Hurdzan and others discuss the state of the industry, the impact of technology and other subjects

Architects Weigh In

BY GEOFF SHACKELFORD

Golfdom informally polled a variety of architects by e-mail and asked their thoughts on the state of the golf industry, the impact technology is having on restoration projects and what the architects would do to put the distance fire out. Here's what they said:

Architects on the state of the golf industry and other comments


John LaFoy: "When I think of the golf industry in recent years, I think of Alan Greenspan's comment about the stock market during much of that same period — 'irrational exuberance.' Not that I blame anyone or anything for being exuberant about their profession, but at some point you have to wonder if things were not going a little too fast. My personal observation is that the golf industry is still strong in the renovation and remodeling sectors. The biggest difference is that many of the architects and contractors that once were not interested in this type of work are now very interested in it."

Michael Hurdzan: "I still don't think the entire golf industry (clubs, shoes, carts, resorts, clothing, instructors, etc.) realizes that golf


Continued on page 60

Architects are worried that some classic courses, like Merion Golf Club, have become outdated because they don't have room to expand.

HOW PREDICTABLE IS YOUR ORGANIC FERTILIZER?



UNIVERSITY RESEARCH



RESULT

On average, over the two year study, 85% of the nitrogen in Nature Safe 8-3-5 released in a 10 week period versus 53% for sewage sludge and less than 30% for the manure product.

Research conducted by North Carolina State University, Dr. Charles Peacock



An organic fertilizer's nutrient value and release characteristics are determined by the amino acid content of the ingredients and their related digestibility. The fish, meat, bone, blood and feather meals in Nature Safe have the highest amino acid concentration of any commercial fertilizer, delivering exceptional nutrition in every bag.

Analysis: **Nature Safe 12-2-0**
Amino Acid Content: **72.43%**

vs.

Analysis: **Sewage Sludge 6-2-0**
Amino Acid Content: **29.60%**

RESULT:

Nature Safe has twice the nutrient value with approximately 2 1/2 times the amino acid content.

Nature Safe is specifically formulated to ensure product consistency and nutrient availability, resulting in an extremely predictable release. The USDA approved ingredients are part of the food chain and contain vitamins, enzymes, sugars and carbohydrates along with chelated phosphorus and calcium for a balanced feed. In the adjacent North Carolina State study, Nature Safe released 60% more available nitrogen than the sewage sludge product and almost three times more than the manure-based fertilizer, confirming Nature Safe's unparalleled fertilizer efficiency.

For the most consistent and predictable organic fertilizer, find out why leading superintendents make **Nature Safe their Natural Choice for Optimum Soil and Plant Nutrition!**

Nature Safe

Natural & Organic Fertilizers

Call (800) 252-4727 or on the web at www.naturesafe.com

Continued from page 58

and the environment is everybody's problem. The entire industry needs to support those who are trying to fund the research to show that golf courses can be even better neighbors to Mother Nature. Somehow they see it as the superintendent's problem alone.

With the water shortage in many parts of the country, golf courses will be one of the first to be cut off, denied permits or pay a high price in one form or another, with the result being fewer golf courses or at least higher-priced golf. Tied to that is the need to keep golf affordable and accessible, and here again most of the industry sees it as someone else's problem, which it's not. We all need to support the growth of the game, make it a family-friendly sport and be aware of its overall image."

Tim Liddy: "The length issue is getting tiresome, but it's a big problem, better stated by others like Jack Nicklaus. Only Augusta National has the ability to require a tournament ball, and I hope it does this year. Golf is going the way of tennis and becoming boring to watch. The talented golfer has also lost his advantage with the new technology. Just look at the recent major winners."

John Fought: "The more I work in the design business, the more I appreciate timeless, classic works from the Golden Age. Why do architects feel the need to invent new gimmicks and create loads of courses that neither

Architect Bob Cupp says golf needs more players who play because they "love the sight of the ball against the blue sky or backdrop of trees – not because they want to thump on their chests."



“The entire industry needs to support those who are trying to fund the research to show that golf courses can be even better neighbors to Mother Nature.”


— MICHAEL HURDZAN

look natural nor play well? I find myself constantly looking back and studying."

Tom Johnson: "With regard to the current low level of new-course activity, golf is just going through a normal business phase. We overbuilt from the mid-1980s until 2000, and the cyclical nature of any industry will have to adjust. The most disturbing factor in golf is the Neanderthal reluctance of Augusta National to join the real world. It opens its door to the media and the golfing public each year, yet flaunts its own right to privacy, bigotry and gender bias each year. It's quite disgusting."

Bob Cupp: "First, it would be better for the game if it wasn't so egotistical. Ego drives the cost, and subsequently fewer people play. People should play because they love the sight of the ball against the blue sky or a backdrop of trees, and its bounce and roll toward the flag — not because they want to thump on their chests. Second, we are headed down a dangerous road right now with the equipment."

Continued on page 64



DYNAMIC GRAPHICS