

Floratine – a different breed of dog. Turf is all that we do. And turf strength is our passion. It's reflected in our scientists' designs, our raw materials selection and our representatives' recommendations of exceptional products like Carbon Power, Astron and ProteSyn. Our singular focus is meeting the physiological requirements for grass to be stronger, longer.

Different? Sure, but we think your turf will appreciate the difference.



Patented Chemistry. University Tested.



Picture 2. Shoot and root biomass of Japanese Lawngrass unaffected in response to 240, 480, 720 millimeters of aluminum.

Continued from page 60

with two controls $(0 \,\mu\text{m of Al})$ at pH 4.0 and 6.5 for six weeks under greenhouse conditions.

Treatments were arranged in a randomized complete block design with three replications. Pot dimensions were 15 centimeters (cm) (6 inches) in diameter and 12 cm (4.8 inches) in height with eight holes (0.40 inches) at the bottom to allow for drainage. For each turf species selected (Table 1), 30 single meristem shoots from 2-year old well-rooted turf were planted in 100 percent sand obtained from Golf Agronomics, with roots and shoots clipped to ensure similar length.

All turfgrasses were fertilized daily (50 milliliters [ml] to 100 ml) using a modified halfstrength Hoagland's nutrient solution [Hoagland and Arnon, 1956]. Aluminum was added to the solution as $Al_2(SO_4)_3 18H_2O$ and was adjusted to pH of 4.0 (+/-0.1). Deionised distilled water was used to make all solutions.

Moderate to high exposure

Study II was similar in design as study I, except Al was increased to $480 \,\mu\text{m}$, $960 \,\mu\text{m}$ and $1440 \,\mu\text{m}$ and P concentrations were reduced to $0.0425 \,\text{kilograms}$ (kg) per hectare per day.

Treatments were arranged in a randomized complete block design and the same 10 species were selected as in study I. Unless otherwise stated, materials and experiment duration were identical to study I.

Relative values were used for all data collected due to the inherent variability among turfgrasses selected. Relative nutrient recovery was determined and dry weight of shoots and roots were measured in both studies and used to cal-

Turfgrass Species	Scientific Name	Cultivar
Bahiagrass	Paspalum notatum Flugge.	'Pensacola'
Buffalograss	Buchloe dactyloides [Nutt] Engelm.	
Seashore Paspalum	Paspalum vaginatum Swartz.	'Isle'
Centipedegrass	Eremochloa ophiuroides (Munro) Hack.	
Common Carpetgrass	Axonopus affinis Chase.	
Common Bermudagrass	Cynodon dactylon var. dactylon	'Princess 77'
Japanese Lawngrass	Zoysia japonica Steud.	'Meyer'
Manilagrass	Zoysia matrella (L.) Merr.	
St. Augustinegrass	Stenotaphrum secundatum (Walt.) Kuntze.	'Raleigh'
Hybrid Bermudagrass	Cynodon dactylon (L.) Pers. x C. transvaalensis Burtt-Davy	'TifEagle'

Warm-season turfgrass species and cultivars selected for studies I and II.

TABLE 2						
Species	Relative Root Mass		Relative	1440		
Coorboro Pacoalum	0.70-*	0.71	0.576	0.00-bc	0.65	0 E1d
Common Carpetorass	0.99abc	1.00	0.57C	0.87abc	0.05	131a
Japanese Lawngrass	1.01abc	1.23	0.79bc	0.84bc	0.93	0.74cd
Manilagrass	1.24a	1.03	1.30a	1.03ab	1.04	0.97b
Bahiagrass	0.82c	0.94	0.71c	0.80bc	0.67	0.71cd
Common Bermudagrass	0.86bc	0.79	0.77c	0.69bc	0.59	0.62cd
Buffalograss	0.88abc	0.67	0.45c	0.55c	0.67	0.67cd
Centipedegrass St. Augustinograss	1.23aD	1.10	0.660	1.18a 0.02ab	0.94	0.7200
Hybrid Bermudagrass	0.09C	0.66	0.73c	0.72ab	0.71	0.66cd
LSD	0.38	ns	0.36	0.34	ns	0.24
p-value	0.05**	0.17	0.01	0.05	0.34	0.01

*VALUES WITHIN A COLUMN FOLLOWED BY THE SAME LETTER ARE NOT SIGNIFICANTLY DIFFERENT AT P 0.05 BY PROTECTED LSD. **SIGNIFICANCE AT THE PROBABILITY LEVEL P 0.05.

Relative root and shoot biomass based on control pH 4.0 of 10 warm-season species affected by aluminum at 480, 960 and 1440 millimeters, study II.

culate relative root and shoot mass.

Roots were extracted from the soil and thoroughly washed until all soil was removed. Once clean, roots were clipped from the base of the shoot tissue. All relative values were based on the control pH of 4.0 and calculated:

Relative root mass [RRM = (root mass with Al/root mass without Al) x 100], relative shoot mass [RSM = (shoot mass with Al/shoot mass without Al) x 100], relative nutrient recovery [RNR = (nutrient concentration with Al/nutrient concentration without Al) x 100].

Fresh weight of roots and shoots were collected and placed in an oven at 80 degrees Celsius (C) (176 degrees Fahrenheit [F]) and dried for 48 hours. Once dried, samples were weighed for root and shoot mass and analyzed by the Clemson University soil testing laboratory for P, K, Ca, Mg, sulfur (S), chlorine (Cl), zinc (Zn), copper (Cu), manganese (Mn), iron (Fe), Al and boron (B).

Relative growth parameters

All statistical computations were conducted using analysis of variance (ANOVA) within the Statistical Analysis System (SAS Institute, 1999). Means were separated by the Fisher's Least Significant Difference (LSD) test. An alpha of 0.05 was used for relative shoot and root mass and for all Al and nutrient analyses of soil and plant tissue.

No differences were found in study I for RRM or RSM. In study II, Japanese lawngrass exhibited a 23 percent increase in RRM at 960 μ m of Al, but as Al increased to 1440 μ m, a 21 percent decrease in rooting occurred (Table 2).

Carpetgrass and manilagrass produced greater RRM (15 percent and 30 percent) at 1440 µm of Al compared to the other eight selected species. Buffalograss, seashore paspalum, bahiagrass and St. Augustinegrass, however, consistently performed poorly with a 55 percent, 43 percent, 29 percent and 50 percent decrease in RRM at 1440 µm of Al, respectively. All species had decreased RSM at 1440 µm of Al, except carpetgrass (31 percent increase).

Liu et al. (1995) observed Kentucky bluegrass (*Poa pratensis* L.) cultivars enhanced root and shoot growth when exposed to 320 µm of Al.

Relative nutrient recovery

Under high Al stress in studies I and II, relative concentrations of P and Ca were significantly reduced in the root tissues (Tables 3 and 4). At 720 µm of Al, Japanese lawngrass increased relative root P

TABLE 3						Dist P	
Species	Study I 240	480	720	Study II 480	960	1440	
Seashore Paspalum Common Carpetgrass Japanese Lawngrass Manilagrass Bahiagrass Common Bermudagrass Buffalograss Centipedegrass St. Augustinegrass Hybrid Bermudagrass	0.94bc* 0.87bc 1.56a 0.77bc 0.82bc 1.09ab 0.51c 0.81bc 1.12ab 0.48c	0.81ab 0.78ab 0.56bc 0.52bc 0.66bc 0.79ab 0.35c 0.81ab 1.03a 0.34c	0.60bc 0.55bc 1.05a 0.85ab 0.65bc 0.71ab 0.28c 0.67b 0.28c 0.67b 0.75ab 0.56bc	0.71 1.17 0.99 0.74 1.26 0.90 0.79 0.66 0.91 0.71	0.57 0.66 0.72 0.57 0.87 0.87 0.87 0.57 0.47 0.55 0.43	0.54 0.66 0.65 0.47 0.85 0.77 0.60 0.47 0.64 0.43	
LSD	0.51	0.37	0.37	ns	ns	ns	
p-value	0.01**	0.01	0.04	0.43	0.30	0.66	

*VALUES WITHIN A COLUMN FOLLOWED BY THE SAME LETTER ARE NOT SIGNIFICANTLY DIFFERENT AT P 0.05 BY PROTECTED LSD

**SIGNIFICANCE AT THE PROBABILITY LEVEL P 0.05.

Relative phosphorus (%) recovery in root tissue based on pH 4.0 of 10 warmseason species affected by aluminum at 240, 480 and 720 mm, study I; and 480, 960, and 1440 mm, study II.

TABLE 4						
Species	Study I 240	480	720	Study 480	960	1440
Seashore Paspalum Common Carpetgrass	0.90 0.89	0.82 0.76	0.57bc* 0.76ab	0.40 0.79	0.41e 0.72abcde	0.33d 0.62abcd
Japanese Lawngrass Manilagrass	1.16	0.82	0.77ab 0.81ab	1.16	1.06a 0.65bcde	0.87 0.39cd
Common Bermudagrass Buffalograss	1.15	0.74 0.99 0.97	0.88a 0.84a 0.88a	1.10	1.00ab 1.06a 0.98abc	0.75abc 0.81ab 0.82ab
Centipedegrass St. Augustinegrass	1.02	0.88	0.86a 0.71ab	0.97	0.61cde 0.93abcd	0.33d 0.86ab
Hybrid Bermudagrass LSD	0.50 ns	0.59 ns	0.40c 0.24	1.03 ns	0.58de 0.38	0.52bcd 0.42
p-value	0.08	0.44	0.01**	0.06	0.01	0.05

*VALUES WITHIN A COLUMN FOLLOWED BY THE SAME LETTER ARE NOT SIGNIFICANTLY DIFFERENT AT P 0.05 BY PROTECTED LSD.

**SIGNIFICANCE AT THE PROBABILITY LEVEL P 0.05.

Relative calcium (%) recovery in root tissue based on pH 4.0 of 10 warm-season species affected by aluminum at 240, 480 and 720 mm, study I; and 480, 960 and 1440 mm, study II.

recovery 5 percent, while manilagrass minimized relative root P loss (15 percent) (Table 3).

However, a 35 percent, 72 percent and 44 percent decrease in relative root P concentration occurred for bahiagrass, buffalograss, and hybrid bermudagrass, respectively. Study I results agree with Foy and Murray's (1998b) observations where P concentrations in shoots and roots of tall fescue (*Festuca arundinacea* Schreb.) cultivars decrease when exposed to Al treatments.

In study II, relative root P concentrations were unaffected by Al levels (Table 3). This was probably because of drastic reductions in P levels for all treatments. However, most species had reduced relative root P concentration as Al treatments increased, which were similar to the results in study I.

Continued on page 64

Continued from page 63

Regardless of species, relative root Ca concentrations were reduced as Al increased in both studies (Table 4). This is consistent with observations reported by Krizek and Foy (1988). No definitive trends for K recovery in shoot or root tissues occurred in studies I or II (Data not shown).

Japanese lawngrass (Picture 2), manilagrass, and carpetgrass are relatively tolerant of Al levels. In study II, carpetgrass and manilagrass were the only warm-season turfgrass species that produced superior rooting at 1440 µm of Al with an increase of 15 percent and 30 percent, respectively. Al impacted nutrient recovery in most turfgrass species.

In both studies, as Al increased, relative Ca recovery generally decreased in root and shoot tissues. K levels, however, were unaffected by Al at any treatment level.

REFERENCES

Clarkson, D.T. 1965. "The effect of aluminum and some other trivalent metal cations on cell division in the root apices of Allium cepa." Ann. Bot. 20:309-315.

Foy, C.D., R.L. Chaney, and M.C. White. 1978. "The physiology of metal toxicity in plants". Ann. Rev. Plant Physio. 29:511-566.

Foy, C.D. and J.J. Murray. 1998a. "Responses of Kentucky bluegrass cultivars to excess aluminum in nutrient solutions." J. Plant Nutr. 21:1967-1983.

Foy, C.D. and J.J. Murray. 1998b. "Developing aluminum-tolerant tall fescue for acid soils." J. Plant Nutr. 21:1301-1325.

Hartwell, B.L. and F. R. Pember. 1918. "The presence of aluminum as a reason for the difference in the effect of so-called acid soils on barley and rye." Soil Sci. 6:259-279.

Hoagland, D. R. and D. I. Arnon. 1956. "The water-culture methods for growing plants without soil." California Agr. Exp. Sta. Circular 347pp.

Kochian, L.V., O.A. Hoekenga, and M.A. Pineros. 2004. "How do crop plants tolerate acid soils? Mechanisms of aluminum tolerance and phosphorus efficiency." Annu. Rev. Plant. Plant. Biol. 55:459-493.

Krizek, D.T. and C.D. Foy. 1988. "Mineral element concentrations of two barley cultivars in relation to water deficit and aluminum toxicity." J. Plant Nutr. 11:369-386.

Liu, H., J. R. Heckman, and J. A. Murphy. 1995. "Screening Kentucky bluegrass for aluminum tolerance." J. Plant Nutri. 18:1797-1814.

Liu, H. 2005. "Aluminum toxicity of seeded bermudagrass cultivars." HortScience. 40(1):221-223

Marschner, H. 1991. "Mechanisms of adaptations of plants to acid soils." Plant and Soil. 134:1-20.

SAS Institute Inc. 1999. SAS Version 8.0. SAS Inst., Cary, NC.

Wu, L., D.R. Huff, and J.M. Johnson. 1981. "Metal tolerance of bermudagrass cultivars." p.35-40. In R.W. Sheard Proc. 4th International Turfgrass Research Conference 19-23 July 1981. University of Guelph. Ont., Canada.

In these greenhouse studies, zoysiagrass and carpetgrass performed well when exposed to acid media with Al. Future research is needed to investigate the two species at the cultivar level to understand their mechanisms of Al tolerance.

Christian Baldwin is a Ph.D. candidate in turfgrass science. Dr. H. Liu is an associate professor and Dr. L.B. McCarty is a professor of horticulture specializing in turfgrass science and management. Dr. W.L. Bauerle is an assistant professor of horticulture, specializing in tree stress physiology and terrestrial ecosystem modeling. All are at Clemson (S.C.) University.

TURFGRASS TRENDS

SECTION STAFE **Managing Editor** Curt Harle

440-238-4556: 440-238-4116 (fax) curt@curtharler.com

Golfdom Staff Contact Thomas Skernivitz 440-891-2613; 440-891-2675 (fax) tskernivitz@questex.com

Online Editor Lynne Brakeman 440-826-2869: 440-891-2675 (fax) lbrakeman@questex.com

Chief Science Editor Dr. Karl Danneberge 614-292-8491; 614-292-3505 (fax) danneberger.1@osu.edu

Production Manager Carol Brisbin 218-279-8842; 218-279-8813 (fax) cbrisbin@questex.com

FIELD ADVISORS

Rob Anthony Southern Methodist University

J. Douglas Barberry Turf Producers International Agronomist

F. Dan Dinelli North Shore CC Merrill J. Frank

Columbia CO Michael Heacock Pacific Golf Management K. K.

Paul B. Latshaw Muirfield Village CC

Kevin Morris National Turfgrass Evaluation Program

EDITORIAL REVIEW BOARD

Dr. A.J. Powell University of Kentucky **Dr. Eliot C. Roberts**

Dr. Garald Horst University of Nebraska **Dr. Eric Nelson**

Cornell University Dr. Richard Hull University of Rhode Island

CONTACT US Editorial: 440-238-4556 Web site: www.turfgrasstrends.com Reprints: TurfgrassTrends@reprintbuyer.com

Graphic Designer Tracie Martinez 440-891-2725; 440-891-2675 (fax) tmartinez@questex.com

Publisher Patrick Roberts 440-891-2609; 440-891-2675 (fax) proberts@questex.com

Vice President & **General Manager**, Industrial & **Specialty Group** Tony D'Avino 440-891-2640; 440-891-2675 (fax) tdavino@questex.com

Corporate & Editorial Office 600 Superior Avenue, East Suite 1100 Cleveland OH 44114

Sean Remington Green Valley Ci

Ken Schwark, CGCS Northern Bay Golf Resort & Marina **Matt Shaffer** Merion G

Wayne Horman

David Irmen The Andersons

Van Cline The Toro Co. **Bill Byrnes** Floratin

Dr. Vic Gibeault University of California **Dr. Pat Vittum** University of Massachusetts Dr. Rick Brandenburg NC State University





The complete package to take care of all your bunker sand maintenance needs. Remove silts, clays, rocks, and remix your sands sieve sizes all in one easy process. Our maintenance free design leaves you with more time and money to focus on your courses needs.

The Sand Storm features a 12 volt DC powered vibratory motor. Our standard units are powered by Vibco. But, we also carry the equivalent motor made by Bosch for our customers residing in the EU certification regions.





So, what happens to the "undesirable" material being separated from your sand? All material larger than USGA spec, along with clay content, rolls off the back of the screen deck and into the removable waste bin. This allows you to dump your debris in the bed of your utility vehicle, or other receptacle, for disposal later. All that's left behind is the bunker spec sand you originally started with.

Phoenix, AZ info@zscreen.com Ph: 623-581-0307 www.zscreen.com





Our home is always welcome to unique visitors. In fact, in January alone, we had $12,040^*$ of them.

Of course... some were more unique than others.



CORPORATE CAPABILITIES

Aquatrols

Aquatrols

1273 Imperial Way Paulsboro, NJ 08066

Phone: 800-257-7797

Fax: 856-537-6018

Web site: www.aquatrols.com

Product focus:

Since 1954, Aquatrols has been the world leader in the development of cutting-edge surfactants and other specialty technologies to address the changing demands of the professional turf industry. We are proud to be an integral part of your management practices, and look forward to helping you meet the challenges ahead by continually striving to improve our product offering.

Technical support, sales, training and customer service:

Aquatrols combines state-of-the-art research and development techniques, a technically knowledgeable sales force, and a long history of industry expertise to better serve the needs of turf professionals worldwide. Our products undergo stringent quality control and assurance testing before shipment to provide the consistent levels of quality and safety that you demand.

Major product lines: Soil Surfactant Products

• Dispatch minimizes water lost to evaporation and runoff, allowing you to maintain turf color and quality with less water. Its unique mode of action also promotes more even penetration and distribution of applied nutrients and soil targeted turf chemicals for more effective use of these inputs. **Revolution** provides better moisture access and gas exchange in the rootzone, improving overall turf health resilience under all types of stresses. Revolution lets you gain control of rootzone conditions so that your turf can perform to its greatest potential. Available in liquid and granular formulations.

Primer Select can be applied monthly or biweekly to enhance turfgrass uni-formity by establishing matrix flow in the rootzone. Available in liquid and granular formulations.

• Aqueduct provides rapid recovery from existing water management problems. Available in liquid and granular formulations.

Specialty Products

Radiance Algaecide prevents algae in ponds, lakes and fountains using a self-dispersing delivery system that stays suspended in the water column.

LeafShield Anti-Transpirant protects turf and plants safely and affordably against winter desiccation. It dries to a thin waxy layer that remains flexible and will not discolor foliage.



QUATROLS Innovative Products for Effective Resource Management

ADVERTISEMENT

BASF Professional Turf & Ornamentals

BASF Professional

Turf & Ornamentals 26 Davis Drive Research Triangle Park, NC 27709

CORPORATE CAPABILITIES

Phone: 800-545-9525

Web site: www.turffacts.com



















The Chemical Company

Better turf comes from better science. Leveraging science from the world's largest chemical company, BASF Professional Turf & Ornamentals adds value to golf course superintendents with a solutions-based offering — both proprietary and post-patent products — which helps superintendents meet course and professional challenges. BASF gives superintendents and its distributor partners leading chemical effectiveness, long-term value, and technical support.

Insignia[®] fungicide

Insignia fungicide gives superintendents an effective, broad-spectrum strobilurin that suppresses dollar spot. Unlike older strobilurins, Insignia fungicide controls golf turf's top diseases, including brown, summer and take all patch diseases, pythium blight, bipolaris and gray leaf spot, snow mold and it suppresses dollar spot. Rainfast in one hour, Insignia is the next generation in strobilurin fungicides, providing superior performance and value in the fungicide rotation.

Emerald[®] fungicide

For golf course superintendents managing premium courses who need to control dollar spot — the No. 1 golf disease — Emerald fungicide is a unique class of chemistry specifically designed for dollar spot control. Emerald gives superintendents a crucial long-lasting alternative to existing dollar spot treatments that are losing their effectiveness. When applied preventively as part of a rotation or as a tank mix, its unique mode of action can actually control dollar spot unlike other fungicides that simply hold the disease in check.

Curalan[®] fungicide

Curalan fungicide offers economical, effective dollar spot control in cool- and warm-season turfgrasses, as well as controlling other diseases including brown patch, leaf spot, red thread, and melting out. Curalan adds further value by providing preventive protection with long residual control.

Iprodione Pro fungicide

Iprodione Pro fungicide prevents and cures key turfgrass diseases, including brown patch and dollar spot, in cool- and warm-season turf. It also controls large patch, fusarium blight, leaf spots, and several other diseases. Superintendents gain additional value with its long residual. Iprodione Pro is also an excellent product for key ornamental diseases in the landscape.

Propiconazole Pro fungicide

Propiconazole Pro fungicide controls nearly 20 key diseases in turfgrass, including dollar spot, brown patch, anthracnose, take-all patch, leaf spot, and pink and gray snow molds. Propiconazole Pro also controls numerous diseases on ornamentals, landscape and nursery plantings, including powdery mildews, rusts and leaf spots.

Basagran[®] herbicide

Basagran is a postemergent herbicide that offers exceptional value for control of annual and perennial nutsedges, kyllinga and numerous broadleaf weeds on cool- and warmseason turfgrasses.

Drive[®] herbicide

Drive postemergent broadleaf and grass herbicide gives superintendents quick-acting cleanup on broadleaf weeds including clover and grassy weeds like crabgrass, and can be used in overseeding with safety on many turfgrass species.

Pendulum AquaCap[®] herbicide

For superintendents who want long-lasting, preemergent control of crabgrass and 40 other grassy and broadleaf weeds, Pendulum AquaCap is an innovative formulation of BASF pendimethalin that greatly reduces odor, staining and eases equipment cleanup after application.

CORPORATE CAPABILITIES

ADVERTISEMENT

Bayer Environmental Science

Baver Environmental Science

2 T.W. Alexander Dr. Research Triangle Park, NC 27709

Phone: 919-549-2000

Fax: 919-549-3947

Web site: www.bayerprocentral.com

E-mail: bcs.gpcinfo @bayercropscience.com

Staff:

Josh Weeks, Vice President, Professional Business Dan Carrothers, Vice President, Chipco Professional Products Nick Hamon, Director, Development and Technical Services Mike Daly, Director of Marketing Scott Welge, Golf Business Lead Dave Ravel, Southern Regional Sales Manager Brian Tepper, Western Regional Sales Manager Jim Davis, Northern Regional Sales Manager

Product focus:

Bayer Environmental Science manufactures a full line of turf and ornamental products for use on golf courses, home lawns, sports fields, commercial landscapes, nurseries and greenhouses. Bayer Environmental Science continues a tradition of innovative research and top-of-the-line professional products, backed by highly trained sales and field development representatives. Some of our innovations in recent years include the Bayer Academy - an industry online training seminar - a service to turf manager and golf course associations, a Turf Scholarship program and a grant program to help superintendents attend the GCSAA show.

Manufacturing facilities:

In the United States, manufacturing facilities are located in Pasadena, TX; St. Louis, MO; and Kansas City, MO.

Technical support, sales, training and customer service:

Bayer Environmental Science supports its green industry products with a dedicated turf and ornamental research facility in Clayton, NC, where industry experts develop and investigate new chemistries. A network of technical development representatives works with university specialists throughout the country. Separate golf and lawn/landscape sales teams form an extensive nationwide network.

Major product lines:

From trusted standards like Merit® insecticide, Ronstar® herbicide, Proxy® plant growth regulator and CompassTM fungicide to newer products like Allectus® insecticide, RevolverTM herbicide and TopChoiceTM insecticide, Bayer Environmental Science offers products to fit all areas of turf maintenance.

A full line of fungicide products includes 26GT[®], SignatureTM, Bayleton[®], ProStar[®] and Banol®. Besides Merit and TopChoice, insecticide products include Dylox®, Chipco® Choice[™], DeltaGard[®], Tempo[®] and Sevin[®]. Herbicidesinclude Finale®, Sencor®, Prograss®, Illoxan®and Acclaim® Extra.

In addition, we continue to create new uses for existing products - additional tree injection uses for Merit, mosquito control for Tempo Ultra, improved turf quality with Signature, and longerlasting results with our fipronil products.



Backed by Bayer:

Bayer By Bayer means you get more than products, you get the confidence you'll succeed. The five pillars of Backed By Baver are:

- Dedicated Business Unit
- Guaranteed Results
- Solutions Approach
- Added Value
- Vision For The Future

MERIT[®] RONSTAR COMPASS allectus

FORBID Armada REVOLVER



CORPORATE CAPABILITIES

ADVERTISEMENT

Bell Laboratories, Inc.

Bell Laboratories, Inc. 3699 Kinsman Blvd. Madison, WI 53704

Phone: 800-323-6628

Fax: 608-241-9631

Web site: www.talpirid.com

E-mail: talpirid@belllabs.com



Company focus:

Bell Laboratories is a world leader in pest control products. From initial product inception and development to manufacturing and subsequent support in product use, Bell focuses on quality, functionality and innovation.

Manufacturing facilities:

Bell is a vertically integrated manufacturing company. All of Bell's products are designed, tested and manufactured in its modern facility in Madison, WI, which houses its manufacturing operation, offices, and biological and chemical testing laboratories.

Technical support, sales, training and customer service:

Bell's dedicated staff is

committed to providing quality pest control products and superior customer service at every stage — from product manufacturing to proper field use. This support includes internal and external technical expertise and technical representatives in the field that consult on difficult control problems.

Major product line:

Talpirid Mole Bait is the result of more than four years of groundbreaking research in Bell's biology and chemistry labs and in the field. It is the first and only mole bait designed, developed and scientifically proven to kill moles.



Talpirid's optimal size, shape and feel allow the bait to be consumed in the same manner as moles consume their natural food source, the earthworm. Special enhancers help ensure immediate attraction and excellent product acceptance. The active ingredient, bromethalin, capitalizes on the mole's physiology. One worm can kill a mole in 24 hours. Talpirid is easy to use and is proven to be extremely effective.

