

Improving Weaknesses of Preemergence Herbicides

By Scott McElroy

Preemergence herbicides are the basis of effective weed management programs. By utilizing preemergence herbicides, one can prevent a problem before it begins. Whereas postemergence herbicides that are safe to the desirable turfgrass must be selected and applied during the correct environmental conditions to limit potential injury, preemergence herbicides are generally safe across all turfgrass species and can decrease the need to use a potentially damaging postemergence herbicide. In light of the industry's upcoming loss of MSMA, the value of preemergence herbicides will become even more important in the future.

The top four preemergence herbicides used in turfgrass are prodiamine, pendimethalin, dithiopyr and oxadiazon. These herbicides vary in the way they are used, chemical characteristics and behavior in the plant.

Prodiamine and pendimethalin are mitotic inhibiting herbicides within the dinitroaniline family. They are orange in color and were some of the first preemergence herbicides utilized in turf. They remain the standards of turfgrass management. Dithiopyr is similar to prodiamine and pendimethalin in that it inhibits mitosis; however, inhibition is at a slightly different site and dithiopyr can also inhibit shoot mitosis. Dithiopyr also has the added benefit that it can control crabgrass (*Digitaria* spp.) postemergence before the plants tiller.

Oxadiazon has a completely different mode of action. Unlike these other preemergence herbicides, oxadiazon is a protox-inhibiting herbicide generating rapid cell degradation as a plant attempts to conduct photosynthesis. Oxadiazon provides more broad-spectrum control, but it tends to be more expensive.

While these herbicides have many benefits, they also have some negative features.

Prodiamine and pendimethalin primarily control annual grasses and small seeded broadleaf weeds. They do not provide preemergence control of a wide spectrum of broadleaf weeds and they offer limited control of sedges. Dithiopyr has a similar weakness in that it primarily controls annual grasses preemerge and provides little to no sedge control. Oxadiazon provides better preemergence control of some broadleaf weeds and sedges; however, it cannot be used in residential turf and it must be applied as a granule to dry turf to avoid potential injury.

To improve upon traditional herbicide performance, the following is needed:

- Improved control of broadleaf weeds.
- Improved sedge control.

A recently introduced herbicide active ingredient, sulfentrazone, addresses these needs and offers some potential benefits over traditional products. Sulfentrazone is a protox-inhibiting herbicide similar to oxidi-

Continued on page 62

Echelon applied for fall *Poa annua* control also provides postemergent *Kyllinga* control. Note the killed *Kyllinga* spp. within the Echelon treated area.



There are three benefits to Echelon that most traditional preemergence herbicides do not provide.

Ad Index

Advertiser	Page
Agraquest	4
Agrium Adv. Tech.	Cvr Tip, CV2-1
The Andersons	25, 39
B A S F	5, 13
Buffalo Turbine	24
Champion Turf Farms	31
Dow AgroSciences	23
Duro Tire Corp	35
FMC	9
Gro Power	51
Hustler Turf	45
John Deere Golf	54
Kochek	47
Lebanon Turf	Insert, CV4
PBI Gordon	7, CV3
Rainbird	15
Redexim Charterhouse	11
Reliable Golf	40
Richway Industries	51
Solu-Cal USA	50
Spectrum Technologies	53
Toro	21
Turfco Mfg	41
Turfline	Insert
White Metal Golf	52
Wireless Solutions	29
Woodbay Turf	34

This index is provided as an additional service. The publisher does not assume any liability for errors or omissions.

Continued from page 61

azon as a preemergent herbicide, but safe to apply as a postemergence herbicide as well. As a postemergence herbicide, sulfentrazone is fast-acting with control occurring within three days after treatment. It is primarily used to control sedge species postemergence, and is sold as Dismiss or Dismiss South (sulfentrazone plus imazethapyr) for enhanced purple nutsedge control.

Sulfentrazone has also been sold as mixtures with broadleaf herbicides (Q4 and Surge) to provide faster activity and as Solitaire (sulfentrazone plus quinclorac) for sedge, crabgrass and broadleaf weed control. Sulfentrazone is also unique in that it can potentially control goosegrass; ongoing research continues to demonstrate this added benefit.

Echelon is a prepackaged mixture of prodiamine and sulfentrazone developed to improve upon the weaknesses of traditional preemergence herbicides. Sulfentrazone has been used in row-crop agriculture for many years as an additive to dinitroniline herbicides to improve preemergence sedge and broadleaf control. A similar type of control is expected and has been seen from Echelon.

As a professor at the University of Tennessee and now at Auburn University, I have conducted approximately 20 studies evaluating the usefulness of sulfentrazone in mixture with prodiamine (as is used in Echelon). In order to better understand its usefulness, it is easiest to simply compare Echelon to standard products.

First, from my data, Echelon offers similar control of crabgrass and *Poa annua* as traditionally used herbicides, such as Barricade. Both provide excellent control of these weeds when used at the proper rate; so if these weeds are your main target then either herbicide will work.

Having said that, there are three benefits to Echelon that most traditional preemergence herbicides do not provide. First, it provides added goosegrass control (Figure 1).

Goosegrass (*Eleusine indica*) is a later germinating annual grass species common in turfgrass. It typically germinates four to eight weeks after crabgrass germination. Using split application programs can

improve goosegrass control, especially if Echelon is used as the second application.

A second benefit from this program is the added preemergence broadleaf control. Typically, this mixture of prodiamine and sulfentrazone is better for winter annual broadleaf weed control, such as henbit and common chickweed, than summer annuals, such as annual lespedeza. Similar to goosegrass control, split application programs that include an initial application of Echelon or a traditional preemergence herbicide followed by Echelon can improve control of annual and non-tuberous sedges, such as annual flat sedge (*Cyperus compressus*) and globe sedge (*Cyperus globulosus*). By using Echelon as the second preemerge application, sedges can germinate, thus allowing the product to provide pre- and postemergence activity. Echelon typically can provide burn back of these weeds, but control throughout the summer may dissipate.

A third benefit is pre- and postemergence sedge control (Figure 2). Similar to goosegrass control, split application programs that include an initial application of Echelon or a traditional preemergence herbicide followed by Echelon can improve control of annual and non-tuberous sedges. Using Echelon for the second application, sedges can germinate, thus allowing both pre- and postemergence activity. Echelon typically can provide burn back of these weeds, but control throughout the summer can dissipate.

The combination of prodiamine and sulfentrazone in Echelon brings some advantages to the turfgrass market for pre-emergence weed control. These benefits are best seen when Echelon is used in a split-application preemergence program as listed in Table 1.

As demonstrated in this article, Echelon provides similar benefits as traditional herbicides for general crabgrass and *Poa annua* control. However, it can improve weed control programs when used as a split application program for sedge and goosegrass control.

Scott McElroy, Ph.D., is an associate professor of turfgrass/weed science in the Department of Agronomy and Soils at Auburn University. He can be reached at jsm0010@auburn.edu.

Golfdom

**Giveaways @
Booth 2181**

golf industry show

Feb. 9-10, 2011

Orlando, FL

**Just Drop Off Your
Business Card for a Chance to Win!**

GRAND PRIZE ALL EXPENSES PAID TRIP TO PINEHURST AND THE GOLFDOM SUMMIT!

November 13-15, 2011

Pinehurst Resort • Village of Pinehurst, NC

www.golfdomsummit.com

The **FIRST** invitation-only, highly focused forum bringing together superintendents, suppliers and dealers of products used for the operation and design of golf courses.




Scan QR Code with your mobile to qualify for the Golfdom Summit today!

www.golfdomsummit.com

2nd PLACE PRIZE

WIN a one-of-a-kind autographed copy of Golfdom magazine's July 1962 issue signed by Arnold Palmer and Jack Nicklaus.



**Stop By to Meet Our New Editor-In-Chief
Seth Jones**

For more details, as well as a schedule of the booth, visit
<http://www.golfdom.blogspot.com/>

Teethrax outbreak!

The time has come for superintendents to get in touch with their subversive side. Only the future of the game is at stake. Let me explain.

Golfers are a sadistic lot and water cooler discussions during the last year have turned to making the game more fun. For any number of reasons, most efforts to return fun to the sport die on the operating room table.

Maybe it's the effete nature of our sport compared to football or even tennis. Perhaps the problem stems from our origins from stiff-upper-lip Scotland, where they pride themselves in only playing real golf in wind and rain. Or maybe golfers just like to be punished. Either way, in the face of modern distance advances the game is constantly trying to man-up and run from anything which might make matters a little more enjoyable. We grow up the rough, add in new back tees and plant more trees. But nowhere is the self-torture more evident than in the tees golfers choose to play.

We all know that an unusually excessive number of golfers insist on playing a course from as far back as possible even as modern designers have offered them more options than ever. And plenty more golfers would never think of going to the red tees, especially older men who think any association with the dreaded red will have them strapping on ballerina dresses and rushing home to watch Oprah.

Female golfers are just as neurotic about their tees, often howling in agony at the thought of architects retrofitting older designs to give them the experience of playing shots into greens

THE SCIENTISTS SAYS THIS INVISIBLE FUNGUS ONLY OVERTAKES TEEING GROUNDS ON GOLF COURSES OVER 6,500 YARDS.

BY GEOFF SHACKELFORD



with clubs that reflect the architectural intent. This is why it's time to forget about the tees altogether.

In the January *Golf Digest* "Fun" issue, writer Ron Kapriske details how he set his manhood aside, first playing a set of 5,301-yard tees and posting an 81. The next day he created his own teeing grounds, moving twenty or so yards in front of the forward tees to get a course at about 4,941 yards. He fired 78 and wrote, "this was the first time I felt a round of golf was truly fun from start to finish."

Kapriske took his fun antics to a public course where, as a single joining a group, he introduced himself on the first tee, then revealed to his new buds that he would be playing the forward tees. After the first three golfers teed off in front of a mini gallery, Kapriske swallowed his pride and walked forward. He didn't look back but you can imagine the snickering that took place when a young, healthy 12 handicapper went so far forward.

So this is where you, the superintendent come in. Because — bare with me here — it's been revealed on various password-protected online discussion boards devoted to turf care that there is a very strange turf disease making its way around

the golf world. It's called Teethrax. The scientists says this invisible fungus only overtakes teeing grounds on golf courses over 6,500 yards. Golfers on these afflicted layouts must not get within 30 yards of any teeing ground until the appropriate chemical compound can be found to combat a perplexing virus known to induce the common cold, as well as the yips, on unsuspecting golfers.

So while the labcoats are hard at work solving this frightening mystery, your job will be to protect your golfers by placing all tees up at the start of fairways for two weeks so that golfers can continue to enjoy their course, even though that's obviously not as fun as slugging it out from the tips. But hey, safety first.

Then just sit back and watch what happens.

Oh sure, there'll be a few traffic jams and maybe even a few safety concerns to monitor with new landing zones. Mostly you'll hear a few laughs, some cheers and most of all, seeing the smiles coming off the 18th green will tell you all you need to know.

Follow Shack, Golfdom's contributing editor, at www.geoffshackelford.com or via [Twitter.com/GeoffShackelford](https://twitter.com/GeoffShackelford).

WHEN YOU NEED **MUSCLE** IN YOUR TANK

**NEW
INSECTICIDE**



CONTROL OF TOUGH TURFGRASS INSECTS!

- CONTROLS ROOT- AND LEAF-FEEDING INSECTS
- RESIDUAL SYSTEMIC CONTROL
- KNOCKDOWN ACTIVITY
- USE IN TROUBLESOME TURF AREAS

EFFECTIVE ON:

- ▶ CHINCH BUG
- ▶ MOLE CRICKET
- ▶ EUROPEAN CRANEFLY
- ▶ ANNUAL BLUEGRASS WEEVIL



COMING SOON
FOR TREES & SHRUBS
ZYLAM™ LIQUID
SYSTEMIC INSECTICIDE

**G pbi / Gordon
corporation**
An Employee-Owned Company
800.821.7925 • pbigordon.com/zylam

ALWAYS READ AND FOLLOW LABEL DIRECTIONS
Zylam is a trademark of PBI-Gordon Corporation.

01169