



Questions From the Floor

By Dr. R. Chris Williamson, Department of Entomology, University of Wisconsin-Madison

1) I've treated my golf course for Japanese beetle grubs for five consecutive seasons; last year there was very little evidence of adults, grubs, or damage. Should I make an insecticide application this year? – Dane County, Wisconsin

Response: Historically speaking, Japanese beetles tend to establish in respective geographic areas for about 5-10 or so years. Various factors influence the length of time which they sustain damaging populations. Such factors include host plant availability (food sources), environmental conditions including soil moisture, natural enemies including insect predators and parasitoids as well as bacterial, fungal, and viral pathogens, and insecticide management practices. Only through a sound Integrated Pest Management (IPM) plan that utilizes routine sampling and monitoring can one effectively make an informed decision as to the need to make an insecticide application. If the data generated from an annual Japanese beetle sampling and monitoring program suggests that populations have been continuing to dramatically decrease, it may be advantageous to forgo making a preventative Japanese beetle insecticide application. Keep in mind, this approach is not fool proof. But, with the development of effective curative (corrective) white grub control products such as chlothianidin (Arena), prophylactic applications of insecticides to putting green surrounds, tee boxes and surrounds, fairways, and rough areas may not be necessary. Ultimately, this IPM approach will reduce the amount of pesticides applied, reduce the monetary cost of controlling Japanese beetles, and minimize potential environmental impact.

2) When do you think or predict the emerald ash borer (EAB) will become a problem in Wisconsin? – Waukesha County, Wisconsin

Response: If I could earnestly predict when and where EAB will be found in Wisconsin, I would likely be considered for some type of distinguished award. Realistically, EAB poses an eminent and serious threat to Wisconsin, especially since Wisconsin borders Michigan (including the Upper Peninsula) and is in relatively close proximity to Indiana where EAB is currently established. EAB was most recently discovered in the eastern Upper Peninsula of Michigan at Brimley State Park. Folks in Wisconsin and Minnesota are particularly concerned about EAB since both states have significantly more ash resources than Michigan or

Indiana, thus stand to lose much more. EAB is thought to spread by four primary means: 1) movement via firewood; 2) transportation via ash (*Fraxinus* spp.) nursery stock; 3) pallets and solid-wood packing material; and 4) adult flight (< 1/2 mile per year). Of these four, the movement of infested firewood from EAB infested areas appears to be the most likely means of introduction into un-infested areas. Thus, it is critical to educate as many people as possible about the threat that EAB poses as well as how its introduction can be minimized. So, to predict where EAB will be discovered first in Wisconsin is anyone's guess; however, the hypothesized geographic areas where EAB is most likely to be discovered are the gateway areas in Wisconsin including but not limited to SE and NE Wisconsin. Other high-risk areas for EAB include commercial properties where ash nursery stock may have been obtained from



**Everything To Make Your
Course Beautiful**

**Turf Seed • Fertilizers • Repair Parts
Golf Course Accessories • Equipment**

LESCO®

800 - 321 - 5325

LESCO is a registered trademark and Grow With Us is a trademark of LESCO Technologies, LLC.