


**You just
can't buy
better gang
mowers.**



**It's been proven time and again, day
after day, acre after acre. You just
can't buy better cutting gang mowers.
For a very convincing demonstration,
call your Jacobsen Distributor.**

**JACOBSEN
TEXTRON**

Jacobsen Division of Textron Inc.

HORST DISTRIBUTING, INC.

444 N. Madison Street, Chilton, Wisconsin 53014 Phone 414-849-2341

Pythium and Sand Topdressing

Clinton F. Hodges

Department of Horticulture Iowa State University Ames, Iowa

(Increasing evidence for Pythium induced root dysfunction of creeping bentgrass grown in high sand content mixes.)

The presentation made by me at the 1981 M.G.C.S.A. annual conference implicated Pythium in a root dysfunction of creeping bentgrass grown on reconstructed high sand content greens. The disease in question killed grass very rapidly and in a manner typical of Pythium "Cottony Blight". Examination of diseased plants, however, failed to yield Pythium or any other pathogen from above-ground portions of the plant. Pythium was found throughout the root system of diseased plants. It is believed that when a green on an old golf course is renovated to sand, Pythium may infest the sand from the collar and apron soil of the old green. The sand probably offers little microbiological competition for the Pythium and increased levels of irrigation and fertilization of sand greens may further promote the Pythium. There also is some evidence that more root mechanical damage may occur in sand than in soil; this could contribute to Pythium infection. This problem has not been observed in the roots of plants grown in traditional soil mixes.

To date, we have isolated four species of Pythium from the roots of creeping bentgrass associated with rapid death of infected plants. We are currently in the process of identifying the isolates to the species level and testing them for pathogenicity. It has been possible to infect roots in greenhouse studies and in some cases reduce the rate of growth of infected plants. Our present observations suggest that the Pythiums in question may infect the roots and co-exist with the plants with minimal damage under mild growing conditions. It may necessitate some form of environmental or cultural stress before death occurs. There are two

outstanding problems for the superintendent relative to this disease. These are diagnosis and control. The rapid death of infected plants appears to be due to foliar pathogen. However, examination of dying leaves usually fails to yield any pathogens and often times even saprophytic organisms are not present. When the root systems are examined they usually appear normal in size and color. Because of this it is assumed that root pathogens are not present. These normal appearing roots can, however, be severely infected by Pythium. The Pythiums associated with this problem do not cause a rot and the degree of discoloration may not be detectable with the naked eye. We have found that when we properly incubate what appears to be a healthy root from these diseased plants, Pythium will grow from the root within six hours. We are suspicious that these Pythiums damage plants by interfering with water relations, not by rotting.

Control of this problem in the field remains elusive. Intense aeration followed by application of Pythium specified fungicides into the aerifier holes may slow the disease. There is some indication that wetting agents used in conjunction with the fungicides may be beneficial. It also appears that following renovation of a green the first time the disease strikes it most severely; each subsequent year it becomes less severe.

Our primary research objectives for the next 2-3 years will be as follows: 1) Continue to collect and identify Pythium species associated with roots. 2) Determine pathogenicity of the Pythium species and the conditions necessary for injury or death to occur. 3) Determine the nature of the pathogenicity; i.e. in that the roots are not rotted, how are the plants ultimately killed. 4) Examine approaches to control.

MILORGANITE

A WISCONSIN ORIGINAL

Slowly released natural organic source of
nitrogen—phosphorus—iron

Granular—Easily Spread—Non-burning

Just right for Dormant Fertilization

Trouble-free Spreader Calibration

Ready Availability—Low Cost

Bulk or Bag Delivery

Ask us about sprayable fine grade for greens and tees.

Wisconsin Turf Equipment Corp.

P.O. Box 708
1917 West Court Street
Janesville, WI 53545
(608) 752-8766

Milwaukee Metropolitan

Sewerage District
735 North Water Street
Milwaukee, WI 53202
(414) 278-2034

Horst Distributing, Inc.

444 North Madison Street
Chilton, WI 53014
(414) 849-2341

1983
**WISCONSIN GOLF COURSE
 SUPERINTENDENTS
 ASSOCIATION**

MEETINGS

Nov. 7th
 Monday

Hartford C.C.
 Hartford, Wi.

Fall Business Meeting
 & Election

LY-MAR
 LTD.

DISTRIBUTOR

**top name golf course
 chemicals & supplies**

8332 NORTH SANTA MONICA BLVD.
 MILWAUKEE, WI 53217 (414) 352-9464

The best in turf supplies
LESCO 100% Sulfur-Coated Fertilizers
 golf course accessories, chemicals including

Lescosan

A highly effective pre-emergence
 crabgrass and poa annua control

* Lescosan (Betasan - Registered TM Stauffer Chemical Co.)

(800-321-5325)
 NATIONWIDE

Available from:

(800-362-7413)
 IN OHIO


Lakeshore
Equipment & Supply Co.
 "Home of LESCO Products"
 300 South Abbe, Elyria, OH 44035

"ENGINEERED by PROFESSIONALS for PROFESSIONALS"

TORO

GREENMASTER 3

NOW AVAILABLE WITH
HAND CONTROLLED
VARIABLE GROUND SPEED KIT



THATCHING REELS

SPIKING REELS

Top ranked in quality of cut because no other greensmower in the world has cutting heads isolated from the traction unit and grass baskets. Engineered for turf care professionals by the professionals in turf care products.

Reinders

TURF EQUIPMENT

13400 W. WATERTOWN PLANK RD., • ELM GROVE, WIS. 53122
PHONES: SALES (414) 786-3163 — OFFICE (414) 786-3300