

TABLE OF CONTENTS

THE PRESIDENT'S MESSAGE	
A Fresh Start	3
GAZING IN THE GRASS	
Liquid Versus Granular Fertilizer Applications For Shaded Tees.	5
WISCONSIN SOILS REPORT	
Nitrogen and Phosphorus Losses from Turf	10
MISCELLANY	
Gypsy Moth Treatment Areas Proposed for 22 Wisconsin Counties	17
WISCONSIN ENTOMOLOGY REPORT	
Know Your Pest!	19
WGCSA	
2002 WGCSA Spring Educational and Business Meeting	20
GOLF IN THE FLATLANDS	
The Five Year Plan	23
PERSONALITY PROFILE	
There's Just Something About Wayne.	29
NOTES FROM THE NOER FACILITY	
You Are At-Risk	37
JOTTINGS FROM THE GOLF COURSE JOURNAL	
J.I. Case: Wisconsin Industrialist.	40
GREEN SECTION	
The Shape of Things to Come	45
THE EDITOR'S NOTEBOOK	
The Wisconsin Seasons Merge	49
WISCONSIN PATHOLOGY REPORT	
Have Less Snow Mold This Year? We're Trying to Find Out Why	53
POA TRIVIA	
.	56

ABOUT THE COVER

Artist Jennifer L. Samerdyke has created a beautiful cover for this early summer issue of The Grass Roots. Entitled "The Moon Over the Sixth at Serenity Bay Country Club," her original creation will evoke memories for Wisconsin golf course superintendents who all have returned to the golf course at night one time or another, usually to deal with an irrigation problem. And who among us

hasn't been struck by the beauty of that landscape usually seen in daylight? The subtleties and shadows are enhanced on those nights when the moon is full.

"To work — to work! It is such infinite delight to know that we still have the best things to do."

*Katherine Mansfield
(1883-1923)*

THE GRASS ROOTS

THE GRASS ROOTS is a bi-monthly publication of the Wisconsin Golf Course Superintendents Association, printed in Waukesha, Wisconsin by Kramer Printing. No part of THE GRASS ROOTS may be reprinted without expressed written permission by the editor.

EDITOR AND PUBLISHER

Monroe S. Miller
Blackhawk Country Club
P.O. Box 5129
Madison, WI 53705
groots@globaldialog.com

EDITORIAL STAFF AND BUSINESS AFFAIRS

Dave Brandenburg
P.O. Box 314
Theresa, WI 53091



BOARD OF DIRECTORS

Back row: (L-R) Kendall Marquardt, Dustin Riley, Jack Tripp, Kris Pinkerton, Brian Ferrie. Front row: (L-R) Mike Lyons, Marc Davison, David Brandenburg, Randy Witt.

2001-2002 WGCSA OFFICERS AND DIRECTORS

PRESIDENT

David Brandenburg
Rolling Meadows GC
P.O. Box 314
Theresa, WI 53091
E-mail: fdclnty@dotnet.com

DIRECTORS

Brian Ferrie
Horseshoe Bay CC
5335 Horseshoe Bay Rd.
Egg Harbor, WI 54209
E-mail: hbaysoup@itol.com

VICE PRESIDENT

Marc Davison
Green Bay CC
2400 Klondike Rd.
Green Bay, WI 54311
E-mail: mdavison@usxchange.net

Kendall Marquardt
Lake Wisconsin CC
6796 Breunig Rd.
Mazomanie, WI 53560
E-mail: kmarqt@speagle.com

TREASURER

Mike Lyons
Old Hickory CC
316 Grove Street
Beaver Dam, WI 53916
E-mail: mlyons@powerweb.net

Jack Tripp
La Crosse CC
N4627 Timbercrest Dr.
Onalaska, WI 54650
E-mail: jttripp@centrytel.net

SECRETARY

Randy Witt
Oneida, G&CC
618 Night Ct.
Green Bay, WI 54313
E-mail: RHWitt@aol.com

Dustin Riley
Oconomowoc GC
247 N. Main Street # 11
Dousman, WI 53118
E-mail: ogriley@voyager.net

PAST PRESIDENT

Kris Pinkerton
Oshkosh CC
11 West Ripple Ave.
Oshkosh, WI 54902
E-mail: occkjp@vbe.com



A Fresh Start

By David Brandenburg, Golf Course Superintendent, Rolling Meadows Golf Course



I feel superintendents in Wisconsin are lucky to have four seasons of weather. The winter season gives us an opportunity to prepare for the next year and not worry so much about the day-to-day grind of course maintenance and player concerns. Best of all the spring season gives us a fresh start with staff, equipment and ideas.

The off-season allows staff to go over equipment so it is painted and polished. Reels and cutting blades are sharp and set perfectly for that first cut. Many of us see new units added to our equipment fleet to help us maintain the property up to expectations. I know our entire staff is excited with the fresh start of a new or redone piece of equipment.

New employees are another highlight of spring. I think back to my first day at Sunset Hills in Beaver Dam and how excited and nervous I was to join the course staff. If you look and listen, you can see that feeling of excitement in most new hires. We have quite a few students on our staff and every year a few move on to their intended careers. For returning seasonal or full time employees, a fresh start means forgetting those little mistakes of last year and a commitment to be better than ever. It also means a renewal of friendships within the staff from the year before.

As managers, we need a fresh start along with our staff and equipment. In winter, I write down goals for improving the operation as they come to mind. It could be new improvements or something we may not be doing to the best of our ability. Spring is a great time to implement new ideas because old habits are easier to break after taking a few months off. As the season approaches, I take the goals and list what we need to do to accomplish them.

For me personally I need to communicate better by expressing my expectations. I explain things to the point I understand them, but I need to go to the next level and explain them until the entire staff understands the idea. I also have a goal of keeping my office organized to reduce time looking for things or handling papers twice.

For our staff, we need to improve our bunker maintenance to protect the borders we work so hard at edging. We also need to develop a system for our rough and trim mowers to cover the entire course effectively. Most of the rough is mowed as it should be, by the proper machine. However, three people mow some of the areas, and then we have those little areas that everyone assumes someone else took care of that are not mowed at all. By coordinating our efforts, we can easily cover each area once without double covering or skipping. We are also starting a first cut of rough around greens and fairways this season to provide a better product.

What are your goals for this year? Take the time this spring while you have a fresh start to reset your goals and priorities. If you do not do it now, spring will slip by and it will be too late. Good luck and God bless in your efforts. ♣



NX-PRO® now offers a complete line of microgreens grade fertilizers. Our three new SGN 100 fertilizers deliver superior particle uniformity and the most efficient methylene urea complex nitrogen source you can find - Meth-Ex® 40. So, now you can have the same excellent NX-PRO results on your greens as you've experienced on tees and fairways. NX-PRO SGN 100 Fertilizers - a terrific microgreens product to strengthen your short game.

Frank Baden
Territory Manager
Bettendorf, IA
(563) 332-9288



EXPERIENCE THE *Reinders* DIFFERENCE

- **Longevity**—We've been serving customer needs since 1866.
- **Stability**—We provide you with "Solutions and Supplies," Always Have, Always Will.
- **Solutions**—We provide recommendations and suggestions to help solve any challenge.
- **Supplies**—We have thousands of quality products in inventory.
- **Experience**
More than 200 years of combined turf industry experience on staff
More than 175 years of combined irrigation industry experience on staff
More than 50 years of combined landscape industry experience on staff
Three Degreed Turf Specialists on staff
One Certified Sports Field Audit Consultant on staff
Seven former Golf Course Superintendents on staff (including 4 CGCSs)
Three former landscape contractors on staff
- **Credibility**—We have worked with hundreds of golf courses, schools, municipalities, landscapers, retailers, etc. for more than 135 years.
- **Delivery**—Products or equipment delivered when you want it, where you want it.
- **Locations**—We have five store locations throughout Wisconsin.
- **Convenience**—High quality products at competitive prices can be picked up or delivered to you.
- **Follow-up**—We provide after sale service to ensure your needs have been met.
- **Education**—We offer industry education programs every year at the Reinders Turf Conference, Reinders Service School and regional seminars.

Reinders

Solutions & Supplies for the Green Industry



Proud Supplier of Turf Equipment and Irrigation Products to the Milwaukee Brewers and Miller Park



ELM GROVE
(262) 786-3301
13400 Watertown Plank Road
(800) 785-3301

MADISON
(608) 244-0200
4217 Nakoosa Trail

APPLETON
(920) 788-0200
900 Randolph Drive

STEVENS POINT
(715) 342-3600
3510 Post Road (Hwy. 54 & 51)
Plover

KENOSHA
(262) 857-3306
20830A 75th Street (Hwy. 50)
Bristol

www.reinders.com

Liquid Versus Granular Fertilizer Applications for Shaded Tees

By Dr. John Stier, Department of Horticulture, University of Wisconsin-Madison

Growing grass in the shade is never easy because most turfgrasses are adapted to full sun conditions. The buffered temperature fluctuations, high humidity and long periods of leaf wetness increase the potential for disease. Lack of suitable light quality and quantity greatly affect turf growth and development, decreasing their traffic tolerance and ability to recover from damage. The cut-off point for many cool-season turfgrasses subjected to traffic and with high quality expectations is approximately 30% full sunlight. When the turf receives

less than 30% full sunlight management becomes difficult if not impossible.

Traditional recommendations have included reducing the amount of nitrogen (N) fertilizer to about half the normal rate. Actual data on the effects of different types of N carriers or formulations is little to nonexistent. Many turf managers, including those at Miller Park in Milwaukee, use foliar-applied N. While foliar N applications can be economical and produce a quick turf response there is little peer-reviewed information available which compared

foliar to granular N sources from an agronomic perspective. Spangenberg et al. (1986) conducted one of the few scientific studies to compare liquid versus granular N applications. On Kentucky bluegrass, growing in full sun, liquid urea applications resulted in better turf color than granular applications, at least until the latter part of summer. Jiang and Hull (1999) showed turf uses energy to absorb and process N in the roots, much of which is ultimately transported to leaves. Bushoven and Hull (2001) reported foliar N applications



syngenta

Consider Primo MAXX® the turf's personal trainer.

Get your turf in top shape. Primo MAXX PGR makes turf thicker and stronger. Preparing it for summer's toughest challenges, like heat, drought, and disease. Not to mention everyday wear and tear. So it can be in prime condition all season long.



Contact Jim Shone at 1-708-217-8509 to learn more
about Syngenta products.

Important: Always read and follow label instructions before buying or using this product.
©2002 Syngenta. Syngenta Professional Products, Greensboro, NC 27419. MAXX®, Primo®, and the Syngenta logo
are trademarks of a Syngenta Group Company.

www.syngentaprofessionalproducts.com

should increase N use efficiency, allowing more energy to be allocated for root growth. Since energy levels on shaded turf are low compared to unshaded turf, it makes sense that foliar N applications should increase turf quality.

A research project was conducted during 2000 and 2001 to provide information for developing a management program for shaded golf course tees. Our goals were to identify the best grass species, application intervals for Primo (trinexapac-ethyl), and foliar versus granular urea applications. Research on Primo effects on turfgrass under reduced light conditions has been previously reported (Stier and Rogers, 2001; Goss et al. 2002).

MATERIALS AND METHODS

Plots were established in August 1999 at the O.J. Noer Turfgrass Research and Educational Facility by placing washed sod on a silt loam soil. The grass species were 'Penncross' creeping bentgrass (*Agrostis stolonifera*), 'Supranova' supina bluegrass (*Poa supina*) and a blend of five elite Kentucky bluegrass (*Poa pratensis*) cultivars. Grasses were arranged in a randomized complete block design with four replications of each species. Plots were split to evaluate the effects of Primo applied at 0.125 oz/1000 ft² monthly, bimonthly, or not at all. Plots were split perpendicular across the Primo applications to evaluate the effects of foliar versus granular urea applications. Urea was applied at 0.5 lb/1000 ft² at 14 day intervals during the growing season. Granular treatments were irrigated in immediately following application, followed by foliar applications to the rest of the plots. Foliar urea was applied by dissolving feed grade urea in water at the rate of 1.1 lb urea per gallon of water and spraying the solution onto the turf using a carrier volume of 2 gallons of water per 1000 ft².

Plots were maintained at 0.5 inch height and irrigated once weekly at 50% estimated evapotranspiration of fully sunlit conditions. Turf was topdressed monthly with 0.062 inch depth of 80:20 sand:peat mixture. Shading was applied by placing an 80% shade cloth approximately 7 feet above the turf each spring to coincide with approximately 80% leaf development on nearby trees. Shade cloth was removed in the autumn to coincide with loss of approximately 80% of the leaves on nearby trees. A datalogger and environmental sensors were placed in the center of the shaded area to monitor air temperature and humidity near the turf canopy, soil temperature at 2 inch depth, and photosynthetically active radiation (PAR; 400-700 nm). A second weather station at the site was situated in full sun to provide comparative data.

Turf quality (density and uniformity) and color were rated biweekly on a one to nine scale, with one equivalent to dead turf and nine equivalent to ideal turf; a rating of six was considered acceptable. Turf density was evaluated each spring, summer, and fall using an optical point quadrat method. Divots were made using a divoting tool each season, backfilled with a sand:peat topdressing, and evaluated for percent regrowth. Nitrogen and chlorophyll concentrations were measured twice in leaf clippings during 2001. Chlorophyll fluorescence measurements were collected several times in both years to provide an indication of photosynthetic efficiency. Core samples were collected each autumn to determine root mass of each treatment. The trial was conducted over two full growing seasons, 2000 and 2001, to help ensure accurate data interpretation.

RESULTS

Environmental data

Shading averaged 80% of full sun

plus or minus 4% during 2000-2001. The lowest PAR values were recorded in early October before the shade cloth was removed (approximately 2 mol m⁻² day⁻¹) and the highest were recorded during the summer solstice in June (approximately 7 to 9 mol m⁻² day⁻¹).

Effect of Nitrogen type

For purposes of brevity, only the data directly concerned with fertility will be described. There were no interactions between fertility and PGR. The type of N fertility did not affect divot recovery, root mass, or photosynthetic efficiency. There were, however, surprising and consistent interactions between type of urea application and grass species.

Turf quality of creeping bentgrass was significantly better when foliar urea was used compared to granular applications (Table 1). Conversely, Kentucky bluegrass responded better to granular applications of urea compared to foliar applications. While creeping bentgrass turf quality was the same or better than Kentucky bluegrass quality, supina bluegrass turf quality was as good or better than creeping bentgrass. When the data were averaged for each year, supina bluegrass appeared to perform similarly regardless of N type. In reality, supina bluegrass quality was enhanced during late summer and fall when foliar applications of urea were used (Fig. 1). Fertilizer effects on turf density were similar to the effects on turf quality.

Color of supina bluegrass was greatly enhanced by Primo to the point that when data were averaged across all Primo treatments, supina bluegrass color was not often significantly different than creeping bentgrass or Kentucky bluegrass (data not shown). Nitrogen content of leaves, though, indicated supina bluegrass had significantly less N than creeping bentgrass or Kentucky

Table 1. Interaction between urea N and species on average turf quality under 80% shade, Verona, WI.

Species	2000		2001	
	-----Nitrogen type-----			
	Granular	Foliar	Granular	Foliar
Creeping bentgrass	6.3	6.9	5.7	6.5
Supina bluegrass	7.2	6.8	6.7	6.7
Kentucky bluegrass	6.5	5.6	6.0	5.4
LSD (0.05)	0.3		0.4	

Table 2. Interaction between urea N type and turf species on total leaf nitrogen content, Verona, WI, 2001. Nitrogen was applied biweekly during season, including 20 July and 4 August at 0.5 lb N per thousand square feet.

Species	Nitrogen concentration (% total N)			
	3 August		15 August	
	Granular	Foliar	Granular	Foliar
Creeping bentgrass	5.4	5.3	5.3	5.2
Supina bluegrass	4.9	4.4	4.8	4.4
Kentucky bluegrass	5.2	4.8	5.2	4.7
LSD (0.05)	0.2		0.2	




Figure 1. Turf quality of supina bluegrass depends on type of urea and time of year, Verona, WI, 2000.

bluegrass. Granular fertilization significantly increased the N content in leaves of both supina bluegrass and Kentucky bluegrass while N content in clippings of creeping bentgrass was not affected by N type (Table 2). Nitrogen content did not seem to correlate with chlorophyll concentration, however, as foliar N resulted in significantly ($p = 0.05$) greater chlorophyll amounts for all species compared to granular N treatments (approximately 26 versus 23 to 24 micrograms of chlorophyll per gram of fresh leaf tissue).

Root masses among the three species were not statistically different in 2000. Root masses in the top 2 inches of turf average 0.635 g for creeping bentgrass, 0.844 for supina bluegrass, and 0.938 g for Kentucky bluegrass. In 2001 Kentucky bluegrass had significantly more root mass (0.488 g) than either supina bluegrass (0.215 g) or creeping bentgrass (0.162 g).

DISCUSSION

The data indicating a differential response to N carrier on turf in shaded conditions is important because it has not previously been reported. The information is especially important because it allows us to solve another “piece of the puzzle” in the development of management programs for shaded turf areas. Although the turf in the study was maintained as a “tee box” turf, the fertility treatments should have similar effects at both lower and higher mowing heights (although to be sure these variables should also be investigated). The fact there were no interactions between PGR and N type is also important as this also had not previously been reported for shaded conditions (or full sun for that matter).

We are currently examining the reasons for the differential species response to nitrogen carriers.



EVERYTHING YOU NEED FOR YOUR BEST SEASON EVER!



Toro Reelmaster® 5500-D

A lightweight fairway mower that's agile and easy on turf, yet has the power and cutting capacity to get the toughest jobs done.



Toro Workman® 2100

This is a serious piece of machinery! It's a hard worker that does what you demand of it, and then comes back for more.



Toro 800S® Sprinklers

The Toro 800S® series sprinklers are enhanced with new technology that defines reliability, durability, performance and the lowest cost of ownership.

**For all of these great solutions
and more, contact MTI Distributing,
the Authorized Toro Distributor for
western Wisconsin**

4830 Azelia Avenue N #100
Brooklyn Center, MN 55429
763-592-5600 800-362-3665

www.mtidistributing.com



Kentucky bluegrass, despite its poor turf quality, had substantially greater root mass than either creeping bentgrass or supina bluegrass, while supina bluegrass had greater root mass than creeping bentgrass. Although data were not collected in the fall during establishment, it appears Kentucky bluegrass lost its root system at a slower rate than supina bluegrass or creeping bentgrass. Beard (1973) indicated Kentucky bluegrass has a perennial root system while creeping bentgrass has an annual root system. The slower root turnover (i.e., greater mass) of bluegrass may have allowed it to absorb more N from granular urea compared to the smaller root mass of creeping bentgrass. Kussow (2001) reported that much of the urea applied to turf foliage in liquid form may be lost due to volatilization or trapped in the plant cuticle. In some cases only 40% of the N may actually be absorbed into the plant. The fine leaf texture of creeping bentgrass may have resulted in more efficient capture and absorption of foliar N compared to the coarser-textured Kentucky bluegrass. Since leaf N content in creeping bentgrass was not affected by N type, however, it is likely there were other factors affecting the N response.

CONCLUSION

The best turf for shaded tees can be obtained using supina bluegrass and fertilizing it with granular urea in the spring and early summer, switching to foliar applications of urea in August. If creeping bentgrass is used, it should be fertilized primarily or only with liquid urea applications directed at the foliage. Kentucky bluegrass should generally not be used for tees subject to 80% or more shade.

Literature Cited

- Beard, J.B. 1973. Turfgrass: Science and culture. Prentice-Hall, Englewood Cliffs, NJ.
- Bushoven, J.T., and R.J. Hull. 2001. Nitrogen use efficiency is linked to nitrate reductase activity and biomass partitioning between roots and shoots of perennial ryegrass and creeping bentgrass. *Int. Turf Soc. Res. J.* 9:245-252.
- Goss, R.M., J.H. Baird, S.L. Kelm, and R.N. Calhoun. 2002. Trinexapac-ethyl and nitrogen effects on creeping bentgrass grown under reduced light conditions. *Crop Sci.* 42:472-479.
- Jiang, Z., and R.J. Hull. 1999. Partitioning of nitrate assimilation between shoots and roots of Kentucky bluegrass. *Crop Sci.* 39:746-754.
- Kussow, W.R. 2001. Foliar, liquid,

or granular. *The Grass Roots.* 30(5):43-45.

Spangenberg, B.G., T.W. Fermanian, and D.J. Wehner. 1986. Evaluation of liquid-applied nitrogen fertilizers on Kentucky bluegrass turf. *Agron. J.* 78:1002-1006.

Stier, J.C., and J.N. Rogers, III. 2001. Trinexapac-ethyl and iron effects on supina and Kentucky bluegrasses under low irradiance. *Crop Sci.* 41:457-465.

ACKNOWLEDGEMENTS

The authors wish to thank Dr. Wayne Kussow from the University of Wisconsin-Madison for his technical assistance in the field and for the use of laboratory space and supplies. We also wish to thank the Wisconsin Turfgrass Association, the Golf Course Superintendents Association of America, the Northern Great Lakes Golf Course Superintendents Association, and the College of Agricultural and Life Sciences at the University of Wisconsin-Madison for funding and support of this research.

Authors Note: Kurt Steinke fulfilled the requirements for the Master of Science degree at the University of Wisconsin-Madison in March of 2002 and is now employed by a private turf company in Wisconsin. He did a masterful job during his program, producing sufficient data for three peer-reviewed publications. One manuscript on herbicide tolerance of supina bluegrass will be published during summer of 2002 in the *Journal of Environmental Horticulture*. Two manuscripts on the shade project will be submitted for publication in *Crop Science* during spring and summer of 2002. ♻

ARMOR
ACCESS BOXES™
 FORMERLY AMETEK® ACCESS BOXES
 Manufactured in
 Sheboygan, Wisconsin
 Specify **ARMOR ACCESS BOXES™**
 – A Wisconsin Product!





Nitrogen and Phosphorus Losses from Turf

By Dr. Wayne R. Kussow, Department of Soil Science, University of Wisconsin-Madison

Introduction

A six year study was conducted at the university's O.J. Noer Turfgrass Research and Education Facility to determine the quantities of nitrogen (N) and phosphorus (P) lost from Kentucky bluegrass established on disturbed soil. The basic soil disturbance conditions simulated were subsoil compaction and topsoil layering over the subsoil. These conditions are common in landscaped urban areas and on golf courses. Two remedial actions were tested. One was chisel plowing of the compacted subsoil. The other was tilling of half the topsoil into the subsoil to disrupt the topsoil-subsoil interface.

Methods

The study was established on an 800-foot long slope from which the topsoil in the study area was stripped off, the remaining soil graded with a bulldozer to provide a nearly uniform slope of 6% and the subsoil compacted in selected areas with a vibrating roller before

replacing the silt loam topsoil. The area was then seeded to a 4-way blend of Kentucky bluegrass cultivars. Once grown in, plots measuring 32 x 8 feet were bounded with plastic lawn edging and runoff water and leachate collection devices installed. The number of runoff plots totaled 18, which were grouped into 3 side-by-side blocks. Volume of runoff water was measured throughout the six years and subsamples analyzed for nitrate + ammoniacal N and water soluble orthophosphate. At no time did the runoff water contain measurable amounts of soil sediment. Percolate was collected from April to December each year, the volume measured and subsamples analyzed for nitrate-N after determining that less than 1 % of the N was in the form of ammonium.

The turf was maintained according to university recommendations. Four pounds of N/1,000 ft² were applied each year in 1.0 lb increments during the periods of May 1 to 15, July 1 to 15, September 1 to 15,

COLUMBIA ParCar

Call for a free demonstration!
And to learn of our special WI course offers at
no cost or obligation!



Eagle Sponsor
of the Golf
Course Owners
of Wisconsin.

- Gas & Electric Golf Cars & Utility Vehicles
- New, Used and Reconditioned
- Full Sales and Service
- Additional Cars for Special Events
- Lease Financing Available

SALES & SERVICE

Madison, WI
(608) 249-6600

Reedsburg, WI
(608) 524-8888

(800) 222-4653



Proudly Manufactured By
Wisconsin Craftsmen