

# TABLE OF CONTENTS

THE PRESIDENT'S MESSAGE  
**Wow!** . . . . . 3

GAZING IN THE GRASS  
**Autumn Turf Management for Best Winter Hardiness** . . . . . 6

TDL  
**The Rise and Likely Fall of PCNB** . . . . . 15

WISCONSIN ENTOMOLOGY REPORT  
**Emerald Ash Borer: Too Close for Comfort!** . . . . . 19

GOLF IN THE FLATLANDS  
**The End of the Line** . . . . . 22

NOTES FROM THE NOER FACILITY  
**Education Abounds at Field Day 2006** . . . . . 27

PERSONALITY PROFILE  
**Anchors Away!** . . . . . 33

THE EDITOR'S NOTEBOOK  
**Transition Time** . . . . . 39

FROM ACROSS THE COUNTRY  
**Ways Golf Courses Deal With Bad News** . . . . . 43

WISCONSIN PATHOLOGY REPORT  
**How Phosphonate Fungicides Affect Bentgrass Stress** . . . . 44

BADGER STATE TURF CLIPPINGS  
 . . . . . 48

## ABOUT THE COVER

Portrait artist Jennifer Samerdyke gives us a cover featuring WGCSA President Mike Lyons.

*"I'm afraid there's more to this than appears on the surface."*

— Mildred Augustine Wirt Benson  
 1905-2002



## THE GRASS ROOTS

THE GRASS ROOTS is a bi-monthly publication of the Wisconsin Golf Course Superintendents Association, printed in Waunakee, Wisconsin by Kramer Printing. No part of THE GRASS ROOTS may be reprinted without expressed written permission by the editor.

### EDITOR AND PUBLISHER

Monroe S. Miller  
 Blackhawk Country Club  
 P.O. Box 5129  
 Madison, WI 53705  
 groots@charter.net

### EDITORIAL STAFF AND BUSINESS AFFAIRS

Dave Brandenburg  
 P.O. Box 314  
 Theresa, WI 53091



Front Row: L-R: Brian Zimmerman, Dustin Riley, Mike Lyons and Jeff Millies. Back Row: L-R: David Oberle, Eric Jasin, Chad Harrington, Randy Dupont & Mike Werth.

### 2006-2007 WGCSA OFFICERS AND DIRECTORS

#### PRESIDENT

Mike Lyons  
 Old Hickory CC  
 316 Grove Street  
 Beaver Dam, WI 53916  
 E-mail: mlyons@powerweb.net

#### DIRECTORS

Eric Jasin  
 Lake Arrowhead GC  
 426 St. Andrews Trail  
 Nekoosa, WI 54457  
 E-mail: arrowheadlakes@hotmail.com

#### VICE PRESIDENT

Dustin Riley  
 Oconomowoc GC  
 W35235 Delafield Rd.  
 Oconomowoc, WI 53966  
 E-mail: ogcriley@voyager.net

Randy Dupont  
 North Hills CC  
 7519 Arthur Rd.  
 Hartford, WI 53027  
 E-mail: rmdnhcc@earthlink.net

#### TREASURER

Brian Zimmerman  
 Milwaukee Co. Parks  
 9480 Watertown Plank Rd.  
 Wauwatosa, WI 53226  
 E-mail: bzimmerman@milwcnty.com

Chad Harrington  
 Autumn Ridge GC  
 335 Berge Street  
 Valders, WI 54245  
 E-mail: puter@lakefield.net

#### SECRETARY

Jeff Millies  
 Edgewood GC  
 W238 S9560 Kuzendorf Ct.  
 Big Bend, WI 53103  
 E-mail: jmillies@wlr.com

Michael Werth  
 Monroe CC  
 2639 Hillcrest Ct.  
 Monroe, WI 53566  
 E-mail: werthashot@tds.net

#### PAST PRESIDENT

Marc Davison  
 Green Bay CC  
 2400 Klondike Rd.  
 Green Bay, WI 54311  
 E-mail: mdavison@usxchange.net



Visit our web site at:  
[www.wgcsa.com](http://www.wgcsa.com)




# Wow!!!

By **Mike Lyons**, Golf Course Superintendent, Old Hickory Country Club

I don't like repeating myself but I feel I need to in this case.

Why you should take advantage or participate more at monthly meetings....

I was very fortunate to have won our local qualifier for BASF-sponsored People vs. the Pro at Lake Arrowhead in June. This year's event for those who qualified, and there were a few ways to qualify, was held at Pinehurst, North Carolina — yes, the historical and famed Pinehurst. Twenty-two superintendents and 178 other qualifiers all had a chance to play either PGA Professional Retief Goosen or Gary McCord. Those under 50 years of age got a chance at Retief Goosen and over 50 got Gary McCord. There was also a superintendent tournament. Two lucky superintendents played for \$10,000 for themselves and \$10,000 for the club in BASF products. Second place winner received \$5,000.



Both tournaments were handicap. Trust me, anyone had a chance. With that said, they look very closely at your handicap and adjusted it so you couldn't sand bag.

The whole event was first class. BASF and everyone at Pinehurst took very good care of us. As my mother would say "it's nice to see how the other half lives" and it was nice. I played four rounds of golf in four days. I have not played four days of golf since I was 12 and day four showed it. My wife was treated to a spa treatment and wine tasting - can't go wrong with that one.

BASF had a reception for the superintendents and their guest every night and the morning breakfast buffet was incredible.

If you are a Donald Ross fan or not, this place is a must see. The history is well preserved and the old photos of who's who are incredible. We did not get a

**syngenta**

Use your points for a worthy cause.

When a GreenPartners® member purchases Syngenta products, they have the option to donate their points to worthy causes. If you haven't registered yet, just log on to [www.greenpartners.com](http://www.greenpartners.com) and find out how rewarding GreenPartners can be. For more information, contact Steve Abler at 920-860-6374.

**Eligible organizations include:**

- Wee One Foundation
- Wisconsin Golf Course Superintendents Association
- Wisconsin Landscape Federation
- Wisconsin Nursery Association
- Wisconsin Turfgrass Association


**GreenPartners®**  
Earn Redeem Donate

©2006 Syngenta. Syngenta Professional Products, Greensboro, NC 27419. GreenPartners® and the Syngenta logo are trademarks of a Syngenta Group Company.



**You're either spending too much . . .**

*Too much time. Too much money.* And you aren't getting better brown patch or dollar spot control for your extra investment. So switch to Chipco® 26GT. It knocks down mycelium within 24 hours of application. And it lasts 14 days or more, twice as long as chlorothalonil. Plus, broad-spectrum Chipco 26GT works as a preventive and a curative against dollar spot and brown patch. You can also tank mix 26GT with other fungicides for season-long control. And 26GT will cost you about 32% less than what you'd spend for chlorothalonil.



**. . . or getting short changed.**

Chipco 26GT was originally known as Chipco 26019, but it was so significantly enhanced in 1997 that it demanded a new name. While the active ingredient remains unchanged, 26GT delivers quicker knockdown of brown patch and dollar spot (up to 48 hours faster than 26019) without sacrificing its excellent 14-day residual.

*Save more, and get more. Use 26GT.*



**Kerry Anderson**  
Sales Representative  
(815) 923-1323





chance to play course two where the Open is played but did play two nice tracks. We were treated to course five and eight. Course eight is a Fazio design and played a little harder than five. They all have that Ross feel. I started out playing well and after two days I was only three back at three over. Day three came and needless to say the wheels fell off, started to leak oil, folded like a cheap card table and whatever. All I know is the arms got short and the o.b. stakes looked about as wide as a goal post. It was ugly. I did play with the gentleman who won. He played to a twelve and played Goosen the next day. I also played two rounds with the superintendent who was playing for the \$10,000. As I am writing this I am not sure how things turned out. But he was a scratch player and had to give nine shots and after nine he




was two down. Goosen and McCord were also up by a few after nine. The PGA guys take this seriously. After three years the matches are all tied at three apiece. McCord is one of the guys who lost and believe me he did not want to lose again. All this will be televised on ESPN 2 in October.

The superintendents were well represented at this great event. At the recognition dinner, Steve Mona from the GCSAA was on hand and Dr. Toni Bucci from BASF gave a very thoughtful speech on how valuable we are to the game of golf and to the environment.

Finally, after all that, the topper came at the dinner put on by BASF for all the superintendents. There was a little problem with the drinking water on the second day at Pinehurst and we had to have the dinner at Pine Needles. I believe

that they will be host to next year's Woman's U.S. Open. Not a bad place to be shipped off to for dinner. After dinner we had heard Arnold Palmer was at a reception next door. After dinner, my wife Dawn and I are outside and the King comes walking out. Dawn tells me to go and get a picture with him but I said she stood a better chance at stopping him than I did. So Dawn and a few other ladies ask for pictures and sure enough, he stops and let's all of us get a picture with him. Great man.

As you can tell, it was an incredible time and opportunity. I truly wish every one of us could have this opportunity. I will say it one last time — participate more and some day you could meet truly one of the greats of this game and be treated like "the other half." Take care. ♣





# Autumn Turf Management for Best Winter Hardiness

By Dr. John Stier, Department of Horticulture, University of Wisconsin-Madison

Autumn is an important period for turf to recover from summer stresses of heat, drought, and golf play. Turfgrasses undergo a number of changes, both visible and within the plant, as daylength shortens and temperatures cool in the autumn. These changes are necessary for the plants to cold-harden sufficiently to survive the winter period when photosynthesis is reduced if not stopped altogether. Most of our turfgrasses have inherently good low temperature tolerance. However, they must also survive a host of adversities ranging from winter desiccation to snow mold diseases. Adhering to good agronomic practices during the autumn can help the turf survive winter and enter the spring looking and playing better.

## Mowing

Turfgrasses thrive on being mowed as it stimulates development of new buds and shoot growth. Each grass species has its own optimal mowing height range (Beard,

2002). Golf course turf is often mowed at the lower end, or even below, the optimal mowing height range. For example, creeping bentgrass putting greens cut below 0.25 inch and Kentucky bluegrass fairways cut at 0.5 inch are surviving below their optimal mowing height. During the growing season irrigation, fungicides, and other practices keep the grass alive. A turf weakened by season-long low mowing may not survive if the winter lasts too long, temperatures get too cold, or fungal diseases are too aggressive. Beard (1973) shows a graph of Kentucky bluegrass having less and less cold tolerance as mowing heights were reduced from an optimum of 2 inches to 1.5, 1.0, and 0.5 inches.

Mowing should continue in the autumn until the turf has stopped producing noticeable clippings. The grass may still be growing, but much of the new growth will be in the form of new bud development on the crowns and stolon, rhizome, or root growth. Mowing height should not be decreased during the autumn as the cool nights and warm days are ideal for photosynthesis.

Much like a squirrel or bear, cool-season turfgrasses are busy storing carbohydrates from photosynthesis to use as an energy source during the long cold days of winter. The carbohydrates may also function as an antifreeze, lowering the freezing point of the plant so it can survive low winter temperatures. The mowing height should be raised in cases where the turf has struggled during the summer due to low mowing heights and/or other stresses such as shade. This will increase the amount of photosynthate production and the storage capacity as the shoots and leaves store over 50% of turfgrass carbohydrates (Hull, 1987).

Tree leaves should be removed from turf as they fall to the ground. Coverage of turf from tree leaves will reduce or prevent photosynthesis during the autumn. Extensive leaf cover may also substitute for snow to allow development of gray snow mold. Tree leaves can usually be mulch-mowed into the turf. Research at Michigan State University indicates leaves of some tree-species can reduce weed populations when mulch-mowed into turf (*J. Rogers, pers. communication*).

## Fertilization

A number of experiments show autumn applications of nitrogen can improve autumn and/or spring color (Wilkinson and Duff, 1972; White and Smithberg, 1980; Wehner et al., 1988). However, fertilizer timing and rate can be critical. A number of other experiments show overfertilization throughout the year or fertilizing



## Water Well Solutions

Toll Free 888.769.9009

*Having Provided Professional Services*

*Blackwolf Run  
North Shore Golf Club  
Oconomowoc Golf Club  
Riverview Country Club*



Design, Service, & Installation  
High Capacity Wells  
Well & Booster Pumps (All Brands)  
Same Day Service Response  
Annual Maintenance & Startup Programs  
Well Rehabilitation:  
AirBurst, Aquateam & Hy-Rip

Authorized Dealer



Oconomowoc, WI

Beecher, IL

WWW.WATERWELLSOLUTIONS.COM

during the cold acclimation period reduces low temperature survival of cool-season grasses (Carroll and Welton, 1939; Wilkinson and Duff, 1972; Welterlen and Watschke, 1985). High N rates create more succulent tissues and crowns having a higher moisture content which makes plants more likely to die from freezing injury (Gusta et al., 1975; Tompkins et al., 2000; Webster and Ebdon, 2005).

Nitrogen fertilizer should be applied at rates and times recommended by Dr. Kussow. Nitrogen can be supplied towards the end of the summer golf season as this nitrogen will be useful for turf recovery from summer stresses, development of new buds for the following spring, and stolon/rhizome/root growth. September 1 is usually a good target date in Wisconsin though this may shift a week or two later for the southern part of the state, particularly the Milwaukee area.

Cold acclimation will begin early to late September and nitrogen applied while the foliage is still capable of growth will reduce cold hardiness. A final application of nitrogen, called a dormant application, should be made after foliage has stopped visibly growing (Webster and Ebdon, 2005), or about the time mowers are put away for the year. The turf is still capable of photosynthesizing and will use the nitrogen for proteins, bud development, and root growth.

Root growth can occur at soil temperatures close to freezing (Stuckey, 1941) while most leaf growth stops at approximately 40°F. Rubisco is the main protein which drives photosynthesis: it is also the primary user of nitrogen as it makes up approximately half of the entire protein mass in the plant.

Cold tolerance is enhanced by production of specialized antifreeze proteins which prevent ice from forming in cells (Griffith et al., 2005). Other proteins called lectins protect membranes from damage so cells can survive winter conditions (Hincha et al. 1997). A number of other proteins associated with freezing tolerance are also formed during cold acclimation (Thomashow, 1999).

Water-soluble nitrogen is probably ideal for the dormant application because it can be quickly metabolized by the plant. Slow-release nitrogen sources are not necessarily bad, though most require microbial activity to release the nitrogen and microbial activity is suppressed at low temperatures. Approximately 1 lb N/1000 ft<sup>2</sup> is ideal, though somewhat greater rates may be suitable for slow release forms. More work is needed in this area to accurately identify the effects of nitrogen types and rates. Some research also indicates that using a fertilizer with equal to twice as much potassium to nitrogen enhance cold tolerance (Hurto and Troll, 1980; Webster and Ebdon, 2005).

#### Irrigation

The cold hardening process involves a natural dehydration of plant tissues which lowers the freezing

potential. Mild drought stress increases cold hardiness (Welterlen and Watschke, 1985). Consequently, irrigation should be reduced as turf begins to enter the cold acclimation period and eventually stopped before the beginning of winter dormancy.

Although the acclimation period is not well defined and varies depending on the species, it appears to begin as daytime temperatures decrease to approximately 50°F. In the southern half of Wisconsin cold acclimation will usually begin in mid to late September and somewhat sooner in the north. Exceptions to withholding irrigation should be made only if turf is being established or to help turf recover from summer drought or other stresses. Even in those cases it is wise to reduce irrigation before low temperatures cause turf to stop visible growth in order to allow the turf to cold harden as much as possible.

## BULK TRUCK LOAD

# \$4,995

For Better Results.  
Naturally.

**Milorganite**<sup>®</sup>

www.milorganite.com  
1-800-237-9645

**FERTILIZE 62 ACRES**

FROM THESE AUTHORIZED DEALERS ONLY:

**HORST**  
DISTRIBUTING, INC.

WISCONSIN  
*Turf*

(920) 849-2341 (608) 752-8766

# DHD TREE PRODUCTS

JUNEAU, WI

*"COMMITTED TO YOUR SUCCESS"*

*Stress Phiter*



**WHITE GOLD®**

*Nature  
Safe®*

**Emerald Isle, Ltd.**



**REGAL**  
CHEMICAL COMPANY



**Salsco, INC.**  
LEADER BY DESIGN



Hydraway 2000



**Kwik-Trench®**

**Wedgle Direct-Inject**  
TREE TREATMENT SYSTEM



**866 - 343-8733**

(866 - DHD-TREE)

**DQ MOBILE: 920 - 210-8616**



### Drainage

Poorly drained areas should be renovated during the growing season to prevent development of ice cover during the winter. Ice covers can often cause turf death, especially on closely mown turfs like putting greens composed of annual bluegrass (Tompkins et al., 2004; Minner et al., 2006) for reasons that are not completely understood. Likely causes include lack of oxygen and/or increases in toxic gases such as carbon dioxide or cyanide (Andrews, 1996; Lebeau, 1966; Dionne et al., 2002).

Internal drainage will be irrelevant during the winter as soil will usually be frozen if an ice cover exists (though exceptions can exist). Consequently, provisions need to be made to allow surface drainage as rainfall occurs or ice and snow melt. Areas of standing water which are allowed to freeze will be the most likely to die from ice cover.

### Turf establishment

Areas damaged during the summer may require seeding or sodding if the remaining turf has not filled in by early autumn. Seeding should be done by early to mid-September to give the turf sufficient time to mature before winter. If this is not possible dormant seeding may be necessary.

Dormant seeding is performed late enough in the year that the chance of germination has passed: the seed is expected to germinate early the following spring. Dormant seeding does not have the success rate of late summer seeding as the seed may be killed by low winter temperatures or washed away during winter storms or thaws, but results may be better than waiting until spring.

All seeding requires good seed to soil contact for success, so seeding should be done in conjunction with aeration, topdressing, or raking. Slit-seeding is an excellent method if the equipment is available. It may be better to sod areas expected to have traffic during the autumn as seed will not sufficiently establish.

### Core aeration

Little research has been conducted on the effects of core aeration on winter turf tolerance. Core aeration during the growing season is known to stimulate root growth and result in denser turf. Core aeration in the autumn should also stimulate root growth by providing oxygen to the roots which will continue to grow long after foliage growth has ceased. On putting greens and tees, especially those on sand-based root zones, it will be important to fill holes with topdressing unless the holes close on their own before winter to avoid winter desiccation of turf around the holes.

Historically core aeration in the fall has been favored as it is less likely to interfere with play than summer or spring aerations. Murphy et al. (2005) found seeding bentgrass in September or October resulted in significantly more annual bluegrass than

seeding in July or August. Consequently, fall aeration may increase the amount of annual bluegrass unless core aeration is conducted late enough that seedlings won't survive the winter, e.g., after the average temperature has dropped to below 50°F. More research is needed to determine the effect of timing of core aeration on annual bluegrass populations in Wisconsin.

### Herbicides

Many perennial weeds such as violet and creeping Charlie are best controlled by herbicides applied in the autumn during the cold acclimation period. Daytime temperatures above 50°F give better results than colder temperatures. Presumably the cold acclimation process causes more of the herbicide to be transported to the growing points (meristems) than when applied in the spring or summer. In addition, weeds that survive the herbicide may be so weakened they can't survive winter. Most herbicides cannot be applied to recently seeded areas. If the situation requires autumn application of herbicides along with new turf establishment, the treated area should either be dormant-seeded or sodded.

### Plant growth regulators

Many superintendents use plant growth regulators such as Primo™ during the year to reduce clipping



Midwest  
Turf Products

1444 Bernard Drive • Addison, Illinois 60101

Proud Distributors of:



**Cubby O'Brien**

Cell 630.742.5124

Office 630.268.8871

Fax 630.268.1357

[cubby@chicagolandturf.com](mailto:cubby@chicagolandturf.com)



yields and to improve turf quality. Research at the O.J. Noer facility shows Primo can enhance cold tolerance of cool-season grasses (Steinke and Stier, 2004). The final application of the year needs to be applied before the cold acclimation period (early to mid-September) as later applications can delay spring greenup (*J. Stier, unpublished data*).

### Fungicides

Snow mold diseases are one of, if not the most, important diseases in Wisconsin. Although a number of cultural management practices can affect the severity of the disease under certain conditions, fungicides remain the only truly effective control of snow molds. Without fungicides, closely mown turf can be completely killed by snow mold diseases in Wisconsin. Dr. Jung and his staff at the University of Wisconsin-Madison have passionately studied the various snow mold diseases and published a number of articles on their biology and control (Scheef et al., 2002; Abler and Jung, 2003; Jung et al., 2004; Jo and Jung, 2006). Fungicide controls and strategies continue to change as new products and information become available. Currently one of the best control strategies includes a combination of iprodione, PCNB, and chlorothalonil (Abler and Jung, 2003; Smiley et al., 2005; Vargas, 2005). Fungicides to prevent snow mold should be

applied as late as possible in the year while daytime temperatures are above freezing in order to allow fungicides such as iprodione to penetrate into the leaf tissue for longer-lasting protection.

### Traffic control

Foot and vehicle traffic on turf covered with frost should be avoided. The pressure crushes the leaf tissues and recovery is slow to none due to cool autumn temperatures. If ski or snowmobile paths cross the golf course these should be clearly marked before snowfall. Avoid allowing traffic across high profile areas such as greens, approaches, and tees that will interfere with play during spring repair or be extremely costly to repair. The course should only be open to skiing or snowmobiling when there is at least 3 inches of snow on the ground; lesser amounts can allow extensive turf kill (Eaton and Beard, 1986).

### The final days of autumn

Autumn can be an important time for turf recovery from summer stresses. Using the best management practices can help avoid turf loss during the winter. Keep mowing until the grass stops growing, fertilize before and after the cold acclimation period, and help the grass cold acclimate by reducing irrigation early in the autumn. Get fungicide applications on all critical turf areas before the grass freezes or snow fall occurs.



**We're Part of the Game**



**Premium, Professional Fertilizer.**

Visit us at [springvalleyusa.com](http://springvalleyusa.com) or call us at 1-800-635-2123