

TABLE OF CONTENTS

THE PRESIDENT'S MESSAGE Another Great One Retires	3
GAZING IN THE GRASS New Formulation and Application Strategies for <i>Poa annua</i> Control with Velocity Herbicide	5
WISCONSIN SOILS REPORT NR 151 Update: 'Tis the Season for Soil Testing	13
THE SPORTS PAGE Golf Courses Are Not Killing Us	20
WISCONSIN ENTOMOLOGY REPORT Emerald Ash Borer: Insecticide Options for Protecting Ash Tress and Their Effectiveness	23
TDL Fungicide Resistance in Turfgrass - An Introduction	32
NOTES FROM THE NOER FACILITY The Big Show Will Be Like No Other	36
GOLF IN THE FLATLANDS A Series of Events	39
JOTTINGS FROM THE GOLF COURSE JOURNAL Our Unsung Heroine	44
THE EDITOR'S NOTEBOOK Cruising Into Summer	46
BADGER STATE TURF CLIPPINGS	50

ABOUT THE COVER

The excitement returns as the USGA returns to Wisconsin for one of their championship tournaments. Our cover celebrates the 2007 U.S. SENIOR OPEN at Whistling Straits with Bev Bergemann's colorful rendition of the tournament logo.

*"Every clod feels a stir of might,
An instinct within it that reaches and towers,
And, groping blindly above it for light,
Climbs to a soul in grass and flowers."*

JUNE
By James Russell Lowell



Visit our web site at:
www.wgcsa.com

THE GRASS ROOTS

THE GRASS ROOTS is a bi-monthly publication of the Wisconsin Golf Course Superintendents Association, printed in Wausau, Wisconsin by Kramer Printing. No part of THE GRASS ROOTS may be reprinted without expressed written permission by the editor.

EDITOR AND PUBLISHER

Monroe S. Miller
Blackhawk Country Club
P.O. Box 5129
Madison, WI 53705
groots@charter.net

EDITORIAL STAFF AND BUSINESS AFFAIRS

Dave Brandenburg
P.O. Box 314
Theresa, WI 53091



Front Row: L-R: Matt Schmitz, Brian Zimmerman, Mike Lyons, Dustin Riley, and Jeff Millies.
Back Row: L-R: Chad Harrington, David Swift, Jim VanHerwynen, David Oberle, Marc Davison.

2007-2008 WGCSA OFFICERS AND DIRECTORS

PRESIDENT Mike Lyons Old Hickory CC 316 Grove Street Beaver Dam, WI 53916 E-mail: mike@oldhickorycc.com	DIRECTORS Chad Harrington Autumn Ridge GC 335 Berge Street Valders, WI 54245 E-mail: putter@lakefield.net
VICE PRESIDENT Dustin Riley Oconomowoc GC 762 Summit Avenue Oconomowoc, WI 53066 E-mail: ogcsuper@bizwlrr.com	Matt Schmitz The Bruce Company 8120 Martinsville Road Cross Plains, WI 53528 E-mail: mschmitz@bruceco.com
TREASURER Brian Zimmerman Milwaukee Co. Parks 7350 South 69th Franklin, WI 53132 E-mail: bzimmerman@milwcnty.com	Jim VanHerwynen South Hills Golf and C.C. 1134 South Park Avenue Fond du Lac, WI 54935 E-mail: jkvanherwynen@charter.net
SECRETARY Jeff Millies Edgewood GC W238 S9560 Kuzendorf Ct. Big Bend, WI 53103 E-mail: jmillies@wi.rr.com	David Swift Whistling Straits GC 4106 North 46th Street Sheboygan, WI 53083 E-mail: Swift@Kohler.com
LIAISON David Oberle BASF Corp. 4250 Wexford Way Eagan, MN 55122 E-mail: oberleD@BASF.com	PAST PRESIDENT Marc Davison Green Bay CC 2400 Klondike Rd. Green Bay, WI 54311 E-mail: mdavison@greenbaycc.org



Another Great One Retires

By **Mike Lyons**, Golf Course Superintendent, Old Hickory Country Club

Hi everyone. Hope all is well. As many of you may know, the WSGA office has moved to a new location. However, what you may not know is that one of their truly great employees is retiring. Diane Haas has been with the WSGA for over 25 years. Diane's husband Gene retired as the Executive Director of the WSGA a few years back. Now it is Diane's turn. Diane has been a very key component of our association. Those who have served our association in one way or another know her well. She has been taking care of our membership dues, directory, financials and anything else we could throw at her. Diane was a key element in our early beginnings. She will be truly missed by us all and we wish her the best. Thank you again, Diane.



Like everyone else the grass is growing here in Beaver Dam, dandelions are popping and green, tee and fairway aeration are nearing completion. For the most part, staying ahead of or trying to figure out what Mother Nature will bring us is always a challenge. Getting things completed in a timely fashion is never easy. It is either too wet, too windy or there just is not enough time. But it sure does feel nice to be outside.

Our first couple meetings have been well attended. I hope this trend will continue for the rest of the year. It is never too early to volunteer your course for next season. We are always open to education ideas as well. Hope everyone is ready for the summer months and that they treat you and the turf well. Take care. 🌿



syngenta

A little goes a long way.


Now there's a fairway fungicide that protects against all major turf diseases. What's better, it offers a low 0.75 oz fairway rate for control of dollar spot and brown patch. New Headway™ fungicide's dual mode of action provides complete systemic protection, so the whole course can have the full protection it needs.



Contact Steve Abler at 920-860-6374 to learn more.

www.syngentaprofessionalproducts.com

Important: Always read and follow label instructions before buying or using this product.
©2007 Syngenta. Syngenta Professional Products, Greensboro, NC 27419.
Headway™ and the Syngenta logo are trademarks of a Syngenta Group Company.



**TO MAKE IT
ON THIS TURF,
YOU NEED THE
POWER MORE
PROFESSIONALS
DEMAND.**



Find out why more people who cut grass for a living prefer to do it with KOHLER[®] powered riding equipment. Visit KohlerEngines.com, or call 800-544-2444, ext GR2

KOHLER[®]

New Formulation and Application Strategies for *Poa annua* Control with Velocity Herbicide



By Dr. John Stier, University of Wisconsin-Madison, Department of Horticulture

Back in the 2004 September/October issue of *The Grass Roots* we first introduced research on a new turf herbicide, bispyribac-sodium, to remove annual bluegrass from creeping bentgrass fairways (Stier, 2004). The herbicide has been available to Wisconsin superintendents for the past two years under the trade name Velocity™.

In our 2004 article we demonstrated that an 80% wettable powder (WP) formulation of bispyribac-sodium provided up to 80% annual bluegrass control. A WDG (wetable dry granule) formulation, while much more desirable to handle for mixing/loading

purposes, performed relatively poorly and provided only about 30% control. In 2006 we conducted another battery of tests to evaluate the efficacy of a new SG (soluble granule) formulation. The SG formulation is more desirable from a handling standpoint than a WP as it's less messy to measure and contamination of clothing and surrounding surfaces is virtually nil compared to a WP. In addition, the solubility of the SG formulation means constant agitation is not needed as with WPs. Finally, the SG formulation is less abrasive to nozzles than the WP.

Climatic conditions, receptive-

ness of weeds to be controlled by an herbicide, and spray conditions or technique can often influence product effectiveness. In our 2004 report we found that two or three applications (14 days apart) of bispyribac-sodium at 30 or 45 g ai/A per application provided significantly better control than four applications (14 days apart) of 20 g ai/A per application or 1 or 2 applications of either 45 or 30 g ai/A, respectively. However, sometimes multiple applications at a lower rate can be more effective than a single or fewer applications at a higher rate because more actual product might eventually

syngenta

Use your points for a worthy cause.

When a GreenPartners® member purchases Syngenta products, they have the option to donate their points to worthy causes. If you haven't registered yet, just log on to www.greenpartners.com and find out how rewarding GreenPartners can be. For more information, contact Steve Abler at 920-860-6374.

Eligible organizations include:

- Wee One Foundation
- Wisconsin Golf Course Superintendents Association
- Wisconsin Landscape Federation
- Wisconsin Nursery Association
- Wisconsin Turfgrass Association

GreenPartners
Earn Redeem Donate

©2007 Syngenta. Syngenta Professional Products, Greensboro, NC 27419. GreenPartners® and the Syngenta logo are trademarks of a Syngenta Group Company.

get into the plant. Furthermore, multiple applications of a lower herbicide rate can sometimes be less injurious to desirable grass (e.g., creeping bentgrass) than fewer applications applied at a higher rate.

Our 2004 study also showed bispyribac-sodium was capable of controlling dollar spot disease, a type of activity highly unusual for an herbicide. Enough research has been conducted on this unique aspect of the product that dollar spot control is now listed on the Velocity label.

Our objectives this past year were to determine the effectiveness of an SG formulation, applied at either 10 or 30 g ai/A per application, at either 2 or 14 day intervals, to achieve a total of 60 g ai/A for the year. We also collected data on dollar spot control of the SG.

MATERIALS AND METHODS

We conducted the study at the O.J. Noer Turfgrass Research and Education Facility in Verona, Wisconsin on a mixed stand of creeping bentgrass (*Agrostis stolonifera*) and annual bluegrass (*Poa annua*). Most if not all of the *P. annua* appeared to be the annual biotype though some perennial biotype may have been present. The soil type was a silt loam with pH about 7.5. Established about 1997, the turf has been mown at 1/2" three times per week, receives 3 lbs N/M/year and is not treated with additional fungicide. Irrigation was applied to prevent drought stress.

Six treatments were arranged in a randomized complete block design with four replications (Table 1). Treatments were applied beginning on 8 June, 2006 with a CO₂-powered backpack sprayer, using XR TeeJet 8004 VS nozzles, at 38 PSI, in water carrier to provide 1.5 gallons spray volume/1000 square feet. Bentgrass tolerance, percent *Poa annua* control, and dollar spot

Table 1. Application methods to compare efficacy of 60 g ai/A of bispyribac-sodium for *Poa annua* Control with Velocity Herbicide trial at the O.J. Noer Turfgrass Research and Education Facility in Verona, WI, 2006.

Trt	Product	Rate/Application	Timing	Application Dates
1	Velocity 17.6 SG	10 g ai/A	6 apps 2 day interval	June 8, 10, 12, 14, 16, 19
2	Velocity 17.6 SG	10 g ai/A	6 apps 7 day interval	June 8, 15, 23, 29, July 5, 11
3	Velocity 17.6 SG	30 g ai/A	2 apps 2 day interval	June 8, 10
4	Velocity 17.6 SG	30 g ai/A	2 apps 7 day interval	June 8, 15
5	Velocity 17.6 SG	30 g ai/A	2 apps 14 day interval	June 8, 23
6	Untreated Control	---	---	None

Table 3. Percent *Poa annua* control, Verona, WI, 2006.

Treatment	Jun 11	Jun 16	Jun 23	Jun 29	Jul 11	Jul 26	Aug 8	Sep 9
1	0	3.8	8.8 b	52.5 a	89.3a	86.3 a	90.3 a	77.1 ab
2	0	17.7	3.1 b	11.5 b	50.4 b	77.9 a	94.6 a	87.3 a
3	0	15.8	23.3 a	44.2 a	57.5 b	68.3 a	72.5 bc	58.3 b
4	0	10.8	20.0 a	37.1 a	65.4 b	61.3 a	67.9 c	56.7 bc
5	0	17.3	4.2 b	7.3 b	67.1 ab	67.1 a	83.4 ab	79.4 ab
6	0	0.0	0.0 b	0.0 b	0.0 c	21.9 b	42.7 d	32.3 c
LSD (0.05)	ns	ns	10.1	20.7	23.8	25.9	13.8	24.4

Percent *P. annua* control is presented here as the percentage of the initial *P. annua* in each treatment that was controlled by the treatment application. Means followed by the same on letter within columns are not significantly different at $P \leq 0.05$. The shading emphasizes the best treatments for *P. annua* control.

control were evaluated at approximately 3, 7, 14, 21, 28, 42, and 56 days after initial treatments began. In addition bentgrass quality was rated multiple times throughout the growing season on a scale from 1-9, where 1=dead turf, 9=best quality turf, and 6=acceptable turf quality. Percent *Poa annua* control was determined by visually rating the percentage of living *Poa annua* in the plot on a given rating date then using the number in the following equation to determine percent control: $100 * [1 - (\%poa \text{ on rating date} / \text{initial } \%poa)]$. Bentgrass tolerance was rated on a scale from 1-9, where 1=no injury, 9=dead turf, and 3 or greater = unacceptable injury. Dollar spot control was determined by counting the number of dollar spots on a given plot and comparing to the control plot in the same replication using the following equation: $100 * [1 - (\# \text{ spots on trt. } X / \# \text{ spots on control plot})]$. Soil (2" depth) and air temperatures were recorded at each application date.

RESULTS AND DISCUSSION

Percent *Poa annua* control for each treatment is presented in Table 3. In this study all of the treatments received the same overall rate of Velocity, however, treatments were split into two or six applications at various timings (Table 1). Air temperatures ranged from ideal (20.6°C = 69°F) to supraoptimal (30°C = 86°F) for turf growth during the treatment period. Some research suggests that *P. annua* control may be improved when Velocity is applied at slightly above optimal temperatures because *P. annua* is less heat-tolerant than creeping bentgrass and control will therefore be improved. While temperature at time of application is important to maximize product absorption, the effect of temperatures in the days following application are unknown. Studies are needed to determine the translocation and metabolism of bispyribac-sodium under different temperature regimes in controlled environments such as growth chambers.




Figure 2. Dollar spot control between 16 June and 26 July, Verona, WI 2006. (Scale: 0-100%).

Soil temperatures ranged between 20-26°C (68-79°F) which were above the optimum for cool-season turf growth during the treatment period (ideally 10-18°C = 50-60°F). By eight days after the initial treatment (DAIT) all treatments were providing some control of *P. annua*. At 14 DAIT treatments 3 and 4, which were 2 applications on a 2 and 7 day interval respectively, were providing over 20% *P. annua* control which was greater than all other treatments. At around 21 DAIT, treatment 1, which consisted of 6 applications on a two day interval, had caught up to treatments 3 and 4 in terms of *P. annua* control. On 11 July (42 DAIT), the last of the treatment applications had been made and treatments 1 and 5 were showing the greatest *Poa annua* control. The rating taken on 26 July showed that all Velocity treatments were providing statistically equal control of *Poa annua* and all treatments were significantly better than the control. Control of *Poa annua* in treatment 6 (the untreated control) was due to severe dollar spot disease which

was affecting the *P. annua* more than the creeping bentgrass. The *Poa annua* control ratings between 26 July and 9 September were probably the most meaningful because at this point all treatments had been made and had been given time to take effect. Ultimately, multiple applications of 10 g ai/A at 2 and 7 day intervals gave better than 90% control while two applications of 30 g ai/A at 14 day intervals also gave good control, approximately 80%. *P. annua* populations partially recovered later in the summer when 30 g ai/A treatments were made at 2 and 7 day intervals. The multiple lower rates probably gave better control than higher rates with fewer applications because the higher rates killed more of the leaf tissue quickly, probably reducing the total amount of product to enter the plant over the course of the study. The 30 g ai/A rate was most successful when applied at a 14 day intervals, likely because new leaf growth occurred during the interval which was able to absorb more of the product at the second application. The overall control of

P. annua (highs of 80-90%) was similar to our previous work and mirrors what many other tests have found. What is uncertain is if the remaining 10-20% of the *P. annua* population was resistant to the bispyribac-sodium, or if these were "escapes" which received insufficient chemical because they were growing under an existing turf canopy and/or had large enough crowns that effectively diluted the chemical and allowed some buds to survive, or if the plants were seedlings that germinated after the Velocity application. Ultimately we need to know if the perennial biotypes are more resistant than the annual types, and if annual use results in development of resistant populations.

Beginning June 23, dollar spot pressure was uniform enough to detect differences in terms of dollar spot control between the treatments. All treatments provided some dollar spot control, ranging from approximately 60 to 90% (Fig. 2). There may be an interaction between application rate, frequency, and dollar spot pressure though our test was not designed to provide this information. Dollar spot control decreased within 3-4 weeks after the final application when disease pressure was moderate (June and July). Dollar spot control decreased within one week after the final application of 10 g ai/A (Trt 2) when dollar spot pressure became severe in August (Table 4). Treatment 2, 10 g ai/A applied at 7 day intervals for a 42 day period, provided excellent control (>90%) beginning two to three weeks after the initial treatment and which lasted until 8 August, about 1 week after the final treatment. These data suggest that 10 g ai/A of bispyribac-sodium may substitute for conventional dollar spot fungicides, though control may need to be supplemented with fungicides early in the season if dollar spot

King of the Hill.



JACOBSEN® AR-5™

When it comes to providing superior conditions in primary undulated roughs, nothing compares to the Jacobsen® AR-5™ fine-cut rotary mower. Its advanced lift-arm and caster wheel design plus its narrow 22 inch decks provide **excellent range of motion** for following undulating terrain and delivering superior quality-of-cut without scalping. On-demand weight transfer combined with a parallel-series traction circuit allows the AR-5 to **climb steep hills with ease**. And, its convertible mulching decks **make quick work of debris**. See for yourself why the AR-5 is at the top of the heap – call your local dealer today or visit jacobsen.com for more information.

1.888.922.TURF • jacobsen.com **TEXTRON FINANCIAL**



The Official Turf Equipment Supplier to The PGA of America and
The Exclusive Turf Equipment Supplier to PGA Golf Properties

©2007 Jacobsen, A Textron Company. All rights reserved.

Phone: 1-800-443-1840

Fax: 1-608-752-1530

email: sales@wisconsinturf.com

www.wisconsinturf.com



EQUIPMENT CORP.

Wisconsin Turf Equipment Corp.

P.O. Box 708

Janesville, WI 53548

JACOBSEN

A Textron Company

When Performance Matters.™

Table 4. Percent dollar spot control, Verona, WI, 2006.

Treatment	Jun 11	Jun 16	Jun 23	Jun 29	Jul 11	Jul 26	Aug 8	Ave 6/23-7/26
1	33.3 a	42.6 a	87.1 a	97.0 a	96.4 a	53.4 ab	0	83.5 ab
2	30.0 a	51.7 a	76.2 ab	92.0 a	99.1 a	93.8 a	30.0	90.2 a
3	29.9 a	34.5 a	88.4 a	89.2 a	35.7 a	27.5 bc	20.0	60.2 b
4	4.9 a	44.5 a	57.6 b	99.2 a	59.9 a	32.1 bc	0	62.2 b
5	24.7 a	33.8 a	65.0 ab	68.2 a	93.6 a	58.9 ab	0	71.4 ab
6	0.0 a	0.0 a	0.0 c	0.0 b	0.0 b	0.0 c	0	0.0 c
LSD (0.05)	ns	ns	27.5	32.4	30.2	41.1	ns	23.8

Percent dollar spot control is presented here as percent control in comparison to the level of disease pressure present in the control plot of each replication. Means followed by the same letter within columns are not significantly different at $P \leq 0.05$. Green shading emphasizes the best treatment for dollar spot control.

pressure is severe. Given the public pressure to reduce pesticide use, information on dollar spot management with reduced fungicide input during a Velocity-treatment program would be beneficial to golf course superintendents. If the early spring fungicide application is sufficient to significantly reduce the initial amount of the dollar spot fungus, its possible a Velocity program beginning in June would be sufficient to control dollar spot until late July or early August, resulting in perhaps only 2 to 4 fungicide

applications during the year for dollar spot control. Granted, other diseases may still require fungicides to be used May through July though such applications might only need to be on a case-by-case basis as dollar spot is the most commonly treated disease.

Similar to our 2004 study, we saw some moderate but short-term phytotoxicity to bentgrass (Table 5). A rating of 3 or above was given if there was unacceptable phytotoxicity to the creeping bentgrass present in a given treatment. High rates or low rates at 2 day



CHAMPION GREENS

Aerification Services

Greens • Tees • Fairways
 Deep tine with hollow or solid tines
 18 Greens, \$1200 plus tines used
Travel charges may apply

GOT THATCH?
 Get **SISIS VEEMO**
 Unrivaled Fairway Dethatching
 High Speed • Scalproof
Rental Available

KEITH 920.894.4857
JOHN 920.467.9737
www.championgreens.net



DHD TREE PRODUCTS

JUNEAU, WI

"COMMITTED TO YOUR SUCCESS"

Stress Phiter



WHITE GOLD®

Nature Safe®

Emerald Isle, Ltd.



REGAL
CHEMICAL COMPANY



Salsco, INC.
LEADER BY DESIGN



Hydraway 2000



BANNERMAN



Wedgle Direct-Inject
TREE TREATMENT SYSTEM

866 - 343-8733

(866 - DHD-TREE)

DQ MOBILE: 920 - 210-8616