



“I’ll make dollar spot pay.”



“Even the best turf can fall victim to disease. But with **Honor® Intrinsic™ brand fungicide**, I won’t go down easily. On top of unsurpassed disease control, research shows that **Honor Intrinsic’s plant health benefits** give me a better root system so I can stand up to stresses like drought and moisture events, extreme temperatures, and aerification—better than ever.”

Intrinsic brand fungicides don’t just fight disease; they give turf the resilience to endure stress. Find out more by contacting Randy Lusher, 630.810.1832, randy.lusher@basf.com.



TABLE OF CONTENTS

WISCONSIN PATHOLOGY REPORT

Impact of Winter Covers on Snow Mold Development - Year 2 Update. 4

TURFGRASS DIAGNOSTIC LAB

2012-2013 Snow Mold Fungicide Trial Results 8

BADGER STATE TURF CLIPPINGS

News and Notes From WGCSA Members 16

WGCSA

Couples Outing and Family Weekend. 18

ASSISTANTS AVENUE

A First Tee-rrific Project. 20

WGCSA

Joint Meeting at Stevens Point Country Club 24

USGA GREEN SECTION

In House Projects - Don't Bite Off More Than You Can Chew To Save A Buck Or Two 26

MISCELLANY

Exploring The Green Industry 30

COVER STORY

Wisconsin State Open Visits Tuckaway Country Club. . 32

MISCELLANY

BMW Championship At Conway Farms Golf Club 38

EDITORS NOTEBOOK

Long Days, Short Weeks 42

EVENT SCHEDULE 37

ABOUT THE COVER

The 9th Hole at Tuckaway Country Club, Host of the 2013 Wisconsin State Open.

"We are proud to help the grass grow - for golf."

By Charles Granville Wilson, Agronomist, 1920-2013

Charlie Wilson proclaimed this to the audience upon receiving the 1982 USGA Distinguished Service Award.

THE GRASS ROOTS

is the bi-monthly publication of the Wisconsin Golf Course Superintendents Association. No part of the THE GRASS ROOTS may be used without the expressed written permission of the editor.

EDITOR

David A Brandenburg, CGCS
Rolling Meadows Golf Course
PO 314 Theresa, WI 53091
grassroots@wgcsa.com



2013 WGCSA OFFICERS AND DIRECTORS

Front Row: Chad Harrington, Colin Seaburg, Jim Van Herwynen, Jon Canavan. Back Row: Brett Grams, Jeff Millies, Jeff Barlow, Josh LePine, Jake Schneider, Scott Bushman, Kevin Knudtson. (Not pictured Steve Wasser.)

PRESIDENT

Chad Harrington
Autumn Ridge GC
335 Berge Street
Valders, WI 54245
putter@lakefield.net

DIRECTORS

Scott Bushman
Fox Valley GC
W1759 CTH UU
Kaukauna, WI 54130
foxvalleygolf@netzero.com

VICE PRESIDENT

Jim Van Herwynen, CGCS
South Hills Golf & CC
1175 Fond du Lac Ave.
Fond du Lac, WI 54935
jvkvanherwynen@charter.net

Josh Lepine, CGCS
Maple Bluff CC
500 Kensington Drive
Madison, WI 53704
mrpinier@aol.com

TREASURER

Colin Seaburg
Ozaukee CC
1175 N River Road
Mequon, WI 53092
colin@auzaukeecc.com

Kevin Knudtson
Geneva National GC
2200 Petrie Rd.
Lake Geneva, WI 53125
kknudtson@gnresort.com

SECRETARY

Jon Canavan
Milwaukee County Parks
9480 Watertown Plank Rd
Wauwatosa, WI 53226
jon.canavan@milwcnty.com

Jeff Barlow CGCS
Waupaca CC
1330 Ware St.
Waupaca, WI 54981
jbarlow@solarus.net

VENDOR LIAISON

Steve Wasser
John deere Golf
1700 Peters Road
Kaukauna, WI 54130
swasser@johndeeregolf.com

PAST PRESIDENT

Jeff Millies, CGCS
Edgeweod GC
W238 S9560 Kuzendorf Ct
Big Bend, WI 53103
jmillies@wi.rr.com

Class C Representative

Jake Schneider
Blackhawk CC
3606 Blackhawk Dr.
Madison, WI 53705
Email: jake@blackhawkcc.com

CHAPTER MANAGER

Brett Grams
N9122 Virginia Dr.
Waupaca, WI 54981
bgrams@wgcsa.com

Visit our Website at

www.wgcsa.com

Impact of Winter Covers on Snow Mold Development – Year 2 Update

By Dr. Paul Koch, Department of Pathology, University of Wisconsin - Madison

Due to the cool conditions that persisted for much of this past summer, the UW turf pathology team inoculated and covered many of the disease-research plots to encourage intense disease development. For the dollar spot research trials, the daily process would entail placing an Evergreen® cover down in the afternoon prior to leaving for the day (to increase humidity at night) and removing them upon arrival the next morning (to prevent temperatures getting too hot for dollar spot during the day). This process was repeated daily for over a month to different degrees, and became affectionately (or begrudgingly) known as ‘tucking or untucking the kids.’

While superintendents don’t ‘tuck in their kids’ (i.e. putting greens) during the summer months to increase their disease pressure and challenge themselves as men and women, many superintendents in Wisconsin and beyond do cover their putting greens during the winter months. They don’t do this to really challenge the effectiveness of their snow mold fungicide program, they do this to protect against the various ailments that winter can hurl at fragile putting surfaces. The most common ailments include desiccation (Figure 1) and crown hydration (Figure 2), but toxic gas buildup and low temperature kill can also occur. In addition, winter covers can offer up earlier spring green-up, giving the superintendent a potential (but also potentially risky) leg-up on the fickle spring season.

But are superintendents unintentionally challenging their snow mold fungicide programs when they install these winter covers each fall, and could they be setting themselves up for more severe damage as a result of snow mold than they would have received from the various abiotic forms of winterkill (Figure 3)? That is precisely the question that Dr. Jim Kerns (former UW pathologist, now at North Carolina State University) and myself set out to answer during the fall of 2011. Dr. Kerns delivered a first-year update in the Sept/Oct 2012 issue of *The Grass Roots* (Kerns and Koch, 2012), and this article provides an update from the second and final year of the study.



Figure 1. Windswept areas in winters of relatively low snowfall can dehydrate the plants and cause desiccation injury.



Figure 2. Rapid freezing of free water near the base of the plant can rupture cells in the turfgrass crown, killing the plants. Annual bluegrass is most susceptible to this type of winter injury.

Office: 847.537.2177

www.arthurclesen.com



ARTHUR
CLESEN INC

Quality Products. Quality Service. Quality People.

Fertilizers

Andersons
Grigg's
Lebanon
Redox
Shaw's
UMaxx

Chemicals

Aquatrols
Bayer
Phoenix
Syngenta
SePro
Valent

Seed

Classics
"Blue Tag"
Commercial line

Tree Care

Arborjet

Scott Bartosh, Golf Sales Representative • 262-221-5524 • s.bartosh@arthurclesen.com



John Anderson, Sales Manager • 630-669-0519 • j.anderson@arthurclesen.com

The study was conducted on the chipping green at Antigo Bass Lake Country Club. Host superintendent Dave van Auken provided and installed two types of winter covers; a permeable Evergreen® cover and an impermeable GreenJacket® cover with foam insulation underneath. Fungicide treatments consisted of a non-treated control and a tank-mixture of Interface (4.0 fl oz/1000 ft²) and Triton Flo (0.85 fl oz/1000 ft²) applied at only an early timing, only a late timing, or at half rate and applied at both the early and late timing. The early application for year 2 was made on October 4th, 2012 and the late timing for year 2 was made on October 29th, 2012. Covers were installed the same day as the late fungicide application.

The results from the winter of 2012-2013 provide an interesting comparison to the previous winter of 2011-2012. In 2011-2012, snow mold severity on non-treated plots was increased on plots covered with a GreenJacket® cover but relatively similar between the no cover treatment and the Evergreen® treatment. In 2012-2013, however, snow mold severity was higher under the Evergreen® cover compared to no cover, and was again highest under the Green Jacket® cover (Figure 3). Why did the Evergreen® cover increase snow mold severity relative to no cover in year 2 but not year 1? The answer likely lies in the different environmental conditions present during each winter. The winter of 2011-2012 provided a relatively early and constant snow cover in the Antigo area, whereas the winter of 2012-2013 saw a significant thaw during mid-January that saw snow depth fall to near 0. As a result, in conditions of constant snow cover there was little difference in snow mold severity under no cover or under an Evergreen® cover, but under conditions of less consistent snow cover Evergreen® covers increased snow mold pressure.

Did fungicide timing have an impact on snow mold development under the covers? In year 1, the answer was not really. While spraying at only the early fungicide timing did not provide acceptable snow mold control (an expected result), there were not significant differences in the late fungicide timing or the split early + late fungicide timing. However, under the heaviest disease pressure found under the Green Jacket® cover there was a slight decrease in snow mold under the split application compared to the late timing. Year 2 provided almost an exact replica of year 1, with highest disease breakthrough on the early treatments and no statistical differences between early + late and late (Figure 4). As in year 1, however, slight decreases in snow mold under the Green Jacket® were observed with the split timing treatment compared to the late timing.




Figure 3. Snow mold severity under no cover, Evergreen® cover, or Green Jacket® cover in spring of 2013.





Figure 4. Snow mold severity on fungicide treatment applied as early, late, or early + late under no cover, Evergreen® cover, or Green Jacket® cover in spring of 2013. Fungicide treatment consisted of Interface applied at 4.0 fl oz per 1000 ft² tank-mixed with Triton FLO at 0.85 fl oz per 1000 ft². Fungicides were applied at half rate for each application in split early + late application. Early application as made on October 4th, 2012 and late application was made on October 29th, 2012.

WISCONSIN PATHOLOGY REPORT

The research presented here clearly demonstrates that winter covers, especially Green Jacket®, increase snow mold severity compared to no cover. But the differences in snow mold severity under the Evergreen cover from year 1 to 2 suggest that the degree of impact the covers have will vary based on snow depth and snow cover duration. Those areas where the snow cover is deep and persistent won't see as large a difference between covered greens and non-covered greens, whereas those areas in the southern portion of Wisconsin where snow cover is more shallow and less consistent will be more likely to see dramatic increases in snow mold severity under winter covers.

What are the environmental conditions that these winter covers impact to

increase snow mold severity? Alas, further analysis of the winter environmental data is needed before a final answer to that question can be provided. A final write-up including analysis of the

environmental data will be provided to *The Grass Roots* in 2014 following publication of the study in a scientific journal. Until then, sleep tight this winter little putting greens. 



Acknowledgements:

Special acknowledgement to Antigo Bass Lake CC superintendent Dave van Auken for providing the host site, covers, and installation of the covers. In addition, special thanks to the WGCSA Par 4 Research program for providing the financial support required to conduct this study.

References:

Kerns, J. P., Koch, P. L. 2012. Influence of Winter Covers on Snow Mold Severity: A Summary of Year 1. *The Grass Roots* 41(5): 20-22.

ONE.

AND DONE.

**One early application for
season-long crabgrass
and grub control.**

Season-long crabgrass and grub control in one early application. Shaw's® Turf Food with Acelepryn® and Dimension®

One early shot takes care of crabgrass and grubs for the whole season. So you can move on to other things. To learn more, call (574) 772-6275.

Shaw's is a registered trademark of Knox Fertilizer Company, Inc. Acelepryn is a registered trademark of E.I. DuPont de Nemours and Company, Dimension is a registered trademark of Dow AgroSciences LLC. ©2012 Knox Fertilizer Company, Inc.



KNOX FERTILIZER COMPANY
www.knoxfert.com

PROGRO solutions
ABSOLUTE CONFIDENCE

Brett Ziegler - 847.302.9673 brett@progro-solutions.com

Mike Werth - 608.214.7011 mike@progro-solutions.com

www.progro-solutions.com

2012-2013 Snow Mold Fungicide Trial Results

By **Bruce Schweiger**, Turfgrass Diagnostic Lab Director, O.J. Noer Turfgrass Research and Education Facility and **Dr. Paul Koch**, Department of Pathology, University of Wisconsin - Madison

As the fall fast approaches it is time to begin thinking about Typhula blight (*Typhula ishikariensis* and *Typhula incarnata*) and Microdochium patch (*Microdochium nivale*). We urge you to review the results of past trials on the Turfgrass Diagnostic Lab website (tdl.wisc.edu) to aid in developing your fungicide program for the upcoming winter. Though results dating to 2004 are posted on the website, I will focus on the most recent results from the winter of 2012-2013.

Disease was low in most of the trials last winter, and in fact was completely absent in some trials. This was surprising to many, as the duration of snow cover well exceeded 60 days in many parts of the state and would seem to encourage snow mold development. In hindsight the key to last year's lack of snow mold can be directly attributed to the lack of a suitable environment. In many areas snow cover came in mid-December and remained until April or later. But a significant thaw in January left many areas of the state with a shallow cover of snow and ice. Under this scenario the ground had little insulation from the harsh temperatures of winter. Lack of snow cover did not insulate the surface of the soil, allowing the soil to freeze. We know that the fungi that cause snow mold thrive in cold conditions, but not in below-freezing temperatures. The insulating effect of snow cover is what allows snow mold fungi to cause disease, and this past winter our surface temperatures were too cold and unfavorable for widespread disease development.

The trials at Odana Hills Golf Course in Madison and Sentry World Golf Course in Stevens Point did not provide a significant test of the fungicide treatments. In both of these trials nearly all the treatments provided acceptable control of snow mold. One interesting aspect of the SentryWorld trials was the differentiation of color after snow melted due to the applications of turf colorants the previous fall. With a wide array of effective snow mold fungicides to choose from, superintendents are increasingly interested in characteristics aside from merely excellent disease control. Spring color has been one of these characteristics that has garnered significant interest in recent years, and at last year's Snow Mold Field Day there was a fair amount of discussion as to why this could be an important tool for golf course superintendents (Figure 1). It may not improve control of Snow Mold but there are other benefits. The discussion focused on what color was most desired by superintendents and how the early spring golfer would see the greener turf in early spring and how this could potentially bring turf health benefits, increased play, or both.



Figure 1: Plots at Sentry World. Notice the green plots from the Fungicide additive or colorant added to the spray



HARTMAN
Golf & Sports Turf
Renovation & Restoration

8099 Bavaria Rd.
Victoria, MN 55386
P - 952.443.2990
www.hartmancompanies.com

THE MOWER THEY DON'T WANT YOU TO SEE.



The competition doesn't want you to see the Jacobsen ECLIPSE® 322 riding greens mower. They can't match its **ZERO leak points**. They can't provide **individual reel control** to vary cut patterns. They can't offer **programmable frequency-of-clip** from an LED screen. And they can't get anywhere near the **true hybrid fuel savings** or **legendary quality-of-cut**. When it comes to the ECLIPSE® 322, the competition can't do *anything* except hope you don't see it. I guess we just ruined that for them, too.



Phone: 800-279-2341
Fax: 920-849-9576

www.horstdistributing.com
HORST DISTRIBUTING, INC.
CHILTON, WI

JACOBSEN

1.888.922.TURF | www.jacobsen.com

TURFGRASS DIAGNOSTIC LAB



**Figure 2: Wawanowin 10th Fairway
May 8, 2013**

Higher disease pressure at Wawanowin Country Club in Champion, MI provided a slightly stiffer test of the fungicide treatments. The snow mold pressure was higher there than at the other trials but did not provide the level of disease we are accustomed to at that site (Figure 2). Of the 56 treatments 42 allowed less than 5% disease. All these treatments provided acceptable provided disease control and turf quality.

Of these treatments, 23 provided complete control. Of all the treatments most of them contained 3 active ingredients. The full results can be seen in the adjoining tables, and we encourage you to look over the results and compare with results from previous years.

We want to thank all of our cooperators from last years trials for all their assistance and allowing us to use their facility; Jeff Jushka at Odana Hills GC, Gary Tanko at SentryWorld GC, Andy Hakaarinen at Wawanowin CC Brent Belanger at Les Bolstad GC, and Matt Mckinnon at Cragun's Resort. Without their generous support we would not be able to provide you with the independent research you need to make informed decisions.

Snow is coming and our schedules will be slowing down. Take the time to reconnect with family and friends and continue your turfgrass education. If you have any questions regarding the trials or about snow mold in general, please don't hesitate to contact Bruce (bschweiger@wisc.edu) or Paul (plkoch@wisc.edu).



When you use the best, it shows.

With a portfolio of products unmatched in the industry, Syngenta is a necessity for every great business. From herbicides to fungicides to growth regulators, we have everything you need to maintain healthy turf.

Contact Phil Spitz to learn more about Syngenta products.
Cell: 414-429-2015 philip.spitz@syngenta.com

syngenta.

www.greencastonline.com

©2011 Syngenta Crop Protection, LLC, 410 Being Road, Greensboro, NC 27409. Important: Always read and follow label instructions before buying or using Syngenta products. The label contains important conditions of sale, including limitations of remedy and warranty. Meridian®, Renown®, and Tenacity® are not currently registered for use in all states. Please check with your state or local extension service prior to buying or using these products. Scimitar® GC is a Restricted Use Pesticide. AWARD®, Banner MAXX®, Barricade®, Concert®, Daron®, LUSTRO®, Rapra Weather Staff®, Cobalt® Golf®, Depature®, Falcob®®, Freedom®, Heritage®, Instrate®, Medallion®, Medallion®, Monument®, Renown®, SARGOLUX®, Primo MAXX®, Precept® Liquid®, Renown®, Revard®, Scimitar® GC, Subdue MAXX®, Tenacity®, Trinell®, and the Syngenta logo are trademarks of a Syngenta Group Company.

