## **IENTIFIC & PRACTICAL** PRACTICAL

Dean Sime Golf Course Superintendent Interlachen Country Club

On July 13, 1932 Dr. A. A. Granovsky inspected a grub infestation on a Minneapolis golf course. This grub had caused considerable damage by killing the grass here and there on greens and fairways. Diggings in one six inch square yielded 29 grubs, 8 pupae and one freshly emerged adult. The name of the beetle, Ataenius spretulus.

Thus begins our story about a little monster who has been lying in the weeds for forty some years and not gaining much recognition until he became known as the Cincinatle flash. In 1975 he ravaged a number of golf courses in Ohio and had we recognized the symptoms, the same could be said for Minnesota. And what's his favorite meal? Why, of course, a nice succulent Poa annua fairway.

Generally, you will first become aware that you have a problem when turf begins to wilt in rather general areas. It will become noticeable in July when weather problems are generally at their worst. Eventually, the turf dies and you continue to be disgruntled with that "failure" grass.

Having attended various conferences, you begin to think of the possibility of Ataenius and desperately go about tearing at the dead turf. It lifts easily as if there were no roots, but upon digging further you do not discover anything really pertinent as to the cause. So again, the obvious, "failure" grass under stress. The real irony about this is that when this state is reached, the Ataenius has developed into the pupae stage and has buried itself deep in the soil. You may look all day and find only a few beetles in the Larvae stage but had you looked three weeks earlier, the cause of death could have easily been determined.

The beetle will remain in the pupae stage until mid August when it will emerge as an adult. In more southern areas it is possible to have two life cycles per year. However, in Minnesota the second cycle is not evident. It over winters only as an adult in leaves and underbrush adjacent to the fairways. In Spring it begins flying again and may be attracted to lights. In mid May it begins to deposit eggs in your favorite turf and again, its life cycle proceeds.

Dr. Harry Niemczyk, at Ohio State University, began studying this creature in 1975. Little was known prior to that time, primarily, I believe, because the cause of death had simply been attributed to failure grass. Dr. Niemczyk has determined that the most feasible means of control is to go after the adult beetle in the egg laying stages. This, in Minnesota, is early May, especially the week that Spirea and the Black Locust are in full bloom. Using Diazinon at a rate of 6 lbs. active ingredient per acre has shown very promising results. This was done at Interlachen in 1977 with the supervision of Dr. Niemczyk and our local Extension Entomologist, Mark Ascerno. I was encouraged as essentially no Larvae were found on treated fairways and a less significant number were found on two check fairways. Damage was evident on the check fairways but not nearly as severe as the preceding season. I should also report that some tip burn did occur using the liquid formulation. When using this product, water should be applied to wash it from the leaf surface.

Continued on Page 10

Continued from Page 8

To those of you who are not sure if you have this problem, ask yourself what happened on your course last season. If you had problems with Poa annua wilting and dying, purchase a lighted insect trap such as those used for flies and other insects. Place it somewhere out on the course where it is feasible and monitor the results every morning. This should begin in the first week of May and continue for about three weeks. If you have a significant number of Aetenius, call Mark Ascerno at 373-1059 for further information. Remember timing is extremely important when attempting to control the adult stage.

Some work has been done in controlling the beetle in the Larvae stage. A product called Mocap has proven to be most effective. This is an extremely volatile product and protective gear must be used in its application. Also your course must be closed to play the day of application. I would suggest its use only in what is deemed an extreme emergency. Other products used for control in the Larvae stage become bound in the thatch layer and the results are mediocre.

In summary, first determine if you have a problem. I tend to believe that since this creature has been around for many years that its population tends to cycle with the extreme winter conditions. I also feel that courses surrounded by heavily wooded and natural areas may have more problems due to the winter protection these areas offer. If you determine you have a problem, call Mark Ascerno for positive identification and up to date control recommendations.

