

not kill the branch. However, after several years of needle infection and needle loss the branch may die on its own.

To control *Rhizosphaera* needlecast, apply chlorothalonil or bordeaux mixture when the new needles are 1/2 to 2 inches long and again in three to four weeks. To minimize the impact of adverse environmental conditions on spruce trees, keep the trees well watered during dry periods and do not plant young trees too close together.

Note: An excellent color brochure on "How to Identify and Control Rhizosphaera Needlecast" can be obtained by writing to: North central Forest Experiment Station, 1992 Folwell Avenue, St. Paul, MN 55108.

DEER TICK / LYME DISEASE SEASON DRAWS NEAR

by **Jeffrey D. Hahn**
Extension Educator, Entomology
Minnesota Extension Service

Memorial weekend marks the beginning of the high risk season for Lyme disease which is vectored by the deer tick. Although infected ticks are present from April through November, most Lyme disease reports come in June or July.

The deer tick is small with a brownish body and black legs. The tick is found in tall grass and underbrush and can be readily picked up by campers and hunters. The adults feed from March to June and August to November, while the nymphs feed from June to August.

Avoid heavily wooded areas when you can. If this is not possible, take these appropriate safety precautions. Wear protective clothing, such as long-sleeved shirts and pants tucked into heavy socks. Apply repellents to clothing for further protection. A new, effective repellent, called Permanone will be available to homeowners this summer. Check yourself carefully after returning from areas when deer ticks may be present. Nymphal deer ticks are very small and can be difficult to detect.

See a doctor immediately if you believe that you have been bitten by a tick carrying Lyme disease. The first sign of Lyme disease is a red skin lesion accompanied by flu-like symptoms, although there are other indications. Save the tick that bit you as an expert can identify it. correct identification of the tick is very important for a proper diagnosis of lyme disease.

TREE FERTILIZATION

by **Bert T. Swanson and Carl Rosen**
Horticulture Science and Landscape Architecture
Minnesota Extension Service

The following article was excerpted from a newly revised Extension Folder entitled "Tree Fertilization" AG-FO-2421 by Bert T. Swanson and Carl Rosen. This publication is now available from your local County Extension Office.

Most trees in Minnesota have a single flush of growth in the spring and spring is the time when trees have the greatest need for nutrients. Early spring, consequently, is the time when nutrients must be available. Fall application is the easiest and probably the most effective, because the ground is easier to work and the nutrients will be available to the tree very early in the spring when growth begins. Fertilizer may be applied from late September until about mid-November. Spring applications may be made as soon as the ground is workable until late April or early May. if soil conditions are extremely dry, water the soil prior to and after fertilization. Nitrogen should be applied to sandy soils only in the spring or much of it can be leached out in the late fall and early spring.

If a tree is showing symptoms of deficiency, fertilizer may be applied at any time during the growing season to correct the problem. Care must be taken, however, to provide sufficient water for absorption of the nutrients by the plant and prevent fertilizer burn of the roots. During periods of hot, dry weather, two to three inches of water should be applied every two to three weeks to wet the top 12 to 18 inches of an average soil. Heavy clay soils require more water at less frequent intervals while light, sandy soils require less water at more frequent intervals. Do not apply fertilizer in late August as plants may force a new flush of growth in early September. Likewise do not allow plants to go into the winter under a nutrient stress as this will also increase winter injury.

Fertilizers require moisture and oxygen to dissolve and be absorbed by the plant. If excess moisture or lack of oxygen exists, nutrient uptake cannot take place even with adequate nutrients available. Continued fertilization under such conditions will result in excess fertilizer levels. Then as the soil dries or becomes aerated, excess uptake may occur. Excess uptake will stimulate excessive succulent growth that is structurally weak, less likely to produce flowers, and more susceptible to disease and insects, such as fire blight or aphids. The high soluble salt concentrations of excessive fertilizer may also damage the tree causing root or leaf burn. Newly planted trees generally should be fertilized at planting time, providing that certain precautions are followed. Fertilization at this time allows deep placement of phosphorus and potassium. Because these nutrients do not move readily in the